



www.cdc.gov/hepatitis

October 4, 2006

Hepatitis A Fact Sheet

SIGNS & SYMPTOMS	Adults will have signs and symptoms more often than children.	
	<ul style="list-style-type: none"> ● jaundice ● fatigue ● abdominal pain ● loss of appetite 	<ul style="list-style-type: none"> ● nausea ● diarrhea ● fever
CAUSE	<ul style="list-style-type: none"> ● Hepatitis A virus (HAV) 	
LONG-TERM EFFECTS	<ul style="list-style-type: none"> ● There is no chronic (long-term) infection. ● Once you have had hepatitis A you cannot get it again. ● About 15% of people infected with HAV will have prolonged or relapsing symptoms over a 6-9 month period. 	
TRANSMISSION	<ul style="list-style-type: none"> ● HAV is found in the stool (feces) of persons with hepatitis A. ● HAV is usually spread from person to person by putting something in the mouth (even though it may look clean) that has been contaminated with the stool of a person with hepatitis A. 	
PERSONS AT RISK OF INFECTION	<ul style="list-style-type: none"> ● Household contacts of infected persons ● Sex contacts of infected persons ● Persons, especially children, living in areas with increased rates of hepatitis A during the baseline period from 1987-1997. (view map) ● Persons traveling to countries where hepatitis A is common ● Click here for more information about traveler's health ● Men who have sex with men ● Injecting and non-injecting drug users 	

visited on 5/3/2007

PREVENTION	<ul style="list-style-type: none">● Hepatitis A vaccine is the best protection.● Short-term protection against hepatitis A is available from immune globulin. It can be given before and within 2 weeks after coming in contact with HAV.● Always wash your hands with soap and water after using the bathroom, changing a diaper, and before preparing and eating food.
VACCINE RECOMMENDATIONS	<p>Vaccine is recommended for the following persons from 12 months of age and older:</p> <ul style="list-style-type: none">● Travelers to areas with increased rates of hepatitis A Click here for more information about traveler's health● Men who have sex with men● Injecting and non-injecting drug users● Persons with clotting-factor disorders (e.g. hemophilia)● Persons with chronic liver disease● Children living in areas with increased rates of hepatitis A during the baseline period from 1987-1997. (view map)
TRENDS & STATISTICS	<ul style="list-style-type: none">● Occurs in epidemics both nationwide and in communities● During epidemic years, the number of reported cases reached 35,000.● In the late 1990s, hepatitis A vaccine was more widely used and the number of cases reached historic lows.● One-third of Americans have evidence of past infection (immunity).● Click here for hepatitis A disease burden information