

visited 7/25/2008

[Skip navigation](#)[Search MedlinePlus](#)[About MedlinePlus](#) | [Site Map](#) | [FAQs](#) | [Contact Us](#)[Home](#) [Health Topics](#) [Drugs & Supplements](#) [Encyclopedia](#) [Dictionary](#) [News](#) [Directories](#) [Other Resources](#)[español](#)

Medical Encyclopedia

Other encyclopedia topics: [A-Ag](#) [Ah-Ap](#) [Aq-Az](#) [B-Bk](#) [Bl-Bz](#) [C-Cg](#) [Ch-Co](#) [Cp-Cz](#) [D-Di](#) [Dj-Dz](#) [E-Ep](#) [Eq-Ez](#) [F](#) [G](#) [H-Hf](#) [Hg-Hz](#) [I-In](#) [Ii-Iz](#) [J](#) [K](#) [L-Ln](#) [Lo-Lz](#) [M-Mf](#) [Mg-Mz](#) [N](#) [O](#) [P-Pl](#) [Pm-Pz](#) [Q](#) [R](#) [S-Sh](#) [Si-Sp](#) [Sq-Sz](#) [T-Tn](#) [To-Tz](#) [U](#) [V](#) [W](#) [X](#) [Y](#) [Z](#) [0-9](#)


Peripheral neuropathy

 [Printer-friendly version](#)  [E-mail to a friend](#)


Contents of this page:

- [Illustrations](#)
- [Alternative Names](#)
- [Definition](#)
- [Causes](#)
- [Symptoms](#)
- [Exams and Tests](#)
- [Treatment](#)
- [Support Groups](#)
- [Outlook \(Prognosis\)](#)
- [Possible Complications](#)
- [When to Contact a Medical Professional](#)
- [Prevention](#)
- [References](#)

Illustrations



[Nervous system](#)



[Central nervous system](#)

Alternative Names [Return to top](#)

Peripheral neuritis; Neuropathy - peripheral; Neuritis - peripheral

Definition [Return to top](#)

Peripheral neuropathy is a problem with the nerves that carry information to and from the brain and spinal cord. This produces pain, loss of sensation, and inability to control muscles.

visited 7/25/2008

- "Peripheral" means nerves beyond the brain and spinal cord.
- "Neuro" means nerves.
- "Pathy" means abnormal.

Causes [Return to top](#)

The peripheral nerves relay information from your central nervous system (brain and spinal cord) to muscles and other organs and from your skin, joints, and other organs back to your brain. Peripheral neuropathy occurs when these nerves fail to function properly, resulting in pain, loss of sensation, or inability to control muscles.

In some cases, the failure of nerves that control blood vessels, intestines, and other organs results in abnormal blood pressure, digestion problems, and loss of other basic body processes. Peripheral neuropathy may involve damage to a single nerve or nerve group ([mononeuropathy](#)) or may affect multiple nerves (polyneuropathy).

There are numerous reasons for nerves to malfunction. In some cases, no cause can be identified. Damage to nerves can result from one of the specific conditions associated with neuropathy, including:

- Hereditary disorders
 - Charcot-Marie-Tooth disease
 - Friedreich's ataxia
- Systemic or metabolic disorders
 - [Diabetes \(diabetic neuropathy\)](#)
 - Dietary deficiencies (especially vitamin B-12)
 - Excessive alcohol use ([alcoholic neuropathy](#))
 - [Uremia](#) (from [kidney failure](#))
 - Cancer
- Infectious or inflammatory conditions
 - [AIDS](#)
 - Hepatitis
 - [Colorado tick fever](#)
 - [Diphtheria](#)
 - [Guillain-Barre syndrome](#)
 - [HIV infection](#) without development of AIDS
 - [Leprosy](#)
 - Lyme disease
 - [Polyarteritis nodosa](#)
 - [Rheumatoid arthritis](#)
 - [Sarcoidosis](#)
 - [Sjogren syndrome](#)
 - [Syphilis](#)
 - [Systemic lupus erythematosus](#)
 - [Amyloidosis](#)
- Exposure to toxic compounds
 - Sniffing glue or other toxic compounds
 - Nitrous oxide

visited 7/25/2008

- Industrial agents -- especially solvents
- Heavy metals (lead, arsenic, [mercury](#), etc.)
- [Neuropathy secondary to drugs](#) (many drugs can cause neuropathy)
- Miscellaneous causes
 - Ischemia (decreased oxygen and blood flow)
 - Prolonged exposure to cold temperature

Peripheral neuropathy is very common. Because there are numerous types and causes of neuropathy and scientists don't always agree on the same definition of neuropathy, the exact incidence cannot be determined precisely.

Some people have a hereditary predisposition for neuropathy.

Prolonged pressure on a nerve is another risk for developing a nerve injury. Pressure injury may be caused by prolonged immobility (such as a long surgical procedure or lengthy illness) or compression of a nerve by [casts](#), [splints](#), braces, crutches, or other devices.

Symptoms [Return to top](#)

The symptoms depend on which type of nerve is affected. The three main types of nerves are sensory, motor, and autonomic. Neuropathy can affect any one or a combination of all three types of nerves. Symptoms also depend on whether the condition affects the whole body or just one nerve (as from an injury).

SENSATION CHANGES

Damage to sensory fibers results in changes in sensation, burning sensations, [nerve pain](#), [tingling or numbness](#), or an inability to determine joint position, which causes incoordination.

For many neuropathies, sensation changes often begin in the feet and progress toward the center of the body with involvement of other areas as the condition worsens.

MOVEMENT DIFFICULTIES

Damage to the motor fibers interferes with muscle control and can cause weakness, loss of muscle bulk, and loss of dexterity. Sometimes, cramps are a sign of motor nerve involvement.

Other muscle-related symptoms include:

- [Lack of muscle control](#)
- Difficulty or inability to move a part of the body ([paralysis](#))
- [Muscle atrophy](#)
- Muscle twitching (fasciculation) or cramping
- Difficulty breathing or swallowing
- Falling (from legs buckling or tripping over toes)
- Lack of dexterity (such as being unable to button a shirt)

AUTONOMIC SYMPTOMS

visited 7/25/2008

The autonomic nerves control involuntary or semi-voluntary functions, such as control of internal organs and blood pressure. Damage to autonomic nerves can cause:

- [Blurred vision](#)
- [Decreased ability to sweat](#)
- [Dizziness](#) that occurs when standing up or [fainting](#) associated with a fall in blood pressure
- [Heat intolerance](#) with exertion (decreased ability to regulate body temperature)
- Nausea or vomiting after meals
- [Abdominal bloating](#) (swelling)
- Feeling full after eating a small amount ([early satiety](#))
- Diarrhea
- Constipation
- Unintentional weight loss (more than 5% of body weight)
- [Urinary incontinence](#)
- Feeling of incomplete bladder emptying
- Difficulty beginning to urinate ([urinary hesitancy](#))
- [Male impotence](#)

Exams and Tests [Return to top](#)

A detailed history will be needed to determine the cause of the neuropathy. Neurologic examination may reveal abnormalities of movement, sensation, or organ function. (See also entries on the specific nerve dysfunction.) Changes in reflexes and muscle bulk may also be present.

Tests that reveal neuropathy may include:

- [EMG](#) (a recording of electrical activity in muscles)
- [Nerve conduction tests](#)
- [Nerve biopsy](#)
- Blood tests to screen for medical conditions, such as diabetes and vitamin deficiency, among others.

Tests for neuropathy are guided by the suspected cause of the disorder, as suggested by the history, symptoms, and pattern of symptom development. They may include various blood tests, x-rays, scans, or other tests and procedures.

Treatment [Return to top](#)

The first steps of treatment are to identify and treat the underlying medical problem (such as diabetes) or remove the cause (such as alcohol). Other goals include controlling symptoms, curing the disorder if possible, and helping the patient gain maximum independence and self-care ability.

Physical therapy, occupational therapy, and orthopedic interventions may be recommended. For example, exercises and retraining may be used to increase muscle strength and control. Wheelchairs, braces, and splints may improve mobility or the ability to use an affected arm or leg.

Safety is an important consideration for people with neuropathy. Lack of muscle control and reduced sensation increase the risk of falls and other injuries. The person may not notice a potential source of injury because he or

visited 7/25/2008

she can't feel it. For this reason, people with decreased sensation should check their feet or other affected areas frequently for bruises, open skin areas, or other injuries, which may go unnoticed (because there is no pain) and become severely infected. Often, a podiatrist can determine if special orthotic devices are needed.

Safety measures for people experiencing difficulty with movement may include railings, various appliances, removing obstacles such as loose rugs, and other measures as appropriate. Safety measures for people having difficulty with sensation include adequate lighting (including lights left on at night), testing water temperature before bathing, use of protective shoes (no open toes, no high heels, and so on) and similar measures. Shoes should be checked often for grit or rough spots that may cause injury to the feet.

People with neuropathy (especially those with polyneuropathy or [mononeuropathy multiplex](#)) are prone to new nerve injury at pressure points (knees and elbows, for example). They should avoid prolonged pressure on these areas from leaning on the elbows, crossing the knees, or assuming similar positions.

Over-the-counter or prescription pain medications may be needed to control [nerve pain](#). Anticonvulsants (phenytoin, carbamazepine, gabapentin, and pregabalin), tricyclic antidepressants (duloxetine), or other medications may be used to reduce the stabbing pains that some people experience. Use the lowest dose possible to avoid side effects.

Adjusting position, using frames to keep bedclothes off tender body parts, or other measures may also be helpful to reduce pain.

The symptoms of autonomic changes will be treated. However, they may be difficult to treat or respond poorly to treatment.

- Postural hypotension (low blood pressure) -- use of elastic stockings and sleeping with the head elevated may help. Fludrocortisone or similar medications may be beneficial in reducing postural hypotension for some people.
- Reduced gastric motility -- medications that increase gastric motility (such as metoclopramide), eating small frequent meals, sleeping with the head elevated, or other measures may help.
- Bladder dysfunction -- manual expression of urine (pressing over the bladder with the hands), intermittent catheterization, or medications such as bethanechol may be necessary.
- Impotence, diarrhea, constipation or other symptoms are treated as appropriate.

Support Groups [Return to top](#)

Additional information can be obtained from the [Neuropathy Association](#).

Outlook (Prognosis) [Return to top](#)

The outcome greatly depends on the cause of the neuropathy. In cases where a medical condition can be identified and treated, the outlook may be excellent. However, in severe neuropathy, nerve damage can be permanent, even if the cause is treated appropriately.

For most hereditary neuropathies, there is no cure. Some of these conditions are harmless, while others progress more rapidly and may lead to permanent, severe complications.

Possible Complications [Return to top](#)

The inability to feel or notice injuries can lead to infection or structural damage. Changes include poor healing,

visited 7/25/2008

[loss of tissue mass](#), tissue erosions, scarring, and deformity. Other complications include:

- Partial or complete [loss of movement](#) (or control of movement)
- Partial or complete [loss of sensation](#)
- [Difficulty breathing](#)
- [Difficulty swallowing](#)
- Cardiac [arrhythmias](#) (uncommon)
- Decreased self esteem
- Relationship problems related to impotence
- Recurrent or unnoticed injury to any part of the body

When to Contact a Medical Professional [Return to top](#)

Call your health care provider if symptoms of peripheral neuropathy are present. In all cases, early diagnosis and treatment increases the possibility that symptoms can be controlled.

Nerve pain, such as that caused by peripheral neuropathy, can be difficult to control. If pain is severe, contact a pain specialist to make sure you get the best and most up-to-date pain treatment.

Emergency symptoms include irregular or rapid heartbeats, difficulty breathing, difficulty swallowing, and fainting.

Prevention [Return to top](#)

If a prolonged procedure or immobility is expected, appropriate measures (such as padding vulnerable areas) can be taken beforehand to reduce the risk of nerve problems.

Some people have a hereditary predisposition for neuropathy. Such people need to be especially careful to limit alcohol and manage other medical problems closely.

All people can reduce the risk of neuropathy through a balanced diet, drinking alcohol in moderation, and maintaining good control of diabetes and other medical problems, if present.

References [Return to top](#)

Delalande S, de Seze J, Fauchais AL, et al. Neurologic manifestations in primary Sjogren syndrome: a study of 82 patients. *Medicine* (Baltimore). 2004 Sep;83(5):280-91.

Mori K, Iijima M, Sugiura M. Sjogren's syndrome associated painful sensory neuropathy without sensory ataxia. *J Neurol Neurosurg Psychiatry*. 2003 Sep;74(9):1320-2.

Update Date: 8/7/2006

Updated by: Daniel Kantor, M.D., Director of the Comprehensive MS Center, Neuroscience Institute, University of Florida Health Science Center, Jacksonville, FL. Review provided by VeriMed Healthcare Network.



A.D.A.M., Inc. is accredited by URAC, also known as the American Accreditation HealthCare Commission (www.uran.org). URAC's [accreditation program](#) is the first of its kind, requiring compliance with 53 standards of quality and accountability, verified by independent audit. A.D.A.M. is among the first to achieve this important

visited 7/25/2008



distinction for online health information and services. Learn more about A.D.A.M.'s [editorial process](#). A.D.A.M. is also a founding member of Hi-Ethics (www.hiethics.com) and subscribes to the principles of the Health on the Net Foundation (www.hon.ch).

The information provided herein should not be used during any medical emergency or for the diagnosis or treatment of any medical condition. A licensed physician should be consulted for diagnosis and treatment of any and all medical conditions. Call 911 for all medical emergencies. Links to other sites are provided for information only -- they do not constitute endorsements of those other sites. Copyright 1997-2008, A.D.A.M., Inc. Any duplication or distribution of the information contained herein is strictly prohibited.

[Home](#) | [Health Topics](#) | [Drugs & Supplements](#) | [Encyclopedia](#) | [Dictionary](#) | [News](#) | [Directories](#) | [Other Resources](#)

[Copyright](#) | [Privacy](#) | [Accessibility](#) | [Quality Guidelines](#)

[U.S. National Library of Medicine](#), 8600 Rockville Pike, Bethesda, MD 20894

[National Institutes of Health](#) | [Department of Health & Human Services](#)

Page last updated: 04 August 2008