

ARE YOU GETTING ENOUGH RELIEF FROM YOUR CURRENT RA THERAPY? YES NO

Pill Identifier App
Now available for the iPhone and iPad.
[View Details](#)



Browse all medications [A](#) [B](#) [C](#) [D](#) [E](#) [F](#) [G](#) [H](#) [I](#) [J](#) [K](#) [L](#) [M](#) [N](#) [O](#) [P](#) [Q](#) [R](#) [S](#) [T](#) [U](#) [V](#) [W](#) [X](#) [Y](#) [Z](#) [Advanced Search](#) [Phonetic Search](#)

Enter a search term

- Home
- Drugs A-Z
- Pill Identifier
- Interactions Checker
- News
- Health Professionals
- Q & A
- Mednotes
- Apps

Home → Conditions → Pain → Naprosyn

[Print](#) [Share](#)

[Sign In](#) or [Register](#)



Generic Name: [naproxen](#) (na PROX en)
Brand Names: *Aleve, Anaprox, Anaprox-DS, EC-Naprosyn, Leader Naproxen Sodium, Midol Extended Relief, Naprelan 375, Naprosyn*

- Overview**
- Side Effects
- Dosage
- Interactions
- For Professionals
- More

Bulging Disc Treatment

Read about our 30-minute treatment for bulging disc back pain relief.



What is Naprosyn?

Naprosyn (naproxen) is a nonsteroidal anti-inflammatory drug (NSAID). Naproxen works by reducing hormones that cause inflammation and pain in the body.

Naprosyn is used to treat pain or inflammation caused by conditions such as rheumatoid arthritis, osteoarthritis, ankylosing spondylitis, tendinitis, bursitis, gout, or menstrual cramps.

Naprosyn may also be used for purposes not listed in this medication guide.

Important information

You should not use Naprosyn if you have a history of allergic reaction to aspirin or other NSAID (nonsteroidal anti-inflammatory drug).

Naprosyn may increase your risk of heart attack or stroke, especially if you use it long term or have heart disease. Do not use this medicine just before or after heart bypass surgery (coronary artery bypass graft, or CABG).

Naprosyn may also cause stomach or intestinal bleeding, which can be fatal. These conditions can occur without warning while you are taking Naprosyn.

Before taking this medicine

Naprosyn may increase your risk of heart attack or stroke, especially if you use it long term or have heart disease. Do not use this medicine just before or after heart bypass surgery (coronary artery bypass graft, or CABG).

Naprosyn may also cause stomach or intestinal bleeding, which can be fatal. These conditions can occur without warning while you are taking naproxen.

You should not use Naprosyn if you are allergic to naproxen, or if you have a history of allergic reaction to aspirin or other NSAIDs.

Ask a doctor or pharmacist if it is safe for you to take this medicine if you have:

ADVERTISEMENT

ARE YOU GETTING ENOUGH RELIEF FROM YOUR CURRENT RA THERAPY? YES NO

Slideshow: Drug Treatment for Rheumatoid Arthritis - What Are Your Options?



Related Information

- OTC Rx** **Availability**
Rx and/or OTC
- C** **Pregnancy Category**
Risk cannot be ruled out
- N/A** **CSA Schedule**
Not a controlled drug
- Approval History**
Drug history at FDA

Harvard Health Publications

Related: [Back Pain](#)

Reviews

[23 User Reviews](#)

Average User Rating

7.4 [Rate it!](#)

Manufacturer

[Roche Pharmaceuticals](#)

Drug Class

[Nonsteroidal anti-inflammatory agents](#)

Related Drugs

Back Pain
[tramadol](#), [Cymbalta](#), [naproxen](#), [aspirin](#), [ibuprofen](#), [Mbre...](#)

Ankylosing Spondylitis
[prednisone](#), [naproxen](#), [Humira](#), [aspirin](#), [Celebrex](#), [diclofenac](#), [Mbre...](#)

Pain
[tramadol](#), [oxycodone](#), [naproxen](#), [acetaminophen](#), [Tylenol](#), [Mbre...](#)

Gout, Acute
[naproxen](#), [ibuprofen](#), [methylprednisolone](#), [indomethacin](#), [Aleve](#), [Advil](#), [Mbre...](#)

[▶ 16 more conditions...](#)

Naprosyn (naproxen) Images

- a history of heart attack, stroke, or blood clot;
- heart disease, congestive heart failure, high blood pressure;
- a history of stomach ulcers or bleeding;
- liver or kidney disease;
- asthma;
- polyps in your nose;
- a bleeding or blood clotting disorder; or
- if you smoke.

FDA pregnancy category C. Before using Naprosyn, tell your doctor if you are pregnant or plan to become pregnant during treatment. Taking naproxen during the last 3 months of pregnancy may result in birth defects. Do not take Naprosyn during pregnancy unless your doctor has told you to.

See also: [Pregnancy and breastfeeding warnings](#) (in more detail)

Naproxen can pass into breast milk and may harm a nursing baby. Do not use this medication without telling your doctor if you are breast-feeding a baby.

Do not give this medicine to a child younger than 2 years old without medical advice.

Age 55+ with Joint Pain?



vital3.com

Join hundreds of thousands who found relief with this new formula!

[Sciatica Exercises?](#) ▾

[Treat Overactive Bladder](#) ▾

How should I take Naprosyn?

Use Naprosyn exactly as directed on the label, or as prescribed by your doctor. Do not use in larger or smaller amounts or for longer than recommended.

The delayed-release tablet is a slower-acting form of Naprosyn and should be used only for treating arthritis or ankylosing spondylitis. Follow your doctor's instructions.

Do not crush, chew, or break an extended-release or enteric-coated tablet. Swallow the pill whole. The extended-release pill is specially made to release medicine slowly in the body. The enteric-coated pill has a special coating to protect your stomach.

Shake the oral suspension (liquid) well just before you measure a dose. Measure the liquid with a special dose-measuring spoon or medicine cup. If you do not have a dose-measuring device, ask your pharmacist for one.

If you use this medication long-term, you may need frequent medical tests at your doctor's office.

Store Naprosyn at room temperature away from moisture and heat.

What happens if I miss a dose?

Since Naprosyn is used when needed, you may not be on a dosing schedule. If you are on a schedule, use the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not use extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid?

Ask a doctor or pharmacist before using any cold, allergy, or pain medication. Many medicines available over the counter contain aspirin or other medicines similar to naproxen (such as ibuprofen or ketoprofen). Taking certain products together can cause you to get too much of this type of medication. Check the label to see if a medicine contains naproxen, aspirin, ibuprofen, or ketoprofen.



Naprosyn (naproxen) 500 mg



Naprosyn (naproxen) 500 mg

[View all 10 images](#)

Get Updates by Email

Email me:

News and Warnings related to this drug

FDA Medwatch Alerts for all Medications

News Roundup

Related News and Articles

Health Tip: Vary Your Exercise Regimen

February 13, 2015

Repeating the same exercises day after day can lead to an overuse injury, and cause you to miss exercise when you're not feeling well. Variety is the...

Ketorolac Tromethamine Injection by Hospira: Recall - Particulate in Glass Vials

February 11, 2015

ISSUE: Hospira announced a voluntary recall of ketorolac tromethamine injection, USP in the United States and Singapore due to potential particulate. The...

Mercury in Seafood May Raise Risk of Autoimmune Diseases in Women: Study

February 10, 2015

The mercury found in some seafood may be linked to autoimmune disorders among women of childbearing age, new research suggests. Autoimmune diseases develop...

Avoid drinking alcohol. It may increase your risk of stomach bleeding.

Avoid exposure to sunlight or tanning beds. Naprosyn can make you sunburn more easily. Wear protective clothing and use sunscreen (SPF 30 or higher) when you are outdoors.

If you're 50+ read this

Hear what doctors have to say
about the latest joint relief trick



Naprosyn side effects

Get emergency medical help if you have any of these signs of an allergic reaction to Naprosyn: hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Stop using Naprosyn and call your doctor at once if you have:

- chest pain, weakness, shortness of breath, slurred speech, problems with vision or balance;
- black, bloody, or tarry stools;
- coughing up blood or vomit that looks like coffee grounds;
- swelling or rapid weight gain, little or no urinating;
- nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes);
- bruising, severe tingling, numbness, pain, muscle weakness;
- fever, headache, neck stiffness, chills, increased sensitivity to light, purple spots on the skin, and/or seizure (convulsions) or
- severe skin reaction -- fever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain, followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling.

Common Naprosyn side effects may include:

- upset stomach, mild heartburn or stomach pain, diarrhea, constipation;
- bloating, gas;
- dizziness, headache, nervousness;
- skin itching or rash;
- blurred vision; or
- ringing in your ears.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

See also: [Side effects](#) (in more detail)

What other drugs will affect Naprosyn?

Ask your doctor before using Naprosyn if you take an antidepressant such as citalopram, escitalopram, fluoxetine (Prozac), fluvoxamine, paroxetine, sertraline (Zoloft), trazodone, or vilazodone. Taking any of these medicines with an NSAID may cause you to bruise or bleed easily.

Ask a doctor or pharmacist if it is safe for you to use this medicine if you are also using any of the following drugs:

- a blood thinner (warfarin, Coumadin);
- lithium;
- methotrexate;
- a diuretic or "water pill";
- steroids (prednisone and others);
- aspirin or other NSAIDs--ibuprofen (Advil, Motrin), celecoxib, diclofenac, indomethacin, meloxicam, and others; or

- heart or blood pressure medication--benazepril, candesartan, enalapril, lisinopril, losartan, olmesartan, quinapril, ramipril, telmisartan, valsartan, and others.

This list is not complete. Other drugs may interact with Naprosyn, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible interactions are listed in this medication guide.

[Google](#) [Facebook](#) [Twitter](#) [Print](#) [Email](#) [+ Add to Mednotes](#)

Next Page → [Side Effects](#)

More about Naprosyn (naproxen)

- [Side Effects](#)
- [During Pregnancy](#) or [Breastfeeding](#)
- [Dosage Information](#)
- [Drug Images](#)
- [Drug Interactions](#)
- [Support Group](#)
- [Pricing & Coupons](#)
- [En Espanol](#)
- [23 Reviews](#) - [Add your own review/rating](#)
- [Generic Availability](#)

Consumer resources

- [Naprosyn](#)
- [Naprosyn suspension](#)
- [Naprosyn](#) (Advanced Reading)
- *Other brands:* [Aleve](#), [Anaprox](#), [Naprelan](#), [Anaprox-DS](#), [More \(2\)](#) »

Professional resources

- [Naprosyn](#) (FDA)
- [Naproxen Sodium](#) (AHFS Monograph)

Other formulations

- [EC-Naprosyn](#)

Related treatment guides

- [Back Pain](#)
- [Headache](#)
- [Chronic Myofascial Pain](#)
- [Ankylosing Spondylitis](#)
- [Pain](#)
- [More \(16\)](#) »

Where can I get more information?

- Your pharmacist can provide more information about Naprosyn.

Bulging Disc Treatment

Read about our 30-minute treatment for bulging disc back pain relief.



Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Naprosyn only for the indication prescribed.

Disclaimer: Every effort has been made to ensure that the information provided by Cerner Multum, Inc. ("Multum") is accurate, up-to-date, and complete, but no guarantee is made to that effect. Drug information contained herein may be time sensitive. Multum information has been compiled for use by healthcare practitioners and consumers in the United States and therefore Multum does not warrant that uses outside of the United States are appropriate, unless specifically indicated otherwise. Multum's drug information does not endorse drugs, diagnose patients or recommend therapy. Multum's drug information is an informational resource designed to assist licensed healthcare practitioners in caring for their patients and/or to serve consumers viewing this service as a supplement to, and not a substitute for, the expertise, skill, knowledge and judgment of healthcare practitioners. The absence of a warning for a given drug or drug combination in no way should be construed to indicate that the drug or drug combination is safe, effective or appropriate for any given patient. Multum does not assume any responsibility for any aspect of healthcare administered with the aid of information Multum provides. The information contained herein is not intended to cover all possible uses, directions, precautions, warnings, drug interactions, allergic reactions, or adverse effects. If you have questions about the drugs you are taking, check with your doctor, nurse or pharmacist.

Support About Drugs.com Your Privacy

- [Advertise With Us](#)
- [Frequently Asked Questions](#)
- [Ask a Question](#)
- [Content Submissions](#)
- [Mobile Apps](#)
- [About Drugs.com](#)
- [Contact Us](#)
- [Sitemap](#)

Follow:     

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment.

Data sources include Micromedex® (updated Feb 4th, 2015), Cerner Multum™ (updated Jan 15th, 2015), Wolters Kluwer™ (updated Feb 4th, 2015) and others. To view content sources and attributions, refer to our [editorial policy](#).

Copyright © 2000-2015 Drugs.com. All rights reserved.



We comply with the HONcode standard for trustworthy health information.
[Verify here](#)



Ad Choices

[Terms of Use](#) · [Privacy Policy](#)