



The Depression Treatment Device



Use for 20 minutes, once or twice a day.

ADVERTISEMENT

Home › Conditions › Muscle Spasm › Flexeril

Print Share

Flexeril

Generic Name: [cyclobenzaprine](#) (sye kloe BEN za preen)

Brand Names: *Amrix, Fexmid, Flexeril*

1 Trick to Fibromyalgia

See How to Relieve Fibromyalgia With This Fast and Easy Trick



Overview

Side Effects

Dosage

Interactions

Professional

More

What is Flexeril?

Flexeril (cyclobenzaprine) is a muscle relaxant. It works by blocking nerve impulses (or pain sensations) that are sent to your brain.

Flexeril is used together with rest and physical therapy to [treat](#) skeletal muscle conditions such as pain or injury.

Flexeril may also be used for other purposes not listed in this medication guide.

ADVERTISEMENT

New Sciatica Treatment

northamericanspine.com/Sciatica

Read about our 30-minute, minimally invasive treatment for sciatica.

Male Catheter Supplies

Important information

Do not take Flexeril if you have used an MAO inhibitor such as isocarboxazid (Marplan), tranylcypromine (Parnate), phenelzine (Nardil), or selegiline (Eldepryl, Emsam) within the past 14 days. Serious, life-threatening side effects can occur if you take Flexeril before the MAO inhibitor has cleared from [your body](#).

You should not take Flexeril if you have recently had a heart attack, or if you have a heart rhythm disorder, congestive heart failure, heart block, or an overactive thyroid.

Flexeril can cause side effects that may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be awake and alert. Avoid drinking alcohol, which can increase some of the side effects of Flexeril.

Before taking this medicine

Do not take Flexeril if you have used an MAO inhibitor such as isocarboxazid (Marplan), tranylcypromine (Parnate), phenelzine (Nardil), or selegiline (Eldepryl, Emsam) within the past 14 days. Serious, life-threatening side effects can occur if you take Flexeril before the MAO inhibitor has cleared from your body. Do not use Flexeril if you have recently had a heart attack, or if you have:

- a heart rhythm disorder;
- congestive heart failure;
- heart block; or
- an overactive thyroid.

Before using Flexeril, tell your [doctor](#) if you are allergic to any drugs, or if you have:

- problems with urination;
- enlarged prostate;

[1 Trick to Fibromyalgia](#)

[Adult Overactive Bladder](#)

[STELARA® \(ustekinumab\)](#)

[Reduce Nerve Pain Now](#)


[Apply for Disability](#)

[3 Herbs That Beat Anxiety](#)

[7 Hidden Dementia Signs](#)


[Parkinson's Bike Therapy](#)

Slideshow: Fighting the Fight: Fibromyalgia Explained



- glaucoma; or
- liver disease.

If you have any of these conditions, you may need a dose adjustment or special tests to safely take Flexeril.

FDA pregnancy category B. Flexeril is not expected to be harmful to an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant during [treatment](#) . It is not known whether cyclobenzaprine passes into breast milk or if it could harm a nursing baby. Do not use this medication without telling your doctor if you are breast-feeding a baby.

See also: [Pregnancy and breastfeeding warnings](#) (in more detail)

Older adults may be more sensitive to the side effects of Flexeril.

Get Sciatica Pain Relief

One 30-minute, minimally
invasive pain relief procedure,
learn more



How should I take Flexeril?

Take this medication exactly as it was prescribed for you. Do not take Flexeril in larger amounts, or take it for longer than recommended by your doctor. Follow the directions on your prescription label.

Take this medicine with a full glass of water.

Flexeril is only part of a complete program of treatment that may also include rest, physical therapy, or other pain relief measures. Follow your doctor's instructions.

Store Flexeril at room temperature away from moisture, heat, and light.

What happens if I miss a dose?

Take the missed dose as soon as you remember. If it is almost time for your next dose, skip the missed dose and take the medicine at the next regularly scheduled time. Do not take extra medicine to make up

the missed dose.

What happens if I overdose?

Seek emergency medical attention if you think you have used too much of this medicine. An overdose of Flexeril can be fatal.

Overdose symptoms may include drowsiness, fast heartbeat, tremors or shaking, slurred speech, confusion, nausea, vomiting, hallucinations (seeing things), chest pain, or seizure (convulsions).

What should I avoid?

Flexeril can cause side effects that may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be awake and alert. Avoid drinking alcohol, which can increase some of the side effects of Flexeril. Cold or allergy medicine, narcotic pain medicine, sleeping pills, and medicine for seizures, depression or anxiety can add to sleepiness caused by Flexeril. Tell your doctor if you regularly use any of these medicines, or any other muscle relaxer.

Parkinson's Bike Therapy

New Therapy Brings Hope &
Results. Learn More, Request a
Free eBook!



Flexeril side effects

Get emergency medical help if you have any of these signs of an allergic reaction to Flexeril: hives; difficulty breathing; swelling of your face, lips, tongue, or throat. Stop using Flexeril and call your doctor at once if you have any of these serious side effects:

- fast, pounding, or uneven heartbeats;
- chest pain or heavy feeling, pain spreading to the arm or shoulder, nausea, sweating, general ill feeling;

sudden numbness or weakness, especially on one side of the body;

- sudden headache, confusion, problems with vision, speech, or balance;
- feeling light-headed, fainting;
- confusion, weakness, lack of coordination;
- nausea, stomach pain, low fever, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes);
- seizure (convulsions);
- unusual thoughts or behavior, hallucinations (seeing things); or
- easy bruising or bleeding, unusual weakness.

Less serious Flexeril side effects may include:

- dry mouth or throat;
- blurred vision;
- drowsiness, dizziness, tired feeling;
- loss of appetite, stomach pain, nausea;
- diarrhea, constipation, gas; or
- muscle weakness.

This is not a complete list of side effects and others may occur. Tell your doctor about any unusual or bothersome side effect. You may report side effects to FDA at 1-800-FDA-1088.

See also: [Side effects](#) (in more detail)

What other drugs will affect Flexeril?

Many drugs can interact with Flexeril. Below is just a partial list. Tell your doctor if you are using:

- atropine (Donnatal, and others), benztropine (Cogentin), dimenhydrinate (Dramamine), methscopolamine (Pamine), or scopolamine (Transderm-Scop);
- a bronchodilator such as ipratropium (Atrovent) or tiotropium (Spiriva);
- glycopyrrolate (Robinul);
- guanethidine (Ismelin);
- mepenzolate (Cantil);
- tramadol (Ultram);
- bladder or urinary medications such as darifenacin (Enablex), flavoxate (Urispas), oxybutynin (Ditropan, Oxytrol), tolterodine (Detrol), or solifenacin (Vesicare); or

- irritable bowel medications such as dicyclomine (Bentyl), hyoscyamine (Anaspaz, Cystospaz, Levsin, and others), or propantheline (Pro-Banthine).

This list is not complete and there may be other drugs that can interact with Flexeril. Tell your doctor about all your prescription and over-the-counter medications, vitamins, minerals, herbal products, and drugs prescribed by other doctors. Do not start a new medication without telling your doctor.

Next → [Side Effects](#)



Add to My Med List

More about Flexeril (cyclobenzaprine)

- [Side Effects](#)
- [During Pregnancy or Breastfeeding](#)
- [Dosage Information](#)
- [Drug Images](#)
- [Drug Interactions](#)
- [Support Group](#)
- [En Espanol](#)
- [125 Reviews - Add your own review/rating](#)

Consumer resources

- [Flexeril](#)
- [Flexeril](#) (Advanced Reading)
- *Other brands:* [Amrix](#), [Fexmid](#)

Professional resources

- [Flexeril](#) (AHFS Monograph)
- [Flexeril](#) (FDA)

Related treatment guides

- [Muscle Spasm](#)
- [Sciatica](#)

Where can I get more information?

- Your pharmacist can provide more information about Flexeril.

4 veggies to never eat:

Cut down a bit of stomach fat every day by never eating these 4 foods



Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Flexeril only for the indication prescribed.

Disclaimer: Every effort has been made to ensure that the information provided is accurate, up-to-date, and complete, but no guarantee is made to that effect. Drug information contained herein may be time sensitive. Multum's drug information is an informational resource designed to assist licensed healthcare practitioners in caring for their patients and/or to serve consumers viewing this service as a supplement to, and not a substitute for, the expertise, skill, knowledge and judgment of healthcare practitioners. The absence of a warning for a given drug or drug combination in no way should be construed to indicate that the drug or drug combination is safe, effective or appropriate for any given patient. The information contained herein is not intended to cover all possible uses, directions, precautions, warnings, drug interactions, allergic reactions, or adverse effects.

Copyright 1996-2015 Cerner Multum, Inc. Version: 3.11. Revision Date: 4/12/2009 4:38:46 PM.

DRUG STATUS



Availability
Prescription only



Pregnancy Category
No proven risk in humans



CSA Schedule
Not a controlled drug



Approval History
Drug history at FDA

Drug Class

[Skeletal muscle relaxants](#)

Related Drugs

Muscle Spasm
[cyclobenzaprine](#),
[Soma](#), [baclofen](#),
[diazepam](#), [Valium](#),

ADVERTISEMENT

tizanidine,
methocarbamol,
Robaxin, Zanaflex,
carisoprodol,
metaxalone,
Skelaxin, More...

Sciatica

cyclobenzaprine,
naproxen, Tylenol,
acetaminophen,
aspirin, ibuprofen,
diclofenac,
Paracetamol,
indomethacin,
Voltaren, Advil,
Aleve, More...

Flexeril Rating

125 User Reviews

6.7/10

Flexeril Images



Flexeril (cyclobenzaprine) 10 mg



[View larger images](#)

Get Updates by Email

Sign up for daily, weekly or monthly emails to receive news updates to your inbox!

News and Warnings related to this drug

FDA Medwatch Alerts for all Medications

News Roundup **Daily**

Related News and Articles

Steroids No Better for Sciatica Pain Than Placebo, Study Finds

May 19, 2015

Spinal Stimulation System Relieves Pain Without Tingling

May 11, 2015

Medication Errors Occur Every 8 Minutes in U.S. Children

October 20, 2014

Drugs.com Mobile Apps

The easiest way to lookup drug information, identify

pills, check interactions and set up your own personal medication records. Available for Android and iOS devices.

Explore Apps



SUPPORT

Help Center
Frequent Questions
Sitemap
Contact Us

ABOUT

About Drugs.com
Advertising Policy
Content Submissions
Drugs.com Blog

TERMS & PRIVACY


Editorial Policy
Privacy Policy
Terms of Use
Attribution & Citations

CONNECT



Subscribe to receive email notifications whenever new articles are published.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex® (updated May 8th, 2015), Cerner Multum™ (updated May 18th, 2015), Wolters Kluwer™ (updated May 6th, 2015) and others. To view content sources and attributions, please refer to our [editorial policy](#).



We comply with the HONcode standard for trustworthy health information - verify here

Copyright © 2000-2015 Drugs.com. All rights reserved.