

[Home](#)[Disorders A - Z](#)[Bell's Palsy](#)[Information Page](#)[Publications](#)[Organizations](#)[News](#)[Research literature](#)[Organizations](#)[Professional](#)[Societies](#)[Government Resources](#)[NINDS Clinical](#)[Trials](#)[En Español](#)[Order Brochures](#)[News From NINDS](#)[Funding Information](#)[Research Programs](#)[Training & Career Awards](#)[Enhancing Diversity](#)[Find People](#)[About NINDS](#)

Bell's Palsy Fact Sheet

[See a list of all NINDS Disorders](#)[Get Web page suited for printing](#)[Email this to a friend or colleague](#)[Request free mailed brochure](#)[Parálisis de Bell](#)

Table of Contents (click to jump to sections)

[Listen](#)[What is Bell's Palsy?](#)[What are the Symptoms?](#)[What Causes Bell's Palsy?](#)[Who Gets it?](#)[How is it Diagnosed?](#)[How is it Treated?](#)[What is the Prognosis?](#)[What Research is Being Done?](#)[Where can I get more information?](#)

What is Bell's Palsy?

Bell's palsy is a form of temporary facial paralysis resulting from damage or trauma to the facial nerves. The facial nerve—also called the 7th cranial nerve—travels through a narrow, bony canal (called the Fallopian canal) in the skull, beneath the ear, to the muscles on each side of the face. For most of its journey, the nerve is encased in this bony shell.

Each facial nerve directs the muscles on one side of the face, including those that control eye blinking and closing, and facial expressions such as smiling and frowning. Additionally, the facial nerve carries nerve impulses to the *lacrimal* or tear glands, the saliva glands, and the muscles of a small bone in the middle of the ear called the *stapes*. The facial nerve also transmits taste sensations from the tongue.

When Bell's palsy occurs, the function of the facial nerve is disrupted, causing an interruption in the messages the brain sends to the facial muscles. This interruption results in facial weakness or paralysis.

Bell's palsy is named for Sir Charles Bell, a 19th century Scottish surgeon who described the facial nerve and its connection to the condition. The disorder, which is not related to stroke, is the most common cause of facial paralysis. Generally, Bell's palsy affects only one of the paired facial nerves and one side of the face, however, in rare cases, it can affect both sides.

What are the Symptoms?

Because the facial nerve has so many functions and is so complex, damage to the nerve or a disruption in its function can lead to many problems. Symptoms of Bell's palsy can vary from person to person and range in severity from mild weakness to total paralysis. These symptoms may include twitching, weakness, or paralysis on one or rarely both sides of the face. Other symptoms may include drooping of the eyelid and corner of the mouth, drooling, dryness of the eye or mouth, impairment of taste, and excessive tearing in one eye. Most often these symptoms, which usually begin suddenly and reach their peak within 48 hours, lead to significant facial distortion.

Other symptoms may include pain or discomfort around the jaw and behind the ear, ringing in one or both ears, headache, loss of taste, hypersensitivity to sound on the affected side, impaired speech, dizziness, and difficulty eating or drinking.

What Causes Bell's Palsy?

Bell's palsy occurs when the nerve that controls the facial muscles is swollen, inflamed, or compressed, resulting in facial weakness or paralysis. Exactly what causes this damage, however, is unknown.

Most scientists believe that a viral infection such as viral meningitis or the common cold sore virus—*herpes simplex*—causes the disorder. They believe that the facial nerve swells and becomes inflamed in reaction to the infection, causing pressure within the Fallopian canal and leading to ischemia (the restriction of blood and oxygen to the nerve cells). In some mild cases (where recovery is rapid), there is damage only to the myelin sheath of the nerve. The myelin sheath is the fatty covering—which acts as an insulator—on nerve fibers in the brain.

The disorder has also been associated with influenza or a flu-like illness, headaches, chronic middle ear infection, high blood pressure, diabetes, sarcoidosis, tumors, Lyme disease, and trauma such as skull fracture or facial injury.

Who Gets it?

Bell's palsy afflicts approximately 40,000 Americans each year. It affects men and women equally and can occur at any age, but it is less common before age 15 or after age 60. It disproportionately attacks people who have diabetes or upper respiratory ailments such as the flu or a cold.

How is it Diagnosed?

A diagnosis of Bell's palsy is made based on clinical presentation -- including a distorted facial appearance and the inability to move muscles on the affected side of the face -- and by ruling out other possible causes of facial paralysis. There is no specific laboratory test to confirm diagnosis of the disorder.

Generally, a physician will examine the individual for upper and lower facial weakness. In most cases this weakness is limited to one side of the face or occasionally isolated to the forehead, eyelid, or mouth. A test called electromyography (EMG) can confirm the presence of nerve damage and determine the severity and the extent of nerve involvement. Blood tests can sometimes be helpful in diagnosing other concurrent problems such as diabetes and certain infections. A magnetic resonance imaging (MRI) or computed tomography (CT) scan can eliminate other structural causes of pressure on the facial nerve.

How is it Treated?

Bell's palsy affects each individual differently. Some cases are mild and do not require treatment as the symptoms usually subside on their own within 2 weeks. For others, treatment may include medications and other therapeutic options. If an obvious source is found to cause Bell's palsy (e.g., infection), directed treatment can be beneficial.

REVIEWED*By Chris Tighe at 12:25 pm, Aug 16, 2016*

Recent studies have shown that steroids such as the steroid **prednisone** -- used to reduce inflammation and swelling -- are effective in treating Bell's palsy. Other drugs such as acyclovir -- used to fight viral herpes infections -- may also have some benefit in shortening the course of the disease. Analgesics such as aspirin, acetaminophen, or ibuprofen may relieve pain. Because of possible drug interactions, individuals taking prescription medicines should always talk to their doctors before taking any over-the-counter drugs.

Another important factor in treatment is eye protection. Bell's palsy can interrupt the eyelid's natural blinking ability, leaving the eye exposed to irritation and drying. Therefore, keeping the eye moist and protecting the eye from debris and injury, especially at night, is important. Lubricating eye drops, such as artificial tears or eye ointments or gels, and eye patches are also effective.

Other therapies such as physical therapy, facial massage or acupuncture may provide a potential small improvement in facial nerve function and pain.

In general, decompression surgery for Bell's palsy -- to relieve pressure on the nerve -- is controversial and is seldom recommended. On rare occasions, cosmetic or reconstructive surgery may be needed to reduce deformities and correct some damage such as an eyelid that will not fully close or a crooked smile.

What is the Prognosis?

The prognosis for individuals with Bell's palsy is generally very good. The extent of nerve damage determines the extent of recovery. Improvement is gradual and recovery times vary. With or without treatment, most individuals begin to get better within 2 weeks after the initial onset of symptoms and most recover completely, returning to normal function within 3 to 6 months. For some, however, the symptoms may last longer. In a few cases, the symptoms may never completely disappear. In rare cases, the disorder may recur, either on the same or the opposite side of the face.

What Research is Being Done?

Within the Federal Government, the National Institute of Neurological Disorders and Stroke (NINDS), part of the National Institutes of Health (NIH), is responsible for supporting and conducting research on brain and nervous system disorders, including Bell's palsy. The NINDS conducts research in its laboratories at the NIH, in Bethesda, Maryland, and supports research through grants to major medical institutions across the country.

The NINDS conducts and supports an extensive research program of basic science to increase understanding of how the nervous system works and what causes the system to sometimes go awry, leading to dysfunction. Part of this research program focuses on learning more about the circumstances that lead to nerve damage and the conditions that cause injuries and damage to nerves. Knowledge gained from this research may help scientists find the definitive cause of Bell's palsy, leading to the discovery of new effective treatments for the disorder. Other NINDS-supported research is aimed at developing methods to repair damaged nerves and restore full use and strength to injured areas, and finding ways to prevent nerve damage and injuries from occurring.

Where can I get more information?

For more information on neurological disorders or research programs funded by the National Institute of Neurological Disorders and Stroke, contact the Institute's Brain Resources and Information Network (BRAIN) at:

BRAIN
P.O. Box 5801
Bethesda, MD 20824
800-352-9424
<http://www.ninds.nih.gov>

Information also is available from the following organizations:

[National Organization for Rare Disorders \(NORD\)](#)
55 Kenosia Avenue
Danbury, CT 06810
orphan@rarediseases.org
<http://www.rarediseases.org> ⓘ
Tel: 203-744-0100; Voice Mail: 800-999-NORD (6673)
Fax: 203-798-2291

"Bell's PalsyFact Sheet", NINDS, Publication date April 2003.

NIH Publication No. 03-5114

Back to [Bell's Palsy Information Page](#)

[See a list of all NINDS Disorders](#)

Publicaciones en Español

► [Parálisis de Bell](#)

Prepared by:
Office of Communications and Public Liaison
National Institute of Neurological Disorders and Stroke
National Institutes of Health
Bethesda, MD 20892

NINDS health-related material is provided for information purposes only and does not necessarily represent endorsement by or an official position of the National Institute of Neurological Disorders and Stroke or any other Federal agency. Advice on the treatment or care of an individual patient should be obtained through consultation with a physician who has examined that patient or is familiar with that patient's medical history.

All NINDS-prepared information is in the public domain and may be freely copied. Credit to the NINDS or the NIH is appreciated.

Last Modified February 5, 2016





NIH...Turning Discovery Into Health®