

# anterior cervical discectomy & fusion (ACDF)

A+ | Reset | A-

# Overview

Anterior cervical discectomy and fusion (ACDF) is a surgery to remove a herniated or degenerative disc in the neck area of the spine. The incision is made in the front of the spine through the throat area. After the disc is removed, a bone graft is inserted to fuse together the bones above and below the disc space. Your doctor may recommend a discectomy if physical therapy or medications fail to relieve your neck or arm pain caused by pressure on the spinal nerves. Patients typically go home the same day; recovery time takes 4 weeks.



Figure 1, top. (top view of vertebra) Degenerative disc disease causes the discs (purple) to dry out. Tears in the disc annulus can allow the gel-filled nucleus material to escape and compress the spinal cord causing numbness and weakness. Bone spurs may develop which can lead to a narrowing of the nerve root canal (foraminal stenosis). The pinched spinal nerve becomes swollen and painful.

# What is an anterior cervical discectomy & fusion (ACDF)?

Discectomy literally means "cutting out the disc." A discectomy can be performed anywhere along the spine from the neck (cervical) to the low back (lumbar). The surgeon reaches the damaged disc from the front (anterior) of the spine through the throat area. By moving aside the neck muscles, trachea, and esophagus, the disc and bony vertebrae are exposed. Surgery from the front of the neck is more accessible than from the back (posterior) because the disc can be reached without disturbing the spinal cord, spinal nerves, and the strong neck muscles. Depending on your particular symptoms, one disc (single-level) or more (multi-level) may be removed.

After the disc is removed, the space between the bony vertebrae is empty. To prevent the vertebrae from collapsing and rubbing together, a spacer bone graft is inserted to fill the open disc space. The graft serves as a bridge between the two vertebrae to create a spinal fusion. The bone graft and vertebrae are fixed in place with metal plates and screws. Following surgery the body begins its natural healing process and new bone cells grow around the graft. After 3 to 6 months, the bone graft should join the two vertebrae and form one solid piece of bone. The instrumentation and fusion work together, similar to reinforced concrete.

Bone grafts come from many sources. Each type has advantages and disadvantages.

- Autograft bone comes from you. The surgeon takes your own bone cells from the hip (iliac crest). This graft has a higher rate of fusion because it has bone-growing cells and proteins. The disadvantage is the pain in your hipbone after surgery. Harvesting a bone graft from your hip is done at the same time as the spine surgery. The harvested bone is about a half inch thick the entire thickness of bone is not removed, just the top half layer.
- Allograft bone comes from a donor (cadaver). Bone-bank bone is collected from people who have
  agreed to donate their organs after they die. This graft does not have bone-growing cells or proteins,
  yet it is readily available and eliminates the need to harvest bone from your hip. Allograft is shaped
  like a doughnut and the center is packed with shavings of living bone tissue taken from your spine
  during surgery.
- Bone graft substitute comes from man-made plastic, ceramic, or bioresorbable compounds. Often called cages, this graft material is packed with shavings of living bone tissue taken from your spine during surgery.

After fusion you may notice some range of motion loss, but this varies according to neck mobility before surgery and the number of levels fused. If only one level is fused, you may have similar or even better range of motion than before surgery. If more than two levels are fused, you may notice limits in turning your head and looking up

## **Mayfield services**

At Mayfield, patients who come to us with neck and back problems are given a rapid review of their medical condition within a few days ... not weeks. It's a treatment process called **Priority Consult**.

Nearly 80% of our spine patients are able to recover with nonsurgical treatment. But when physical therapy, medications, and spinal injections fail to relieve neck or arm pain, we help patients with surgery.

We perform around 650 ACDF surgeries each year. Many of these are done as an outpatient procedure, allowing patients to go home the same day.

To make an appointment call 513-221-1100.

Make an Appointment

## Related: <u>Mayfield Sweeps in Cincinnati</u> <u>Magazine's 'Top Doctors' 2015</u>



Learn more about minimally invasive outpatient spine surgery at the spine surgery center.



susan's story

and down. Motion-preserving artificial disc replacements have emerged as an alternative to fusion. Similar to knee replacement, the artificial disc is inserted into the damaged joint space and preserves motion, whereas fusion eliminates motion. Outcomes for artificial disc compared to ACDF are similar, but long-term results of motion preservation and adjacent level disease are not yet proven. Talk with your surgeon about whether ACDF or artificial disc replacement is most appropriate for you.

## Who is a candidate?

# You may be a candidate for discectomy if you have:

- diagnostic tests (<u>MRI, CT, myelogram</u>) show that you have a herniated or degenerative disc
- significant weakness in your hand or arm
- arm pain worse than neck pain
- · symptoms that have not improved with physical therapy or medication

### ACDF may be helpful in treating the following conditions:

- Bulging and herniated disc: The gel-like material within the disc can bulge or rupture through a weak area in the surrounding wall (annulus). Irritation and swelling occurs when this material squeezes out and painfully presses on a nerve.
- <u>Degenerative disc disease</u>: As discs naturally wear out, bone spurs form and the facet joints inflame. The discs dry out and shrink, losing their flexibility and cushioning properties. The disc spaces get smaller. These changes lead to canal stenosis or disc herniation (Fig. 1).

## The surgical decision

Most herniated discs heal after a few months of nonsurgical treatment. Your doctor may recommend treatment options, but only you can decide whether surgery is right for you. Be sure to consider all the risks and benefits before making your decision. Only 10% of people with herniated disc problems have enough pain after 6 weeks of conservative treatment to consider surgery.

Your surgeon will also discuss the risks and benefits of different types of bone graft material. Autograft is the gold standard for rapid healing and fusion, but the hip incision can be painful and at times lead to complications. Allograft (bone-bank) is more commonly used and has proven to be as effective for routine 1 and 2 level fusions in non-smokers.

## Who performs the procedure?

A neurosurgeon or an orthopedic surgeon can perform spine surgery. Many spine surgeons have specialized training in complex spine surgery. Ask your surgeon about their training, especially if your case is complex or you've had more than one spinal surgery.

## What happens before surgery?

You may be scheduled for presurgical tests (e.g., blood test, electrocardiogram, chest X-ray) several days before surgery. In the doctor's office, you will sign consent and other forms so that the surgeon knows your medical history (allergies, medicines/vitamins, bleeding history, anesthesia reactions, previous surgeries). Discuss all medications (prescription, over-the-counter, and herbal supplements) you are taking with your health care provider. Some medications need to be continued or stopped the day of surgery.

Stop taking all non-steroidal anti-inflammatory medicines (Naprosyn, Advil, Motrin, Nuprin, Aleve, etc.) and blood thinners (Coumadin, Plavix, etc.) 1 to 2 weeks before surgery as directed by the doctor. Additionally, stop smoking, chewing tobacco, and drinking alcohol 1 week before and 2 weeks after surgery because these activities can cause bleeding problems. No food or drink is permitted past midnight the night before surgery.

#### Smoking

The most important thing you can do to ensure the success of your spinal surgery is quit smoking. This includes cigarettes, cigars, pipes, chewing tobacco, and smokeless tobacco (snuff, dip). Nicotine prevents bone growth and puts you at higher risk for a failed fusion. Patients who smoked had failed fusions in up to 40% of cases, compared to only 8% among non-smokers [1]. Smoking also decreases your blood circulation, resulting in slower wound healing and an increased risk of infection. Talk with your doctor about ways to help you quit smoking: nicotine replacements, pills without nicotine (Wellbutrin, Chantix), and tobacco counseling programs.

# Morning of surgery

- Shower using antibacterial soap. Dress in freshly washed, loose-fitting clothing.
- Wear flat-heeled shoes with closed backs.
- If you have instructions to take regular medication the morning of surgery, do so with small sips of water.
- Remove make-up, hairpins, contacts, body piercings, nail polish, etc.
- Leave all valuables and jewelry at home (including wedding bands).
- Bring a list of medications (prescriptions, over-the-counter, and herbal supplements) with dosages and the times of day usually taken.
- Bring a list of allergies to medication or foods.

Arrive at the hospital 2 hours before (surgery center 1 hour before) your scheduled surgery time to complete the



gary's story

### **Related links:**

• Neck pain

- Degenerative Disc Disease
- Herniated Cervical Disc

## **Q&A: Artificial disc**

**Q.** Is artificial disc replacement an alternative to fusion?

**A.** One of the downsides of spinal fusion is the loss of mobility in the neck. New on the horizon is the artificial cervical disc, an alternative to fusion that is designed to preserve motion.

There is much debate within the medical community about the efficacy of artificial cervical disc surgery. In clinical trials, the artificial cervical disc had comparable outcomes to ACDF. However, further studies are needed to know the long-term effects. The artificial disc is new technology that many insurers in Ohio and Kentucky are not yet covering. Ask your surgeon if artificial disc replacement is an option for your condition. Then check with your insurance company to determine what costs you may have to pay out of pocket.

See more questions

## clinical trials

Click <u>here</u> for information about clinical trials conducted by our doctors at local Cincinnati hospitals or call 1-800-325-7787.



ACDF Surgery | Anterior Cervical Discectomy and Fusion

necessary paperwork and pre-procedure work-ups. An anesthesiologist will talk with you and explain the effects of anesthesia and its risks. An intravenous (IV) line will be placed in your arm.

# What happens during surgery?

There are seven steps to the procedure. The operation generally takes 1 to 3 hours.

#### Step 1: prepare the patient

You will lie on your back on the operative table and be given anesthesia. Once asleep, your neck area is cleansed and prepped. If a fusion is planned and your own bone will be used, the hip area is also prepped to obtain a bone graft. If a donor bone will be used, the hip incision is unnecessary.

### Step 2: make an incision

A 2-inch skin incision is made on the right or left side of your neck (Fig. 2). The surgeon makes a tunnel to the spine by moving aside muscles in your neck and retracting the trachea, esophagus, and arteries. Finally, the muscles that support the front of the spine are lifted and held aside so the surgeon can clearly see the bony vertebrae and discs.



Figure 2. A 2-inch skin incision is made on the side of your neck.

#### Step 3: locate the damaged disc

With the aid of a fluoroscope (a special X-ray), the surgeon passes a thin needle into the disc to locate the affected vertebra and disc. The vertebrae bones above and below the damaged disc are spread apart with a special retractor.

#### Step 4: remove the disc

The outer wall of the disc is cut (Fig. 3). The surgeon removes about 2/3 of your disc using small grasping tools, and then looks through a surgical microscope to remove the rest of the disc. The ligament that runs behind the vertebrae is removed to reach the spinal canal. Any disc material pressing on the spinal nerves is removed.



Figure 3. The muscles are retracted to expose the vertebra. The disc annulus is cut open and the disc material is removed with grasping tools.

### Step 5: decompress the nerve

Bone spurs that press on your nerve root are removed. The foramen, through which the spinal nerve exits, is

ACDF Surgery | Anterior Cervical Discectomy and Fusion

enlarged with a drill (Fig. 4). This procedure, called a foraminotomy, gives your nerves more room to exit the spinal canal.



Figure 4. (top view) The disc annulus and nucleus are removed to decompress the spinal cord and nerve root. Bone spurs are removed and the spinal foramen is enlarged to free the nerve.

# Step 6. prepare a bone graft fusion

Using a drill, the open disc space is prepared on the top and bottom by removing the outer cortical layer of bone to expose the blood-rich cancellous bone inside. This "bed" will hold the bone graft material that you and your surgeon selected:

- Bone graft from your hip. A skin and muscle incision is made over the crest of your hipbone. Next, a chisel is used to cut through the hard outer layer (cortical bone) to the inner layer (cancellous bone). The inner layer contains the bone-growing cells and proteins. The bone graft is then shaped and placed into the "bed" between the vertebrae (Fig. 5).
- Bone bank or fusion cage. A cadaver bone graft or bioplastic cage is filled with the leftover bone shavings containing bone-growing cells and proteins. The graft is then tapped into the shelf space.



Figure 5. (side view) A bone graft is shaped and inserted into the shelf space between the vertebrae.

The bone graft is often reinforced with a metal plate screwed into the vertebrae to provide stability during fusion. An x-ray is taken to verify the position of the graft, plate, and screws (Fig. 6).



Figure 6. X-ray and illustration showing a metal plate and four screws used to hold the

#### bone graft between the vertebrae while fusion occurs.



Figure 7. Artificial disc replacement preserves motion of the spine segment.

Alternative option: artificial disc replacement (Fig. 7). Instead of a bone graft or fusion cage, an artificial disc device is inserted into the empty disc space. In select patients, it may be beneficial to preserve motion. Talk to your doctor – not all insurance companies will pay for this new technology and out-of-pocket expenses may be incurred.

#### Step 7. close the incision

The spreader retractors are removed. The muscle and skin incisions are sutured together. Steri-Strips or biologic glue is placed across the incision.

# What happens after surgery?

You will awaken in the postoperative recovery area, called the PACU. Blood pressure, heart rate, and respiration will be monitored. Any pain will be addressed. Once awake, you will be moved to a regular room where you'll increase your activity level (sitting in a chair, walking). Patients who have had bone graft taken from their hip may feel more discomfort in their hip than neck incision. Most patients having a 1 or 2 level ACDF are sent home the same day. However, if you have difficulty breathing or unstable blood pressure, you may need to stay overnight.

# **Discharge instructions**

### Discomfort

- After surgery, pain is managed with narcotic medication. Because narcotic pain pills are addictive, they are used for a limited period (2 to 4 weeks). As their regular use can cause constipation, drink lots of water and eat high fiber foods. Laxatives (e.g., Dulcolax, Senokot, Milk of Magnesia) can be bought without a prescription. Thereafter, pain is managed with acetaminophen (e.g., Tylenol).
- Hoarseness, sore throat, or difficulty swallowing may occur in some patients and should not be cause for alarm. These symptoms usually resolve in 1 to 4 weeks.

### Restrictions

- If you had a fusion, do **not** use non-steroidal anti-inflammatory drugs (NSAIDs) (e.g., aspirin; ibuprofen, Advil, Motrin, Nuprin; naproxen sodium, Aleve) for 6 months after surgery. NSAIDs may cause bleeding and interfere with bone healing.
- Do not smoke. Smoking delays healing by increasing the risk of complications (e.g., infection) and inhibits the bones' ability to fuse.
- Do not drive for 2 to 4 weeks after surgery or until discussed with your surgeon.
- Avoid sitting for long periods of time.
- Avoid bending your head forward or backward.
- Do not lift anything heavier than 5 pounds (e.g., gallon of milk).
- · Housework and yard-work are not permitted until the first follow-up office visit. This includes
- gardening, mowing, vacuuming, ironing, and loading/unloading the dishwasher, washer, or dryer.
  Postpone sexual activity until your follow-up appointment unless your surgeon specifies otherwise.

#### Activity

- You may need help with daily activities (e.g., dressing, bathing), but most patients are able to care for themselves right away.
- Gradually return to your normal activities. Walking is encouraged; start with a short distance and gradually increase to 1 to 2 miles daily. A physical therapy program may be recommended.
- If applicable, know how to wear a cervical collar before leaving the hospital. Wear it when walking or riding in a car.

### **Bathing/Incision Care**

- You may shower 1 to 4 days after surgery. Follow your surgeon's specific instructions. No tub baths, hot tubs, or swimming pools until your health care provider says it's safe to do so.
- If you have staples or stitches when you go home, they will need to be removed. Ask your surgeon
  or call the office to find out when.

# When to Call Your Doctor

ACDF Surgery | Anterior Cervical Discectomy and Fusion

- If your temperature exceeds 101° F, or if the incision begins to separate or show signs of infection, such as redness, swelling, pain, or drainage.
- If your swallowing problems interfere with your ability to breathe or drink water.

# **Recovery and prevention**

Schedule a follow-up appointment with your surgeon for 2 weeks after surgery. Recovery time generally lasts 4 to 6 weeks. X-rays may be taken after several weeks to verify that fusion is occurring. The surgeon will decide when to release you back to work at your follow-up visit.

A cervical collar or brace is sometimes worn during recovery to provide support and limit motion while your neck heals or fuses (see <u>Braces & Orthotics</u>). Your doctor may prescribe neck stretches and exercises or <u>physical therapy</u> once your neck has healed.

If you had a bone graft taken from your hip, you may experience pain, soreness, and stiffness at the incision. Get up frequently (every 20 minutes) and move around or walk. Don't sit or lie down for long periods of time.

Recurrences of neck pain are common. The key to avoiding recurrence is prevention:

- Proper lifting techniques
- Good posture during sitting, standing, moving, and sleeping
- Appropriate <u>exercise</u> program
- An ergonomic work area
- Healthy weight and lean body mass
- A positive attitude and relaxation techniques (e.g., stress management)
- No smoking

# What are the results?

Anterior cervical discectomy is successful in relieving arm pain in 92 to 100% of patients [3]. However, arm weakness and numbness may persist for weeks to months. Neck pain is relieved in 73 to 83% of patients [3]. In general, people with arm pain benefit more from ACDF than those with neck pain. Aim to keep a positive attitude and diligently perform your physical therapy exercises.

Achieving a spinal fusion varies depending on the technique used and your general health (smoker). In a study that compared three techniques: ACD, ACDF, and ACDF with plates and screws, the outcomes were [3]:

- 67% of people who underwent ACD (no bone graft) achieved fusion naturally. However, ACD alone results in an abnormal forward curving of the spine (kyphosis) compared with the other techniques.
- 93% of people who underwent ACDF with bone graft placement achieved fusion.
- 100% of people who underwent ACDF with bone graft placement and plates and screws achieved fusion.

# What are the risks?

No surgery is without risks. General complications of any surgery include bleeding, infection, blood clots (deep vein thrombosis), and reactions to anesthesia. If spinal fusion is done at the same time as a discectomy, there is a greater risk of complications. Specific complications related to ACDF may include:

**Hoarseness and swallowing difficulties.** In some cases, temporary hoarseness can occur. The recurrent laryngeal nerve, which controls the vocal cords, is affected during surgery. It may take several months for this nerve to recover. In rare cases (less than 1/250) hoarseness and swallowing problems may persist and need further treatment with an ear, nose and throat specialist.

**Vertebrae failing to fuse**. There are many reasons why bones do not fuse together. Common ones include smoking, osteoporosis, obesity, and malnutrition. Smoking is by far the greatest factor that can prevent fusion. Nicotine is a toxin that inhibits bone-growing cells. If you continue to smoke after your spinal surgery, you could undermine the fusion process.

**Hardware fracture**. Metal screws and plates used to stabilize the spine are called "hardware." The hardware may move or break before the bones are completely fused. If this occurs, a second surgery may be needed to fix or replace the hardware.

**Bone graft migration.** In rare cases (1 to 2%), the bone graft can move from the correct position between the vertebrae soon after surgery. This is more likely to occur if hardware (plates and screws) is not used or if multiple vertebral levels are fused. If this occurs, a second surgery may be necessary.

**Transitional syndrome**. Fusion of a spine segment causes extra stress and load to be transferred to the discs and bones above or below the fusion. The added wear and tear can eventually degenerate the adjacent level and cause pain.

**Nerve damage or persistent pain.** Any spine surgery comes with the risk of damaging the nerves or spinal cord. Damage can cause numbness or even paralysis. However, the most common cause of persistent pain is nerve damage from the disc herniation itself. Some disc herniations may permanently damage a nerve making it unresponsive to surgery. Like furniture on the carpet, the compressed nerve doesn't spring back. In these cases, <u>spinal cord stimulation</u> or other treatments may provide relief.

# Sources & links

If you have more questions, please contact the Mayfield Clinic at 800-325-7787 or 513-221-1100.

#### Sources

- 1. Bose B: Anterior cervical instrumentation enhances fusion rates in multilevel reconstruction in smokers. J Spinal Disord 14:3-9, 2001.
- 2. Hilibrand AS, et al.: Impact of smoking on the outcome of anterior cervical arthrodesis with interbody or strut-grafting. J Bone Joint Surg Am 83-A:668-73, 2001.
- 3. Xie JC, Hurlbert RJ. Discectomy versus discectomy with fusion versus discectomy with fusion and instrumentation: a prospective randomized study. Neurosurgery 61:107-16, 2007.

#### Links

www.spine-health.com www.spineuniverse.com www.knowyourback.org

# **Glossary**

**allograft:** a portion of living tissue taken from one person (the donor) and implanted in another (the recipient) for the purpose of fusing two tissues together.

autograft (autologous): a portion of living tissue taken from a part of ones own body and transferred to another for the purpose of fusing two tissues together.

**bone graft**: bone harvested from ones self (autograft) or from another (allograft) for the purpose of fusing or repairing a defect.

**discectomy:** a type of surgery in which herniated disc material is removed so that it no longer irritates and compresses the nerve root.

foraminotomy: surgical enlargement of the intervertebral foramen through which the spinal nerves pass from the spinal cord to the body.

fusion: to join together two separate bones into one to provide stability.

We comply with the HONcode standard for trustworthy health information: <u>verify here</u>.

interbody cage: a device made of titanium, carbon-fiber, or polyetheretherketone (PEEK) that is placed in the disc space between two vertebrae.

updated > 7.2013 reviewed by > <u>Robert Bohinski, MD</u>, Mayfield Clinic / University of Cincinnati Department of Neurosurgery, Cincinnati, Ohio



CERTIFIED 03/2013 *Mayfield Certified Health Info* materials are written and developed by the Mayfield Clinic & Spine Institute. We comply with the HONcode standard for trustworthy health information. This information is not intended to replace the medical advice of your health care provider.

Mayfield Clinic 506 Oak Street Cincinnati, Ohio 45219 phone: 513.221.1100 email: comments@MayfieldClinic.com



- Mayfield Chiari Center
- Mayfield Imaging Center
- Mayfield Education & Research Foundation
- Glia Media
- Precision Radiotherapy Center
- The Christ Hospital Spine Surgery Center
- University of Cincinnati Department of Neurosurgery

We invite you to join us on Social media:



Our blog: The Mayfield Brain & Spinal Column

Questions or comments? Please contact us

© 1998- 2015 Mayfield Clinic | All rights reserved permissions and licensing Privacy Policy | Disclaimer | Site Map