

If depression is holding you back from what you enjoy, and keeps you feeling sad, helpless and uninterested in your favorite activities,

Important Safety Information Prescribing Information

Important Safety Information About PRISTIQ

Suicidality and Antidepressant Drugs
Antidepressants increased the risk of suicidal thinking and behavior in children, teens, and young adults. Depression and certain other psychiatric disorders are themselves



Browse all medications [A](#) [B](#) [C](#) [D](#) [E](#) [F](#) [G](#) [H](#) [I](#) [J](#) [K](#) [L](#) [M](#) [N](#) [O](#) [P](#) [Q](#) [R](#) [S](#) [T](#) [U](#) [V](#) [W](#) [X](#) [Y](#) [Z](#) [Advanced Search](#) [Phonetic Search](#)

- Home → Drugs A to Z → Effexor
- Drugs A-Z ▾
- Pill Identifier
- Interactions Checker
- News ▾
- Health Professionals ▾
- Q & A ▾
- Mednotes
- Apps

Print Share

Sign In or Register

Effexor

Generic Name: [venlafaxine](#) (VEN la fax een)

Brand Names: *Effexor, Effexor XR*

- Overview
- Side Effects
- Dosage
- Interactions
- For Professionals
- More ▾

Quit Smoking Benefits

Visit to Learn More About The Benefits of Quitting Smoking



What is Effexor?

Effexor (venlafaxine) is an antidepressant in a group of drugs called selective serotonin and norepinephrine reuptake inhibitors (SSNRIs). Venlafaxine affects chemicals in the brain that may become unbalanced and cause depression.

Effexor is used to treat [major depressive disorder](#), [anxiety](#), and [panic disorder](#).

Important information

You should not take Effexor if you have uncontrolled narrow-angle glaucoma, or if you are being treated with methylene blue injection.

Do not use Effexor if you have taken an MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, phenelzine, rasagiline, selegiline, and tranylcypromine. After you stop taking this medicine, you must wait at least 7 days before you start taking an MAOI.

Some young people have thoughts about suicide when first taking an antidepressant. Your doctor will need to check your progress at regular visits while you are using Effexor. Your family or other caregivers should also be alert to changes in your mood or symptoms.

Report any new or worsening symptoms to

Related Information

- Rx** **Availability**
Prescription only
- C** **Pregnancy Category**
Risk cannot be ruled out
- N/A** **CSA Schedule**
Not a controlled drug
- Calendar** **Approval History**
Drug history at FDA

Manufacturer

Wyeth

Drug Class

[Serotonin-norepinephrine reuptake inhibitors \(SNRIs\)](#)

Related Drugs

Major Depressive Disorder
[trazodone](#), [Lexapro](#), [Zoloft](#), [Cymbalta](#), [sertraline](#), [Prozac](#), More...

Depression
[Xanax](#), [trazodone](#), [citalopram](#), [Lexapro](#), [Zoloft](#), [Cymbalta](#), More...

Bulimia
[Prozac](#), [fluoxetine](#), [venlafaxine](#), [Prozac Weekly](#), [Rapiflux](#), More...

Autism
[Abilify](#), [venlafaxine](#), [risperidone](#), [Risperdal](#), [Geodon](#), [Effexor XR](#), More...

▶ [1 more conditions...](#)

Reviews

[239 User Reviews](#)

Average User Rating

7.0 Rate it!

Effexor (venlafaxine) Images



ADVERTISEMENT

Important Safety Information Prescribing Information
Important Safety Information About PRISTIQ® (desvenlafaxine)

Suicidality and Antidepressant Drugs
Antidepressants increased the risk of suicidal thinking and behavior in children, teens, and young adults. Depression and certain other psychiatric disorders are themselves associated with increases in the risk of suicide. Patients of all ages who are started on antidepressant therapy or when the dose is changed should be monitored appropriately and observed closely for clinical worsening, suicidality, or unusual changes in behavior such as becoming agitated, irritable, hostile, aggressive, impulsive, or restless. Should these occur, report them to a doctor right away. PRISTIQ is not approved for use in children under 18.

Do not take PRISTIQ if you are allergic to desvenlafaxine.

ADVERTISEMENT

If depression is holding you back from what you enjoy, and keeps you feeling sad, helpless and uninterested in your favorite activities,

Prescribing Information
Important Safety Information
Important Safety Information About PRISTIQ

Suicidality and Antidepressant Drugs
Antidepressants increased the risk of suicidal thinking and behavior in children, teens, and young adults. Depression and certain other psychiatric disorders are themselves associated with increases in the risk of suicide.

05/05/2015

your doctor, such as: mood or behavior changes, anxiety, panic attacks, trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself.

Do not give this medicine to anyone younger than 18 years old without the advice of a doctor. Effexor is not approved for use in children.

Slideshow: Depression, the Risk of Suicide, and Treatment Options



Before taking this medicine

You should not take Effexor if you are allergic to venlafaxine or desvenlafaxine (Pristiq), or:

- if you have uncontrolled narrow-angle glaucoma; or
- if you are being treated with methylene blue injection.

Do not use Effexor if you have taken an MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, phenelzine, rasagiline, selegiline, and tranylcypromine. After you stop taking Effexor, you must wait at least 7 days before you start taking an MAOI.

To make sure Effexor is safe for you, tell your doctor if you have:

- bipolar disorder (manic depression);
- cirrhosis or other liver disease;
- kidney disease;
- heart disease, high blood pressure, high cholesterol;
- diabetes;
- narrow-angle glaucoma;
- a thyroid disorder;
- a history of seizures;
- a bleeding or blood clotting disorder;
- low levels of sodium in your blood; or
- if you are switching to Effexor from another antidepressant.

Some young people have thoughts about suicide when first taking an antidepressant. Your doctor will need to check your progress at regular visits while you are using Effexor. Your family or other caregivers should also be alert to changes in your mood or symptoms.

FDA pregnancy category C. Taking an SSRI antidepressant during pregnancy may cause serious lung problems or other complications in the baby. However, you may have a relapse of depression if you stop taking your antidepressant. Tell your doctor right away if you become pregnant while taking Effexor. **Do not start or stop taking this medicine during pregnancy without your doctor's advice.**

See also: [Pregnancy and breastfeeding warnings](#) (in more detail)

Venlafaxine can pass into breast milk and may harm a nursing baby. You should not breast-feed while using this medicine.

Do not give this medicine to anyone younger than 18 years old without the advice of a doctor. Effexor is not approved for use in children.

"My
Insomnia
Trick"

Top 5
Wrinkle
Creams

Quit
Smoking
w/



Effexor (venlafaxine) 75 mg



Effexor (venlafaxine) 50 mg

▶ [View all 8 images](#)

Get Updates by Email

Email me:

News and Warnings related to this drug

FDA Medwatch Alerts for all Medications

News Roundup Daily

Related News and Articles

[ADHD Tied to Higher Risk of Eating Disorder in Kids and Teens](#) May 1, 2015

Children with attention-deficit/hyperactivity disorder (ADHD) have an increased risk of a certain type of eating disorder, according to a new study. The...

[Want to Stay Slim? Keep Food Out of Sight](#)

April 30, 2015

With food, seeing it is eating it, a new study suggests. People who keep food in plain sight around the house are more likely to be obese, researchers report,...

[Bullying May Take Bigger Toll Than Child Abuse, Neglect](#) April 28, 2015

Being bullied as a child may take a larger toll on a young adult's mental health than being abused or neglected at home, a new study suggests. Kids who are...



Harvard Health Publications

Related: [Major Depression](#)

[zennight.co...](#)

Naturally Fall Asleep Fast, Stay Asleep All Night—Wake Up Refreshed

[iskincarere...](#)

The best anti-aging creams of 2015 reviewed by customers like you!

[Nicorette](#)[nicorette.com](#)

Double Your Chances of Quitting w/ Help From Nicorette. Learn More!



How should I take Effexor?

Take Effexor exactly as prescribed by your doctor. Follow all directions on your prescription label. Do not take this medicine in larger or smaller amounts or for longer than recommended.

Effexor should be taken with food. Try to take this medicine at the same time each day.

Swallow the controlled-release capsule (Effexor XR) whole, without crushing or chewing. To make the medication easier to swallow, you may open the capsule and sprinkle the medicine into a small amount of applesauce. Swallow all of the mixture without chewing, and do not save any for later use.

Your blood pressure will need to be checked often.

It may take 4 weeks before your symptoms improve. Keep using the medication as directed and tell your doctor if your symptoms do not improve.

You should not stop using Effexor suddenly. Follow your doctor's instructions about tapering your dose.

This medicine can cause you to have a false positive drug screening test. If you provide a urine sample for drug screening, tell the laboratory staff that you are taking Effexor.

Store at room temperature away from moisture and heat.

What happens if I miss a dose?

Take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not take extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking Effexor?

Ask your doctor before taking a nonsteroidal anti-inflammatory drug (NSAID) for pain, arthritis, fever, or swelling. This includes aspirin, ibuprofen (Advil, Motrin), naproxen (Aleve), celecoxib (Celebrex), diclofenac, indomethacin, meloxicam, and others. Using an NSAID with Effexor may cause you to bruise or bleed easily.

Avoid drinking alcohol, which can increase some of the side effects of Effexor.

Effexor may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be alert.

3 Herbs that Beat Anxiety

[a2xanxiety.com](#)

Doctors Reveal 1 Weird Compound to Calm Anxiety

2015 Best Skin Tightener

[shopskin.care...](#)

The One Thing Products To Tighten Loose, Sagging Face

3 Worst Foods for Thyroid

[medixselect.c...](#)

The One Thing You Should Be Eating For Your Thyroid

that May
Surprise YouSkin, Without
Plastic Surgery

Every Morning.



Effexor side effects

Get emergency medical help if you have any of these **signs of an allergic reaction to Effexor**: skin rash or hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety, panic attacks, trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself.

Call your doctor at once if you have:

- blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights;
- easy bruising;
- cough, chest tightness, trouble breathing;
- seizure (convulsions);
- high levels of serotonin in the body - agitation, hallucinations, fever, fast heart rate, overactive reflexes, nausea, vomiting, diarrhea, loss of coordination, fainting;
- low levels of sodium in the body - headache, confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady; or
- severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out.

Common Effexor side effects may include:

- vision changes;
- nausea, vomiting, diarrhea;
- changes in appetite or weight;
- dry mouth, yawning;
- dizziness, headache, anxiety, feeling nervous;
- fast heartbeats, tremors or shaking;
- sleep problems (insomnia), strange dreams, tired feeling;
- increased sweating; or
- decreased sex drive, impotence, or difficulty having an orgasm.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

See also: [Side effects](#) (in more detail)

Effexor dosing information

Usual Adult Dose for Depression:

Immediate release:

Initial dose: 37.5 mg orally twice a day or 25 mg orally 3 times a day

Maintenance dose: May increase in daily increments of up to 75 mg at intervals of no less than 4 days

Maximum dose: (moderately depressed outpatients): 225 mg/day

Maximum dose (severely depressed inpatients): 375 mg/day

Daily dosage may be divided in 2 or 3 doses/day

Extended release:

Initial dose: 75 mg orally once daily

Maintenance dose: May increase in daily increments of up to 75 mg at intervals of no less than 4 days

Maximum dose (moderately depressed outpatients): 225 mg/day

Maximum dose (severely depressed inpatients): 375 mg/day

Usual Adult Dose for Anxiety:

For generalized anxiety disorder or social anxiety disorder:

Extended release:

Initial dose: 75 mg orally once daily

Maintenance dose: May increase in daily increments of 75 mg at intervals of no less than 4 days

Maximum dose: 225 mg/day

Usual Adult Dose for Panic Disorder:

Extended-release:

Initial dose: 37.5 mg once a day

Maintenance dose: May increase dose in daily increments of 75 mg at intervals of no less than 7 days

Maximum dose: 225 mg/day






What other drugs will affect Effexor?

Taking this medicine with other drugs that make you sleepy can worsen this effect. Ask your doctor before taking Effexor with a sleeping pill, narcotic pain medicine, muscle relaxer, or medicine for anxiety, depression, or seizures.

Many drugs can interact with venlafaxine. Not all possible interactions are listed here. Tell your doctor about all your medications and any you start or stop using during treatment with Effexor, especially:

- any other antidepressant;
- cimetidine;
- St. John's wort;
- tramadol;
- tryptophan (sometimes called L-tryptophan);
- a blood thinner - warfarin, Coumadin, Jantoven;
- medicine to treat mood disorders, thought disorders, or mental illness - buspirone, lithium, and many others; or
- migraine headache medicine - sumatriptan, zolmitriptan, and others.

This list is not complete and many other drugs can interact with venlafaxine. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Give a list of all your medicines to any healthcare provider who treats you.

 Google  Facebook  Twitter  Print  Email  Add to Mednotes

Next Page → [Side Effects](#)

More about Effexor (venlafaxine)

- [Side Effects](#)
- [During Pregnancy](#) or [Breastfeeding](#)
- [Dosage Information](#)
- [Drug Images](#)
- [Drug Interactions](#)
- [Support Group](#)
- [En Espanol](#)
- [239 Reviews](#) - [Add your own review/rating](#)

Consumer resources

- [Effexor](#)

- [Effexor](#) (Advanced Reading)

Professional resources

- [Effexor](#) (FDA)
- [Venlafaxine Hydrochloride](#) (AHFS Monograph)

Other formulations

- [Effexor XR](#)

Related treatment guides

- [Major Depressive Disorder](#)
- [Depression](#)
- [Bulimia](#)
- [Autism](#)
- [Postpartum Depression](#)

Where can I get more information?

- Your pharmacist can provide more information about Effexor.

Holistic Addiction Rehab

Alternative to 12-step program Holistic Biochemical
Restoration



Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Effexor only for the indication prescribed.

Disclaimer: Every effort has been made to ensure that the information provided by Cerner Multum, Inc. ('Multum') is accurate, up-to-date, and complete, but no guarantee is made to that effect. Drug information contained herein may be time sensitive. Multum information has been compiled for use by healthcare practitioners and consumers in the United States and therefore Multum does not warrant that uses outside of the United States are appropriate, unless specifically indicated otherwise. Multum's drug information does not endorse drugs, diagnose patients or recommend therapy. Multum's drug information is an informational resource designed to assist licensed healthcare practitioners in caring for their patients and/or to serve consumers viewing this service as a supplement to, and not a substitute for, the expertise, skill, knowledge and judgment of healthcare practitioners. The absence of a warning for a given drug or drug combination in no way should be construed to indicate that the drug or drug combination is safe, effective or appropriate for any given patient. Multum does not assume any responsibility for any aspect of healthcare administered with the aid of information Multum provides. The information contained herein is not intended to cover all possible uses, directions, precautions, warnings, drug interactions, allergic reactions, or adverse effects. If you have questions about the drugs you are taking, check with your doctor, nurse or pharmacist.

Copyright 1996-2015 Cerner Multum, Inc. Version: 13.03. Revision Date: 2014-08-23, 9:34:03 AM.



Support

- [Advertise With Us](#)
- [Frequently Asked Questions](#)
- [Ask a Question](#)
- [Content Submissions](#)
- [Mobile Apps](#)
- [About Drugs.com](#)
- [Contact Us](#)
- [Sitemap](#)

Follow: [g+](#) [f](#) [t](#) [You Tube](#) [RSS](#)

About Drugs.com

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment.

Data sources include Micromedex® (updated Apr 27th, 2015), Cerner Multum™ (updated Apr 20th, 2015), Wolters Kluwer™ (updated Apr 6th, 2015) and others. To view content sources and attributions, refer to our [editorial policy](#).

Copyright © 2000-2015 Drugs.com. All rights reserved.

Your Privacy



We comply with the HONcode standard for trustworthy health information.

[Verify here](#)



Ad Choices

[Terms of Use](#) · [Privacy Policy](#)