

Browse all medications [A](#) [B](#) [C](#) [D](#) [E](#) [F](#) [G](#) [H](#) [I](#) [J](#) [K](#) [L](#) [M](#) [N](#) [O](#) [P](#) [Q](#) [R](#) [S](#) [T](#) [U](#) [V](#) [W](#) [X](#) [Y](#) [Z](#) [Advanced Search](#) [Phonetic Search](#)

Drugs A-Z ▾

Pill Identifier

Interactions Checker

News ▾

Health Professionals ▾

Q & A ▾

Mednotes

Apps

Home → Conditions → Fibromyalgia → Cymbalta

Print
 Share

[Sign In](#) or [Register](#)

Cymbalta

Generic Name: [duloxetine](#) (du LOX e [teen](#))Brand Names: *Cymbalta*

Overview

Side Effects

Dosage

Interactions

For Professionals

More ▾

1 Trick to Fibromyalgia

See How to Relieve Fibromyalgia With This Fast and Easy Trick



What is Cymbalta?

Cymbalta (duloxetine) is a selective serotonin and norepinephrine reuptake inhibitor antidepressant (SSNRI). Duloxetine affects chemicals in the brain that may become unbalanced and cause depression.

Cymbalta is used to treat [major depressive disorder](#) in adults. It is also used to treat [general anxiety disorder](#) in adults and children who are at least 7 years old.

Cymbalta is also used in adults to treat [fibromyalgia](#) (a chronic pain disorder), or chronic muscle or joint pain (such as low back pain and osteoarthritis pain).

Cymbalta is also used to treat pain caused by nerve damage in adults with diabetes ([diabetic neuropathy](#)).

Important information

You should not use Cymbalta if you have untreated or uncontrolled glaucoma, or if you also take thioridazine.

Do not use Cymbalta if you have used an MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, methylene blue injection, phenelzine, rasagiline, selegiline, tranylcypromine, and others. After you stop taking Cymbalta, you must wait at least 5 days before you start taking an MAOI.

Some young people have thoughts about suicide when first taking an antidepressant. Your doctor will need to check your progress at regular visits while you are using Cymbalta. [Your family](#) or other caregivers should also

Slideshow: Top 10 Money-Making Drugs of 2012

Related Information



Availability
Prescription only



Pregnancy Category
Risk cannot be ruled out



CSA Schedule
Not a controlled drug



Approval History
Drug history at FDA

Manufacturer

Eli Lilly and Company

Drug Class

Serotonin-norepinephrine reuptake inhibitors (SNRIs)

Related Drugs

Fibromyalgia

[prednisone](#), [Lyrica](#), [amitriptyline](#), [duloxetine](#), [Elavil](#), [pregabalin](#), More...

Chronic Pain

[tramadol](#), [oxycodone](#), [Percocet](#), [fentanyl](#), [morphine](#), More...

Anxiety

[Xanax](#), [Lexapro](#), [alprazolam](#), [lorazepam](#), [hydroxyzine](#), [atenolol](#), More...

Back Pain

[tramadol](#), [naproxen](#), [aspirin](#), [ibuprofen](#), [Norco](#), More...

[5 more conditions...](#)

Reviews

809 User Reviews

Average User Rating

7.0 [Rate it!](#)

Cymbalta (duloxetine) Images



05/05/2015

be alert to changes in your mood or symptoms.

Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety, panic attacks, trouble [sleeping](#), or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself.



Cymbalta (duloxetine) 30 mg



Cymbalta (duloxetine) 60 mg

[View all 4 images](#)

Before taking this medicine

You should not use this Cymbalta if you are allergic to duloxetine, or if you have untreated or uncontrolled glaucoma.

Do not use Cymbalta if you have used an MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, methylene blue injection, phenelzine, rasagiline, selegiline, tranylcypromine, and others. After you stop taking Cymbalta, you must wait at least 5 days before you start taking an MAOI.

To make sure Cymbalta is safe for you, tell your doctor if you have:

- liver or kidney disease;
- seizures or epilepsy;
- a bleeding or blood clotting disorder;
- high blood pressure;
- narrow-angle glaucoma;
- bipolar disorder (manic depression); or
- a history of drug abuse or suicidal thoughts.

Some young people have thoughts about suicide when first taking an antidepressant. Your doctor will need to check your progress at regular visits while you are using Cymbalta. Your [family](#) or other caregivers should also be alert to changes in your mood or symptoms.

FDA pregnancy category C. It is not known whether Cymbalta will harm an unborn baby. However, Cymbalta may cause problems in a newborn if you take the medicine during the third trimester of pregnancy. Tell your doctor if you are pregnant or plan to become pregnant while using this medicine.

See also: [Pregnancy and breastfeeding warnings](#) (in more detail)

If you are pregnant, your name may be listed on a pregnancy registry. This is to track the outcome of the pregnancy and to evaluate any effects of Cymbalta on the baby.

Duloxetine can pass into breast milk and may harm a nursing baby. You should not breast-feed while taking this medicine.

Do not give this medicine to anyone younger than 18 years old without the advice of a doctor.

3 Herbs that Beat Anxiety

a2xanxiety.com

Doctors Reveal 1 Weird Compound to Calm Anxiety that May Surprise You



Reduce Nerve Pain Now

[neuropathytreat...](#)

Don't Let Neuropathy Symptoms Slow You Down. Get A Free Trial Now!



"My Insomnia Trick"

zennight.com/I...

Naturally Fall Asleep Fast, Stay Asleep All Night—Wake Up Refreshed

Get Updates by Email

Email me:

News and Warnings related to this drug

FDA Medwatch Alerts for all Medications

News Roundup Daily

Related News and Articles

[Want to Stay Slim? Keep Food Out of Sight](#)

April 30, 2015

With food, seeing it is eating it, a new study suggests. People who keep food in plain sight around the house are more likely to be obese, researchers report,...

[Bullying May Take Bigger Toll Than Child Abuse, Neglect](#)

April 28, 2015

Being bullied as a child may take a larger toll on a young adult's mental health than being abused or neglected at home, a new study suggests. Kids who are...

[Depression, Weapons May Be More Common for Bullied Teens](#)

April 27, 2015

Bullied high school students have greater odds for depression and suicidal thoughts than others, and they're also more likely to take weapons to school,...



Harvard Health Publications

Related: [Fibromyalgia](#)



How should I take Cymbalta?

Take Cymbalta exactly as prescribed by your doctor. Follow all directions on your prescription label. Do not take this medicine in larger or smaller amounts or for longer than recommended.

Try to take the medicine at the same time each day. Follow the directions on your prescription label.

Do not crush, chew, break, or open an extended-release capsule. Swallow it whole.

It may take 4 weeks or longer before your symptoms improve. Keep using the medication as directed. Do not stop using Cymbalta without first talking to your doctor. You may have unpleasant side effects if you stop taking this medicine suddenly.

Store at room temperature away from moisture and heat.

What happens if I miss a dose?

Take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not take extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking Cymbalta?

Avoid drinking alcohol. It may increase your risk of liver damage.

Ask your doctor before taking a nonsteroidal anti-inflammatory drug (NSAID) for pain, arthritis, fever, or swelling. This includes aspirin, ibuprofen (Advil, Motrin), naproxen (Aleve), celecoxib (Celebrex), diclofenac, indomethacin, meloxicam, and others. Using an NSAID with Cymbalta may cause you to bruise or bleed easily.

Cymbalta may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be alert. Avoid getting up too fast from a sitting or lying position, or you may feel dizzy. Get up slowly and steady yourself to prevent a fall. Severe dizziness or fainting can cause falls, accidents, or severe injuries.

2015 Best Skin Tightener

S

shopskin.care/Fa...
Products To Tighten Loose, Sagging Face Skin, Without Plastic Surgery



Apply for Disability

disability-benefi...

Are you eligible for up to \$2642/mo in Social Security Disability?



New Depression Therapy

fisherwallace.com

The Fisher Wallace Stimulator@. Cleared by the FDA & Health Canada.



Cymbalta side effects

Get emergency medical help if you have any of these **signs of an allergic reaction to Cymbalta**: skin rash or hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety, panic attacks, trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts

about suicide or hurting yourself.

Call your doctor at once if you have:

- a light-headed feeling, like you might pass out;
- blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights;
- easy bruising, unusual bleeding;
- painful or difficult urination;
- liver problems - nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes);
- high levels of serotonin in the body - agitation, hallucinations, fever, fast heart rate, overactive reflexes, nausea, vomiting, diarrhea, loss of coordination, fainting;
- low levels of sodium in the body - headache, confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady;
- severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors; or
- severe skin reaction - fever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain, followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling.

Older adults may be more sensitive to the side effects of this medicine.

Common Cymbalta side effects may include:

- vision changes;
- dry mouth;
- drowsiness, dizziness;
- tired feeling;
- loss of appetite; or
- increased sweating.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

See also: [Side effects](#) (in more detail)

What other drugs will affect Cymbalta?

Taking this medicine with other drugs that make you sleepy can worsen this effect. Ask your doctor before taking Cymbalta with a sleeping pill, narcotic pain medicine, muscle relaxer, or medicine for anxiety, depression, or seizures.

Many drugs can interact with duloxetine. Not all possible interactions are listed here. Tell your doctor about all your medications and any you start or stop using during treatment with Cymbalta, especially:

- any other antidepressant;
- cimetidine;
- St. John's wort;
- theophylline;
- tryptophan (sometimes called L-tryptophan);
- an antibiotic - ciprofloxacin, enoxacin;
- a blood thinner - warfarin, Coumadin, Jantoven;
- heart rhythm medication - flecainide, propafenone, quinidine, and others;
- narcotic pain medicine - fentanyl, tramadol;
- medicine to treat mood disorders, thought disorders, or mental illness - buspirone, lithium, and many others; or
- migraine headache medicine - sumatriptan, rizatriptan, zolmitriptan, and others.

This list is not complete and many other drugs can interact with duloxetine. This includes

prescription and over-the-counter medicines, vitamins, and herbal products. Give a list of all your medicines to any healthcare provider who treats you.

 Google  Facebook  Twitter  Print  Email  Add to Mednotes

Next Page → [Side Effects](#)

More about Cymbalta (duloxetine)

- [Side Effects](#)
- [During Pregnancy](#) or [Breastfeeding](#)
- [Dosage Information](#)
- [Drug Images](#)
- [Drug Interactions](#)
- [Support Group](#)
- [Pricing & Coupons](#)
- [En Espanol](#)
- [809 Reviews](#) - [Add your own review/rating](#)
- [Generic Availability](#)

Consumer resources

- [Cymbalta delayed-release capsules](#)
- [Cymbalta](#) (Advanced Reading)

Professional resources

- [Cymbalta](#) (AHFS Monograph)
- [Cymbalta](#) (FDA)

Related treatment guides

- [Fibromyalgia](#)
- [Chronic Pain](#)
- [Anxiety](#)
- [Back Pain](#)
- [Osteoarthritis](#)
- [More \(4\) »](#)

Where can I get more information?

- Your pharmacist can provide more information about Cymbalta.

Apply For Disability

Find Out If You May Qualify For Up To \$2,663/mo in Disability

Benefit



Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Disclaimer: Every effort has been made to ensure that the information provided by Cerner Multum, Inc. ('Multum') is accurate, up-to-date, and complete, but no guarantee is made to that effect. Drug information contained herein may be time sensitive. Multum information has been compiled for use by healthcare practitioners and consumers in the United States and therefore Multum does not warrant that uses outside of the United States are appropriate, unless specifically indicated otherwise. Multum's drug information does not endorse drugs, diagnose patients or recommend therapy. Multum's drug information is an informational resource designed to assist licensed healthcare practitioners in caring for their patients and/or to serve consumers viewing this service as a supplement to, and not a substitute for, the expertise, skill, knowledge and judgment of healthcare practitioners. The absence of a warning for a given drug or drug combination in no way should be construed to indicate that the drug or drug combination is safe, effective or appropriate for any given patient. Multum does not assume any responsibility for any aspect of healthcare administered with the aid of information Multum provides. The information contained herein is not intended to cover all possible uses, directions, precautions, warnings, drug interactions, allergic reactions, or adverse effects. If you have questions about the drugs you are taking, check with your doctor, nurse or pharmacist.

Copyright 1996-2015 Cerner Multum, Inc. Version: 10.03. Revision Date: 2014-12-15, 9:43:27 AM.

**Support**

[Advertise With Us](#)
[Frequently Asked Questions](#)
[Ask a Question](#)
[Content Submissions](#)
[Mobile Apps](#)
[About Drugs.com](#)
[Contact Us](#)
[Sitemap](#)

Follow:     

About Drugs.com

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment.

Data sources include Micromedex® (updated Apr 27th, 2015), Cerner Multum™ (updated Apr 20th, 2015), Wolters Kluwer™ (updated Apr 6th, 2015) and others. To view content sources and attributions, refer to our [editorial policy](#).

Copyright © 2000-2015 Drugs.com. All rights reserved.

Your Privacy

We comply with the HONcode standard for trustworthy health information.
[Verify here](#)



Ad Choices

[Terms of Use](#) · [Privacy Policy](#)