



Substance Abuse Cured

Utah Rehab Facility in Resort Area Exclusive treatment, affordable



ADVERTISEMENT

Home › Conditions › Pain › Percocet

Print Share

Percocet

Generic Name: [acetaminophen and oxycodone](#) (a SEET a MIN oh fen and OX i KOE done)

Brand Names: *Endocet, Percocet, Primlev, Roxicet, Xartemis XR*

Bulging Disc Pain Relief

New, minimally invasive procedure for bulging disc pain, learn more



Overview

Side Effects

Dosage

Interactions

Professional

More

What is Percocet?

Percocet contains a combination of acetaminophen and oxycodone. Oxycodone is an opioid pain medication. An opioid is sometimes called a narcotic. Acetaminophen is a less potent pain reliever that increases the effects of oxycodone.


Percocet is used to relieve moderate to severe pain.

Percocet may also be used for purposes not listed in this medication guide.

ADVERTISEMENT



07/09/2015



#1 LIE About Blood Pressure Meds

Here's the ugly truth your doctor will NEVER tell you about blood pressure medication. [Read More](#)

Important information

You should not use Percocet if you have recently used alcohol, sedatives, tranquilizers, or other narcotic medications.

Oxycodone can slow or stop your breathing. Never use Percocet in larger amounts, or for longer than prescribed. Narcotic pain medicine may be habit-forming, even at regular doses. Never share this medicine with another person, especially someone with a history of drug abuse or addiction. Keep the medication in a place where others cannot get to it. **MISUSE OF NARCOTIC MEDICATION CAN CAUSE ADDICTION, OVERDOSE, OR DEATH, especially in a child or other person using the medicine without a prescription.**

Do not use Percocet if you have taken an MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, phenelzine, rasagiline, selegiline, and tranylcypromine.

Do not take more of this medication than is recommended. An overdose of acetaminophen can damage your liver or cause death. Call your doctor at once if you have nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, or jaundice (yellowing of your skin or eyes).

In rare cases, acetaminophen may cause a severe skin reaction. Stop taking Percocet and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling.

Slideshow: 2013 Drug News Round-Up - Top 20 Stories



Before taking this medicine

07/09/2015

You should not use Percocet if you are allergic to acetaminophen (Tylenol) or oxycodone, or if you have recently used alcohol, sedatives, tranquilizers, or other narcotic medications.

Do not use Percocet if you have taken an MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, phenelzine, rasagiline, selegiline, and tranylcypromine.

To make sure Percocet is safe for you, tell your doctor if you have:

- liver disease, cirrhosis, or if you drink more than 3 alcoholic beverages per day;
- a history of alcoholism or drug addiction;
- diarrhea, inflammatory bowel disease;
- bowel obstruction, severe constipation;
- a colostomy or ileostomy;
- kidney disease;
- low blood pressure, or if you are dehydrated;
- a history of head injury, brain tumor, or stroke; or
- asthma, COPD, sleep apnea, or other breathing disorders.

This medicine is more likely to cause breathing problems in [older](#) adults and people who are severely ill, malnourished, or otherwise debilitated.

Oxycodone may be habit forming. Never share Percocet with another person, especially someone with a history of drug abuse or addiction. Keep the medication in a place where others cannot get to it. Selling or giving away Percocet to any other person is against the law.

FDA pregnancy category C. It is not known whether this medication is harmful to an unborn baby, but it could cause breathing problems or addiction/withdrawal symptoms in a newborn. Before you take Percocet, tell your doctor if you are [pregnant](#) or plan to become pregnant during [treatment](#).

See also: [Pregnancy and breastfeeding warnings](#) (in more detail)

Acetaminophen and oxycodone may pass into breast milk and could harm a [nursing](#) baby. Tell your doctor if you are breast-feeding a baby.

Bulging Disc Pain Relief



northamericanspine.com/Bulging_Disc

New, minimally invasive procedure for bulging disc pain, learn more.

07/09/2015

[Need Withdrawal Relief?](#)

[Knee Pain Injections](#)

How should I take Percocet?

Take Percocet exactly as prescribed. Follow all directions on your prescription label. Never take Percocet in larger amounts, or for longer than prescribed. An overdose can damage your liver or cause death. Tell your doctor if the medicine seems to stop working as well in relieving your pain.

Drink 6 to 8 full glasses of water daily to help prevent constipation while you are taking Percocet. Do not use a stool softener (laxative) without first asking your doctor.

Percocet can cause unusual results with certain urine tests. Tell any doctor who treats you that you are using Percocet.

If you need surgery, tell the surgeon ahead of time that you are using acetaminophen and oxycodone. You may need to stop using the medicine for a short time.

Do not stop using Percocet suddenly after long-term use, or you could have unpleasant withdrawal symptoms. Ask your doctor how to avoid withdrawal symptoms when you stop using this medicine.

Store Percocet at room temperature away from moisture and heat.

Keep track of the amount of medicine used from each new bottle. Oxycodone is a drug of abuse and you should be aware if anyone is using your medicine improperly or without a prescription.

Always check your bottle to make sure you have received the correct pills (same brand and type) of medicine prescribed by your doctor. Ask the pharmacist if you have any questions about the medicine you receive at the pharmacy.

Do not keep leftover Percocet tablets. Ask your pharmacist where to locate a drug take-back disposal program. If there is no take-back program, flush any unused tablets down the toilet. Disposal of medicines by flushing is recommended to reduce the danger of accidental overdose causing death. This advice applies to a very small number of medicines only. The FDA, working with the manufacturer, has determined this method to be the most appropriate route of disposal and presents the least risk to human safety.

What happens if I miss a dose?

Since Percocet is taken as needed, you may not be on a dosing schedule. If you are taking the medication regularly, take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not use extra medicine to make up the missed dose.

07/09/2015

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. An overdose of Percocet can be fatal.

The first signs of an acetaminophen overdose include loss of appetite, nausea, vomiting, stomach pain, sweating, and confusion or weakness. Later symptoms may include pain in your upper stomach, dark urine, and yellowing of your skin or the whites of your eyes.

Overdose symptoms may also include extreme drowsiness, pinpoint pupils, cold and clammy skin, muscle weakness, fainting, weak pulse, slow heart rate, coma, blue lips, shallow breathing, or no breathing

What should I avoid?

This medication may impair your thinking or reactions. Avoid driving or operating machinery until you know how Percocet will affect you. Dizziness or severe drowsiness can cause falls or other accidents.

Ask a doctor or pharmacist before using any other cold, allergy, pain, or sleep medication.

Acetaminophen (sometimes abbreviated as APAP) is contained in many combination medicines. Taking certain products together can cause you to get too much acetaminophen which can lead to a fatal overdose. Check the label to see if a medicine contains acetaminophen or APAP.

Avoid drinking alcohol. It may increase your risk of liver damage while taking acetaminophen.

Need Withdrawal Relief?



 [withdrawal-ease.com](http://www.withdrawal-ease.com)

#1 Painkiller Withdrawal Remedy. Home Remedy for Opiate Withdrawal

[Knee Pain Injections](#) ▾

[New Sciatica Treatment](#) ▾

Percocet side effects

Get emergency medical help if you have any signs of an allergic reaction to Percocet: hives; difficulty

07/09/2015

breathing; swelling of your face, lips, tongue, or throat.

In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking Percocet and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen.

Call your doctor at once if you have:

- shallow breathing, slow heartbeat;
- a light-headed feeling, like you might pass out;
- confusion, unusual thoughts or behavior;
- seizure (convulsions);
- problems with urination; or
- nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes).

Common Percocet side effects include:

- drowsiness;
- upset stomach, constipation;
- blurred vision; or
- dry mouth.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

See also: [Side effects](#) (in more detail)

What other drugs will affect Percocet?

Taking this medicine with other drugs that make you sleepy or slow your breathing can cause dangerous or life-threatening side effects. Ask your doctor before taking Percocet with a sleeping pill, narcotic pain medicine, muscle relaxer, or medicine for anxiety, depression, or seizures.

Other drugs may interact with acetaminophen and oxycodone, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell each of your health care providers about all medicines you use now and any medicine you start or stop using.

Next → [Side Effects](#)



More about Percocet (acetaminophen / oxycodone)

- [Side Effects](#)
- [During Pregnancy](#)
- [Dosage Information](#)
- [Drug Images](#)
- [Drug Interactions](#)
- [Support Group](#)
- [173 Reviews - Add your own review/rating](#)

Consumer resources

- [Percocet](#)
- [Percocet \(Advanced Reading\)](#)
- *Other brands: [Endocet](#), [Roxicet](#), [Tylox](#), [Xartemis XR](#), [More \(6\)](#) »*

Professional resources

- [Percocet \(FDA\)](#)

Other formulations

- [Percocet 5/325](#)
- [Percocet 10/325](#)
- [Percocet 7.5/325](#)
- [More \(3\)](#) »

Related treatment guides

- [Chronic Pain](#)
- [Pain](#)

07/09/2015

Where can I get more information?

- Your pharmacist can provide more information about Percocet.

Substance Abuse Cured

Utah Rehab Facility in Resort Area Exclusive treatment, affordable



Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Percocet only for the indication prescribed.

Disclaimer: Every effort has been made to ensure that the information provided by Cerner Multum, Inc. ("Multum") is accurate, up-to-date, and complete, but no guarantee is made to that effect. Drug information contained herein may be time sensitive. Multum information has been compiled for use by healthcare practitioners and consumers in the United States and therefore Multum does not warrant that uses outside of the United States are appropriate, unless specifically indicated otherwise. Multum's drug information does not endorse drugs, diagnose patients or recommend therapy. Multum's drug information is an informational resource designed to assist licensed healthcare practitioners in caring for their patients and/or to serve consumers viewing this service as a supplement to, and not a substitute for, the expertise, skill, knowledge and judgment of healthcare practitioners. The absence of a warning for a given drug or drug combination in no way should be construed to indicate that the drug or drug combination is safe, effective or appropriate for any given patient. Multum does not assume any responsibility for any aspect of healthcare administered with the aid of information Multum provides. The information contained herein is not intended to cover all possible uses, directions, precautions, warnings, drug interactions, allergic reactions, or adverse effects. If you have questions about the drugs you are taking, check with your doctor, nurse or pharmacist.

Copyright 1996-2015 Cerner Multum, Inc. Version: 15.08. Revision Date: 2015-03-11, 10:25:47 AM.

DRUG STATUS



Availability
Prescription only



Pregnancy Category
Risk cannot be ruled out



CSA Schedule
High potential for abuse



Approval History
Drug history at FDA



WADA Class
Anti-Doping Classification



Manufacturer

[Endo](#)
[Pharmaceuticals](#)

ADVERTISEMENT

07/09/2015

Drug Class

[Narcotic analgesic combinations](#)

Related Drugs

Chronic Pain

[tramadol](#), [oxycodone](#),
[Cymbalta](#), [fentanyl](#),
[morphine](#), [duloxetine](#),
[Dilaudid](#), [methadone](#),
[OxyContin](#), [Ultram](#),
[hydromorphone](#),

[More...](#)

Pain

[tramadol](#), [oxycodone](#),
[naproxen](#),
[acetaminophen](#),
[Tylenol](#), [aspirin](#),
[hydrocodone](#),
[amitriptyline](#),
[ibuprofen](#),
[hydroxyzine](#),
[clonidine](#), [More...](#)

Percocet Rating

[173 User Reviews](#)

8.7/10

Percocet Images



Percocet (acetaminophen / oxycodone) 325 mg / 5 mg

07/09/2015



[View larger images](#)

Get Updates by Email

Sign up for daily, weekly or monthly emails to receive news updates to your inbox!

News and Warnings related to this drug

FDA Medwatch Alerts for all Medications

News Roundup **Daily**

Related News and Articles

Painkiller Overdoses Often Involve 'Pharmacy Shopping'

July 8, 2015

Who's Most Likely to Get Addicted to Their Narcotic Painkiller?

July 7, 2015

Many Doctors Underestimate Risks of Prescription Painkillers: Survey

June 26, 2015



Percocet

ADVERTISEMENT

 [Pronunciation](#)

Generic Name: [acetaminophen and](#)

Drugs.com Mobile Apps

The easiest way to lookup drug information, identify pills, check interactions and set up your own personal medication records. Available for Android and iOS devices.

[Explore Apps](#)



SUPPORT

- [Help Center](#)
- [Frequent Questions](#)
- [Sitemap](#)
- [Contact Us](#)

ABOUT

- [About Drugs.com](#)
- [Advertising Policy](#)
- [Content Submissions](#)
- [Drugs.com Blog](#)

TERMS & PRIVACY


- [Editorial Policy](#)
- [Privacy Policy](#)
- [Terms of Use](#)
- [Attribution & Citations](#)

CONNECT



[Subscribe](#) to receive email notifications whenever new articles are published.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex® (updated May 8th, 2015), Cerner Multum™ (updated May 18th, 2015), Wolters Kluwer™ (updated May 6th, 2015) and others. To view content sources and attributions, please refer to our [editorial policy](#).



We comply with the HONcode standard for trustworthy health information - [verify here](#)

07/09/2015

Certified Privacy

CERTIFIED
10/2013

Copyright © 2000-2015 Drugs.com. All rights reserved.