


Browse all medications [A](#) [B](#) [C](#) [D](#) [E](#) [F](#) [G](#) [H](#) [I](#) [J](#) [K](#) [L](#) [M](#) [N](#) [O](#) [P](#) [Q](#) [R](#) [S](#) [T](#) [U](#) [V](#) [W](#) [X](#) [Y](#) [Z](#) [Advanced Search](#) [Phonetic Search](#)



Home → Drugs A to Z → Lidoderm


[Print](#) [Share](#)

[Sign In](#) or [Register](#)

Lidoderm

Generic Name: [lidocaine topical](#) (LYE doe kane TOP i kal)

Brand Names: *AneCream*, *AneCream with Tegaderm*, *Anestacon*, *Bactine*, *LidaMantle*, *Lidocream*, *Lidoderm*, *Lidosense5*, *LMX 4*, *LMX 4 with Tegaderm*, *LMX 5*, *Medi-Quik Spray*, *Senatec*, *Xylocaine Jelly*, *Xylocaine Topical*, *Zilactin-L*



Chronic Back Pain Relief

One groundbreaking, 30-minute procedure may relieve back pain.



What is Lidoderm?

Lidoderm is a local anesthetic (numbing [medication](#)). It works by blocking nerve signals in your body.

Lidoderm is used to relieve post-shingles pain. Apply Lidoderm only to intact [skin](#) with no blisters.

Lidocaine topical in other formulations is used to reduce pain or discomfort caused by skin irritations such as sunburn, insect bites, poison ivy, poison oak, poison sumac, and minor cuts, scratches, hemorrhoids, and burns. Lidocaine topical can also be used to treat sores inside the mouth, during [dental](#) procedures to numb the gums, and to numb the skin for a medical procedure (such as getting stitches).



Important information

An overdose of Lidoderm can cause fatal side effects if too much lidocaine is absorbed through your skin and into your blood. This is more likely to occur when using a numbing medicine without the advice of a medical doctor (such as during a cosmetic procedure like laser hair removal). Overdose [symptoms](#) may include uneven heartbeats, seizure (convulsions), coma, slowed breathing, or respiratory failure (breathing stops). Your body may absorb more lidocaine if you use too much, if you apply it over large skin areas, or if you apply heat, bandages, or plastic wrap to treated skin areas. Skin that is cut or irritated may also absorb more topical medication than healthy skin.

Use the smallest amount of Lidoderm needed to numb the skin or relieve pain. Do not use

Slideshow: Hemorrhoids: A Common Ailment with Frequent Questions

Related Information



Availability
Rx and/or OTC



Pregnancy Category
No proven risk in humans



CSA Schedule
Not a controlled drug



Approval History
Drug history at FDA

Drug Class

ADVERTISEMENT

[Topical anesthetics](#)

Related Drugs

Pain

[tramadol](#), [oxycodone](#), [naproxen](#), [Tylenol](#), [acetaminophen](#), [More...](#)

Hemorrhoids

[lidocaine topical](#), [hydrocortisone topical](#), [phenylephrine topical](#), [Anusol](#), [benzocaine topical](#), [Anusol-HC](#), [More...](#)

Anal Itching

[lidocaine topical](#), [hydrocortisone topical](#), [phenylephrine topical](#), [benzocaine topical](#), [Anusol-HC](#), [Proctosol-HC](#), [More...](#)

Anesthesia

[fentanyl](#), [lidocaine](#), [hyoscyamine](#), [ketamine](#), [propofol](#), [More...](#)

[4 more conditions...](#)

Chronic Back Pain Relief

One groundbreaking, 30-minute procedure may relieve back pain.



Reviews

[22 User Reviews](#)

Average User Rating

7.9 [Rate it!](#)

Get Updates by Email

05/06/2015

large amounts of Lidoderm, or cover treated skin areas with a bandage or plastic wrap without medical advice. Be aware that many cosmetic procedures are performed without a medical doctor present.

Before using Lidoderm

An overdose of numbing medications can cause fatal side effects if too much of the medicine is absorbed through your skin and into your blood.

Overdose is more likely to occur when using a numbing medicine without the advice of a medical doctor (such as during a cosmetic procedure like laser hair removal). Symptoms may include uneven heartbeats, seizure (convulsions), coma, slowed breathing, or respiratory failure (breathing stops).

Do not use Lidoderm if you are allergic to any other type of numbing medicine.

Before using Lidoderm, tell your doctor if you are allergic to any drugs, or if you have liver disease, or broken, swollen, or damaged skin. You may not be able to use Lidoderm, or you may need dosage adjustments or special tests during treatment.

FDA pregnancy category B. Lidoderm is not expected to be harmful to an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant during treatment. Lidocaine can pass into breast milk and may harm a nursing baby. Do not use this medication without telling your doctor if you are breast-feeding a baby.

See also: [Pregnancy and breastfeeding warnings](#) (in more detail)

Reduce Nerve Pain Now



neuropathytreatmentgroup.com

Don't Let Neuropathy Symptoms Slow You Down. Get A Free Trial Now!

2015 Best Skin Tighteners

Best Arthritis Cream



Image Credit: U.S. FDA/Drugs.com

Email me:

News and Warnings related to this drug

FDA Medwatch Alerts for all Medications

News Roundup Daily

Related News and Articles

[AcelRx Pharmaceuticals Provides Zalviso Regulatory Update](#) May 4, 2015

AcelRx Pharmaceuticals, Inc. (Nasdaq: ACRX), a specialty pharmaceutical company focused on the development and commercialization of innovative therapies for...

[Too Many Americans Neglect Backs in Skin Cancer Prevention](#) May 4, 2015

A new survey finds that many people in the United States are forgetting their backs when they try to be forward-thinking about skin cancer prevention. Experts...

[Health Tip: Promptly Treat a Minor Burn](#) April 28, 2015

Burns less than 3 inches in diameter usually can be cared for at home. The U.A. National Safety Council suggests: Hold the burned area under cool running...



Harvard Health Publications

Related: [Pain](#)

How should I use Lidoderm?

Use Lidoderm exactly as directed on the label, or as it has been prescribed by your doctor. Do not use the medication in larger amounts, or use it for longer than recommended.

Your body may absorb more lidocaine if you use too much, if you apply it over large skin areas, or if you apply heat, bandages, or plastic wrap to treated skin areas. Skin that is cut or irritated may also absorb more topical medication than healthy skin.

Use the smallest amount of this medication needed to numb the skin or relieve pain. Do not use large amounts of Lidoderm, or cover treated skin areas with a bandage or plastic wrap without medical advice. Be aware that many cosmetic procedures are performed without a medical doctor present.

Do not apply Lidoderm to swollen skin areas or deep puncture wounds. Avoid using the medicine on skin that is raw or blistered, such as a severe burn or abrasion.

Lidoderm may be applied with your finger tips or a cotton swab. Follow your doctor's instructions.

Store Lidoderm at room temperature away from moisture and heat.

What happens if I miss a dose?

Since Lidoderm is used as needed, you may not be on a dosing schedule. If you are using the medication regularly, use the missed dose as soon as you remember. If it is almost time for your next dose, skip the missed dose and use the medicine at your next regularly scheduled time. Do not use extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention if you think you have used too much of this medicine. Lidoderm applied to the skin is not likely to cause an overdose unless you apply more than the recommended dose. Overdose may also occur if you apply heat, bandages, or plastic wrap to treated skin areas.

Improper use of Lidoderm may result in death.

Overdose symptoms may include drowsiness, confusion, nervousness, ringing in your ears, blurred vision, feeling hot or cold, numbness, muscle twitches, uneven heartbeats, seizure (convulsions), slowed breathing, or respiratory failure (breathing stops).

What should I avoid?

Do not allow Lidoderm to come into contact with your eyes. If it does, rinse with water.

Avoid using other topical medications on the affected area unless your doctor has told you to.

Zovirax Coupon

Get A Free Printable
Discount Now. Visit Us
Online & Save Up To 75%.



Lidoderm side effects

Get emergency medical help if you have any of these signs of an allergic reaction to Lidoderm: hives; difficulty breathing; swelling of your face, lips, tongue, or throat. Stop using Lidoderm and call your doctor at once if you have any of these serious side effects:

- uneven heartbeats;
- drowsiness, confusion;
- tremors, seizure (convulsions); or
- blurred vision.

Less serious Lidoderm side effects include:

- mild irritation, redness, or swelling where the medication is applied;
- numbness in places where the medicine is accidentally applied.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.



See also: [Side effects](#) (in more detail)

What other drugs will affect Lidoderm?

Before using Lidoderm, tell your doctor if you are taking medication to treat a heart rhythm disorder, such as:

- quinidine (Quinidex, Quinaglute);
- disopyramide (Norpace);
- flecainide (Tambocor);
- mexiletine (Mexitil);
- procainamide (Procan, Pronestyl);
- tocainide (Tonocard); or
- propafenone (Rythmol).

This list is not complete and there may be other drugs that can interact with Lidoderm. Tell your doctor about all the prescription and over-the-counter medications you use. This includes vitamins, minerals, herbal products, and drugs prescribed by other doctors. Do not start using a new medication without telling your doctor.

 Google  Facebook  Twitter  Print  Email  Add to Mednotes

Next Page → [Side Effects](#)

More about Lidoderm (lidocaine topical)

- [Side Effects](#)
- [During Pregnancy](#) or [Breastfeeding](#)
- [Dosage Information](#)
- [Drug Interactions](#)
- [Support Group](#)
- [Pricing & Coupons](#)
- [En Espanol](#)
- [22 Reviews](#) - [Add your own review/rating](#)
- [Generic Availability](#)

Consumer resources

- [Lidoderm patch](#)
- [Lidoderm Topical application](#) (Advanced Reading)
- *Other brands: [Xylocaine Jelly](#), [LidoRx](#), [Bactine](#), [Zingo](#), [More \(10\)](#) »*

Professional resources

- [Lidocaine Topical Oral Solution 4%](#) (FDA)

Related treatment guides

- [Pain](#)
- [Anal Itching](#)
- [Anesthesia](#)
- [Burns, External](#)
- [Hemorrhoids](#)
- [More \(3\)](#) »

Where can I get more information?

- Your pharmacist has information about Lidoderm.

Find an Opioid Treatment

Get help and treatment options. Find a doctor near you.



Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Disclaimer: Every effort has been made to ensure that the information provided by Cerner Multum, Inc. ('Multum') is accurate, up-to-date, and complete, but no guarantee is made to that effect. Drug information contained herein may be time sensitive. Multum information has been compiled for use by healthcare practitioners and consumers in the United States and therefore Multum does not warrant that uses outside of the United States are appropriate, unless specifically indicated otherwise. Multum's drug information does not endorse drugs, diagnose patients or recommend therapy. Multum's drug information is an informational resource designed to assist licensed healthcare practitioners in caring for their patients and/or to serve consumers viewing this service as a supplement to, and not a substitute for, the expertise, skill, knowledge and judgment of healthcare practitioners. The absence of a warning for a given drug or drug combination in no way should be construed to indicate that the drug or drug combination is safe, effective or appropriate for any given patient. Multum does not assume any responsibility for any aspect of healthcare administered with the aid of information Multum provides. The information contained herein is not intended to cover all possible uses, directions, precautions, warnings, drug interactions, allergic reactions, or adverse effects. If you have questions about the drugs you are taking, check with your doctor, nurse or pharmacist.

Copyright 1996-2015 Cerner Multum, Inc. Version: 3.02. Revision Date: 4/12/2009 4:37:53 PM.



Support

[Advertise With Us](#)
[Frequently Asked Questions](#)
[Ask a Question](#)
[Content Submissions](#)
[Mobile Apps](#)
[About Drugs.com](#)
[Contact Us](#)
[Sitemap](#)

Follow:     

About Drugs.com

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment.

Data sources include Micromedex® (updated Apr 27th, 2015), Cerner Multum™ (updated Apr 20th, 2015), Wolters Kluwer™ (updated Apr 6th, 2015) and others. To view content sources and attributions, refer to our [editorial policy](#).

Copyright © 2000-2015 Drugs.com. All rights reserved.

Your Privacy



We comply with the HONcode standard for trustworthy health information.

[Verify here](#)



Ad Choices

[Terms of Use](#) · [Privacy Policy](#)