

SUBOXONE® Sublingual Film

(buprenorphine and naloxone) CIII. Find a doctor.



ADVERTISEMENT

Pill Identifier App
Now available for the iPhone and iPad.
[View Details](#)

Browse all medications [A](#) [B](#) [C](#) [D](#) [E](#) [F](#) [G](#) [H](#) [I](#) [J](#) [K](#) [L](#) [M](#) [N](#) [O](#) [P](#) [Q](#) [R](#) [S](#) [T](#) [U](#) [V](#) [W](#) [X](#) [Y](#) [Z](#) [Advanced Search](#) [Phonetic Search](#)[Drugs A-Z](#) ▾[Pill Identifier](#)[Interactions Checker](#)[News](#) ▾[Health Professionals](#) ▾[Q & A](#) ▾[Mednotes](#)[Apps](#)[Home](#) → [Conditions](#) → [ADHD](#) → [Vyvanse](#)[Print](#) [Share](#)[Sign In](#) or [Register](#)

Vyvanse

Generic Name: [lisdexamfetamine](#) (lis dex am FET a meen)Brand Names: [Vyvanse](#)

Overview

[Side Effects](#)[Dosage](#)[Interactions](#)[For Professionals](#)[More ▾](#)

Stem Cell Treatment

Attention Lung Diseases Sufferers Find Out If You're a Candidate



What is Vyvanse?

Vyvanse (lisdexamfetamine) is a central nervous system stimulant. It affects chemicals in the brain and nerves that contribute to hyperactivity and impulse control.

Vyvanse is used to treat attention deficit hyperactivity disorder (ADHD) in adults and in children who are at least 6 years old.

Vyvanse is also used to treat moderate to severe binge eating disorder in adults. This medicine is not to be used for obesity or weight loss.

Important information

Do not use Vyvanse if you have taken an MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, methylene blue injection, phenelzine, rasagiline, selegiline, tranylcypromine, and others.

Vyvanse is a drug of abuse and may be habit-forming. Keep track of the amount of medicine used from each new bottle. You should be aware if anyone is using your medicine improperly or without a prescription. **Using this medication improperly can cause death or serious side effects on the heart.**

ADVERTISEMENT

ADD/ADHD Symptom Relief

SAFE NATURAL HOME REMEDY FOR EAS BALANCE

SYNAPTOL®

LEARN MORE! [hellolife.net](#)

Slideshow: Does Your Child Have ADHD? Recognizing Signs & Treatment Options



Before taking this medicine

You should not use Vyvanse if you are allergic

Related Information



Availability
Prescription only



Pregnancy Category
Risk cannot be ruled out



CSA Schedule
High potential for abuse



Approval History
Drug history at FDA

Reviews

383 User Reviews

Average User Rating

8.2 Rate it!

Manufacturer

[Shire US Inc](#)

Drug Class

[CNS stimulants](#)

Related Drugs

[Binge Eating Disorder](#)
[lisdexamfetamine](#)
More...

ADHD

[Adderall](#)
[methylphenidate](#)
[Concerta](#)
[Ritalin](#)
[Strattera](#)
[Adderall XR](#)
[Focalin](#)
[amphetamine/dextroamphetamine](#)

[Focalin XR](#)
[dexmethylphenidate](#)
[atomoxetine](#)
[lisdexamfetamine](#)
More...

ADVERTISEMENT



Groundbreaking treatments for peripheral arterial disease.

[Click to learn more >](#)

Prakash Makam, MD

CARDIOLOGY ASSOCIATES

Vyvanse (lisdexamfetamine) Images



04/07/2015



to lisdexamfetamine.

Do not use Vyvanse if you have taken an MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, methylene blue injection, phenelzine, rasagiline, selegiline, tranylcypromine, and others.

Some stimulants have caused sudden death in people with serious heart problems or congenital heart defects. Tell your doctor if you have:

- heart disease, heart rhythm disorder;
- high blood pressure;
- coronary artery disease (hardened arteries); or
- history of heart attack.

To make sure this medicine is safe for you, tell your doctor if you have:

- kidney disease;
- circulation problems;
- peripheral vascular disease such as Raynaud's syndrome;
- a personal or family history of mental illness, psychotic disorder, bipolar illness, depression, or suicide attempt; or
- if you take an antidepressant.

FDA pregnancy category C. It is not known whether this medication will harm an unborn baby. However, lisdexamfetamine may cause premature birth, low birth weight, or withdrawal symptoms in a newborn if the mother takes Vyvanse during pregnancy. Tell your doctor if you are pregnant or plan to become pregnant while using this medication.

See also: [Pregnancy and breastfeeding warnings](#) (in more detail)

Lisdexamfetamine can pass into breast milk and may harm a nursing baby. Tell your doctor if you are breast-feeding a baby.

Vyvanse is not approved to treat **ADHD** in a child younger than 6 years old. This medicine is not approved to treat **binge eating disorder** in anyone younger than 18 years old.

Long-term use of Vyvanse can slow a child's growth. Tell your doctor if the child using this medication is not growing or gaining weight properly.

Do not give Vyvanse to a child younger than 6 years old without the advice of a doctor.

[Find a Doctor Near You](#)



[opioiddependence.com](#)

Dependent on opioid painkillers? Get treated in a doctor's office.

[Homeopathic ADHD Remedy](#)



[Printable Coupons Here](#)



How should I take Vyvanse?



Vyvanse (lisdexamfetamine) 50 mg



Vyvanse (lisdexamfetamine) 30 mg

[View all 13 images](#)

Get Updates by Email

Email me:

News and Warnings related to this drug

FDA Medwatch Alerts for all Medications

News Roundup Daily

Related News and Articles

[Prenatal Exposure to Pollutants May Spell Trouble for Kids](#) March 25, 2015

Exposure to common air pollutants before birth may make children more likely to have the symptoms of attention-deficit/hyperactivity disorder (ADHD) and other...

[ADHD Children May Gain More From Family-Centered Care](#) March 23, 2015

For children with attention-deficit/hyperactivity disorder (ADHD), receiving more family-centered, compassionate care may be more effective than standard care,...

[Can Fish Oil Help Boys With ADHD Pay Attention?](#)

March 19, 2015

Boys with attention-deficit/hyperactivity disorder may benefit from the omega-3 fatty acids found in fish and some vegetable oils, a small European study...

Multiple Sclerosis

Get the basics for managing MS.

Watch video ▶▶▶



Take Vyvanse exactly as prescribed by your doctor. Follow all directions on your prescription label. Your doctor may occasionally change your dose to make sure you get the best results. Do not take this medicine in larger or smaller amounts or for longer than recommended.

Lisdexamfetamine is a drug of abuse and may be habit-forming. Keep track of the amount of medicine used from each new bottle. You should be aware if anyone is using your medicine improperly or without a prescription. Using this medication improperly can cause death or serious side effects on the heart.

Vyvanse should be taken in the morning with or without food.

Do not take this medicine in the evening because it may cause sleep problems (insomnia).

Do not crush, chew, break, or divide a capsule. Swallow the capsule whole. To make swallowing easier, you may open the capsule and sprinkle the medicine into a glass of water or mix it with yogurt. After the medicine has dissolved, drink or eat the mixture immediately. Do not save for later use.

Your doctor will need to check your progress on a regular basis. Your blood pressure and heart rate will need to be checked often.

Store at room temperature away from moisture, heat, and light.

Throw away unused or expired Vyvanse in a sealed container or bag. Ask your pharmacist where to locate a community pharmaceutical take back disposal program.

What happens if I miss a dose?

Take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose, or if it is already evening. Taking this medicine late in the day can cause sleep problems. Do not take extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. **An overdose of lisdexamfetamine can be fatal.**

Overdose symptoms may include restlessness, tremor, muscle twitches, rapid breathing, confusion, hallucinations, panic, aggressiveness, muscle pain or tenderness, muscle weakness, fever or flu symptoms, and dark colored urine. These symptoms may be followed by depression and tiredness. Other overdose symptoms include nausea, vomiting, diarrhea, stomach pain, uneven heartbeats, feeling light-headed, fainting, seizure (convulsions), or coma.

What should I avoid?

Vyvanse may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be alert.

Drinking fruit juices or taking vitamin C at the same time you take Vyvanse can make your body absorb less of the medicine.

[Apply for Disability](#)



 disability-benefits-help.org

Are you eligible for up to \$2642/mo in Social Security Disability?

[I Stopped My Hair Loss](#) ▾

[My 1 Trick for Hair Loss](#) ▾

Vyvanse side effects

Get emergency medical help if you have any of these **signs of an allergic reaction to Vyvanse**: hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Stop using Vyvanse and call your doctor at once if you have:

- chest pain, shortness of breath, fast or uneven heart rate;
- confusion, paranoia, unusual thoughts or behavior, hallucinations;
- penis erection that is painful or lasts 4 hours or longer (rare); or
- numbness, pain, cold feeling, unexplained wounds, or skin color changes (pale, red, or blue appearance) in your fingers or toes.

Common Vyvanse side effects may include:

- dry mouth, loss of appetite, weight loss;
- sleep problems (insomnia);
- dizziness, feeling irritable; or
- nausea, vomiting, stomach pain, diarrhea.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

See also: [Side effects](#) (in more detail)

Vyvanse dosing information

Usual Adult Dose for Attention Deficit Disorder:

Initial dose: 30 mg orally once a day in the morning

If necessary, the initial daily dose may be increased by 10 mg to 20 mg approximately once a week up to a maximum daily dose of 70 mg. Maintain patients on their optimal dose.

Usual Pediatric Dose for Attention Deficit Disorder:

6 years or older:

Initial dose: 30 mg orally once a day in the morning







If necessary, the initial daily dose may be increased by 10 mg to 20 mg approximately once a week up to a maximum daily dose of 70 mg. Maintain patients on their optimal dose.

Vyvanse has not been studied in children under 6 years of age. Long-term effects of amphetamines in children have not been well established. Amphetamines are not recommended for use in children under 3 years of age.

What other drugs will affect Vyvanse?

Ask your doctor before using a stomach acid medicine. Some of these medicines can change the way your body absorbs Vyvanse, and may increase side effects.

Other drugs may interact with lisdexamfetamine, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell each of your health care providers about all medicines you use now and any medicine you start or stop using.

 [Google](#)  [Facebook](#)  [Twitter](#)  [Print](#)  [Email](#)  [Add to Mednotes](#)

Next Page → [Side Effects](#)

More about Vyvanse (lisdexamfetamine)

- [Side Effects](#)
- [During Pregnancy](#) or [Breastfeeding](#)
- [Dosage Information](#)
- [Drug Images](#)

- [Drug Interactions](#)
- [Support Group](#)
- [Pricing & Coupons](#)
- [En Espanol](#)
- [383 Reviews](#) - [Add your own review/rating](#)

Consumer resources

- [Vyvanse](#)
- [Vyvanse](#) (Advanced Reading)

Professional resources

- [Vyvanse](#) (AHFS Monograph)
- [Vyvanse](#) (FDA)

Related treatment guides

- [Binge Eating Disorder](#)
- [ADHD](#)

External resources

- [Official Vyvanse Website \(US\)](#)
- [Manufacturer - Shire](#)
- [Attention Deficit Disorder Association](#)

Where can I get more information?

- Your pharmacist can provide more information about Vyvanse.

Never Eat These "5 Foods"

Here are 5 foods you should never eat if You want to lose belly fat



Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Vyvanse only for the indication prescribed.

Disclaimer: Every effort has been made to ensure that the information provided by Cerner Multum, Inc. ('Multum') is accurate, up-to-date, and complete, but no guarantee is made to that effect. Drug information contained herein may be time sensitive. Multum information has been compiled for use by healthcare practitioners and consumers in the United States and therefore Multum does not warrant that uses outside of the United States are appropriate, unless specifically indicated otherwise. Multum's drug information does not endorse drugs, diagnose patients or recommend therapy. Multum's drug information is an informational resource designed to assist licensed healthcare practitioners in caring for their patients and/or to serve consumers viewing this service as a supplement to, and not a substitute for, the expertise, skill, knowledge and judgment of healthcare practitioners. The absence of a warning for a given drug or drug combination in no way should be construed to indicate that the drug or drug combination is safe, effective or appropriate for any given patient. Multum does not assume any responsibility for any aspect of healthcare administered with the aid of information Multum provides. The information contained herein is not intended to cover all possible uses, directions, precautions, warnings, drug interactions, allergic reactions, or adverse effects. If you have questions about the drugs you are taking, check with your doctor, nurse or pharmacist.

Copyright 1996-2015 Cerner Multum, Inc. Version: 7.02. Revision Date: 2015-02-10, 3:07:42 PM.



Support

- [Advertise With Us](#)
- [Frequently Asked Questions](#)
- [Ask a Question](#)
- [Content Submissions](#)
- [Mobile Apps](#)
- [About Drugs.com](#)

About Drugs.com

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment.

Data sources include Micromedex® (updated Feb 4th, 2015), Cerner Multum™ (updated Feb 17th, 2015), Wolters Kluwer™ (updated Apr 6th,

Your Privacy

We comply with the HONcode standard for trustworthy health information.
[Verify here](#)



[Terms of Use](#) · [Privacy Policy](#)

[Contact Us](#)

[Sitemap](#)

Follow:     

2015) and others. To view content sources and attributions, refer to our [editorial policy](#).

Copyright © 2000-2015 Drugs.com. All rights reserved.