

ADVERTISEMENT

**GET THE FACTS ABOUT EPILEPSY**

[Watch the video >](#)



Browse all medications [A](#) [B](#) [C](#) [D](#) [E](#) [F](#) [G](#) [H](#) [I](#) [J](#) [K](#) [L](#) [M](#) [N](#) [O](#) [P](#) [Q](#) [R](#) [S](#) [T](#) [U](#) [V](#) [W](#) [X](#) [Y](#) [Z](#) [Advanced Search](#) [Phonetic Search](#)

- Home
- Drugs A-Z
- Pill Identifier
- Interactions Checker
- News
- Health Professionals
- Q & A
- Mednotes
- Apps

Home → Conditions → Seizure Prevention → Lamictal

[Print](#) [Share](#)

[Sign In](#) or [Register](#)

# Lamictal

Generic Name: [lamotrigine](#) (la MOE tri jeen)  
 Brand Names: *LaMICTal*, *LaMICTal ODT*, *LaMICTal XR*

- Overview
- Side Effects
- Dosage
- Interactions
- For Professionals
- More

## Loved One with Bipolar?

Proven methods to cope & deal with your loved one's bipolar disorder



## What is Lamictal?

Lamictal (lamotrigine) is an anti-epileptic medication, also called an anticonvulsant.

Lamictal is used either alone or in combination with other medications to treat epileptic seizures in adults and children. Lamotrigine is also used to delay mood episodes in adults with bipolar disorder (manic depression).

The immediate-release form of Lamictal (regular tablet and orally disintegrating tablet) can be used in children as young as 2 years old when it is given as part of a combination of seizure medications. However, this form should not be used as a single medication in a child or teenager who is younger than 16 years old.

The extended-release form of lamotrigine (Lamictal XR) is for use only in adults and children who are at least 13 years old.

ADVERTISEMENT

**Stop Depression & Anxiety**

*Or Return Your Device for a Refund*

[Learn More](#)

## Related Information

- Rx** **Availability**  
Prescription only
- C** **Pregnancy Category**  
Risk cannot be ruled out
- N/A** **CSA Schedule**  
Not a controlled drug
- Approval History**  
Drug history at FDA

Learn about treating **BIPOLAR DISORDER**

[LEARN MORE](#)

**Harvard Health Publications**

Related: [Bipolar Disorder \(Manic Depressive Illness or Manic Depression\)](#)

Reviews  
**273 User Reviews**

Average User Rating  
**8.4** Rate it!

Manufacturer  
[GlaxoSmithKline](#)

Drug Class  
[Triazine anticonvulsants](#)

Related Drugs  
[Bipolar Disorder](#)  
[Seroquel](#)

[Lamotrigine](#)  
[Abilify](#)  
[Klonopin](#)  
[Depakote](#)  
[quetiapine](#)  
[risperidone](#)  
 More...  
[Schizoaffective Disorder](#)  
[Seroquel](#)  
[Prozac](#)  
[lithium](#)  
[Abilify](#)  
[fluoxetine](#)  
[quetiapine](#)

ADVERTISEMENT

**The Depression Treatment Device**

Fisher Wallace Stimulator  
 212 688 8100  
[fisherwallace.com](#)

Learn about **causes, treatments and first aid** for epileptic seizures. [Click Here](#)

## Important information

Lamictal may cause a severe or life-threatening skin rash, especially in children and in people who take too high of a starting a dose, or those who also take valproic acid (Depakene) or divalproex (Depakote).

Seek emergency medical attention if you have: fever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain, followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling.


You may have thoughts about suicide while taking this medication. Your doctor will need to check you at regular visits while you are using

**Slideshow: Breaking Down Bipolar Disorder - 12 Things You Need To Know**

04/07/2015


Lamictal. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, depression, anxiety, or if you feel agitated, hostile, restless, hyperactive (mentally or physically), or have thoughts about suicide or hurting yourself.

Do not stop taking Lamictal without first talking to your doctor, even if you feel fine. You may have increased seizures if you stop using Lamictal suddenly. Contact your doctor if your seizures get worse or you have them more often while taking this medicine. Taking this medication during early pregnancy can increase the risk of the baby being born with cleft lip or cleft palate. Tell your doctor if you are pregnant or if you become pregnant during treatment.



risperidone  
lamotrigine  
More...

Seizure Prevention  
clonazepam  
diazepam  
Klonopin  
Valium  
Topamax  
lamotrigine  
topiramate  
magnesium sulfate  
More...



## Before taking this medicine

You should not take Lamictal if you are allergic to lamotrigine.

Lamictal may cause a severe or life-threatening skin rash, especially in children and in people who take too high a dose at the start of treatment. Serious skin rash may also be more likely to occur if you are taking lamotrigine together with valproic acid (Depakene) or divalproex (Depakote).

To make sure Lamictal is safe for you, tell your doctor if you have:

- kidney or liver disease;
- a history of depression or suicidal thoughts or actions; or
- if you are allergic to other seizure medications.

Some people have thoughts about suicide while taking this medicine. Your doctor will need to check your progress at regular visits while you are using Lamictal. Your family or other caregivers should also be alert to changes in your mood or symptoms.

FDA pregnancy category C. It is not known whether Lamictal will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant while using this medicine.

**See also:** [Pregnancy and breastfeeding warnings](#) (in more detail)

If you are pregnant, your name may be listed on a pregnancy registry. This is to track the outcome of the pregnancy and to evaluate any effects of Lamictal on the baby.

Birth control pills can make Lamictal less effective, resulting in increased seizures. **Tell your doctor if you start or stop using birth control pills while you are taking this medicine.** Your lamotrigine dose may need to be changed.

Lamotrigine can pass into breast milk and may harm a nursing baby. Tell your doctor if you are breast-feeding a baby.

**SUBOXONE®**  
Sublingual Film



[suboxone.com](#)

(buprenorphine and naloxone) CIII. Find a doctor.


[Apply for Disability](#) ▾

[Reduce Nerve Pain Now](#) ▾

## How should I take Lamictal?

Take Lamictal exactly as prescribed by your doctor. Follow all directions on your prescription label.

## Lamictal (lamotrigine) Images



[View all 17 images](#)

## Get Updates by Email

### Email me:

- News and Warnings related to this drug
- FDA Medwatch Alerts for all Medications
- News Roundup
- Daily

## Related News and Articles

### Actavis Receives FDA Approval of Saphris for Pediatric Patients with Bipolar I Disorder

March 13, 2015

Actavis plc (NYSE: ACT) today announced that the U.S. Food and Drug Administration (FDA) has approved its supplemental new drug application (sNDA) for Saphris...

### Study Highlights Complexity of 'Hearing Voices'

March 11, 2015

Rachel Waddingham hears voices. "I hear about 13 or so voices," she said in a news release from Durham University, in England. "Each of them is different --..."

### Epilepsy Surgery Gets High Marks From Patients in Survey

February 26, 2015

More than nine in 10 epilepsy patients who had brain surgery to try to control their seizures are happy they did so, a new survey reveals. The review appears...

**GET THE FACTS ABOUT  
EPILEPSY**

[Watch the video](#) >

Your doctor may occasionally change your dose to make sure you get the best results. Do not take this medicine in larger or smaller amounts or for longer than recommended.

**Taking too much Lamictal at the start of treatment may increase your risk of a severe life-threatening skin rash.**

You may need frequent blood tests to help your doctor make sure you are taking the right dose.

**Extended-release** Lamictal XR may be used for different conditions than **immediate-release** Lamictal is used for. Always check your refills to make sure you have received the correct size, color, and shape of tablet. Ask the pharmacist if you have any questions about the medicine you receive at the pharmacy.

If you switch to Lamictal from another seizure medicine, carefully follow your doctor's instructions about the timing and dosage of your medications. Read all patient information, medication guides, and instruction sheets provided to you. Ask your doctor or pharmacist if you have any questions.

Do not crush, chew, or break a **regular Lamictal tablet**. Swallow it whole.

Do not swallow the orally disintegrating tablet (ODT) whole. Place it on your tongue and move it around in your mouth. Allow the tablet to dissolve in your mouth without chewing. Swallow several times as the tablet dissolves. If desired, you may drink liquid to help swallow the dissolved tablet.

To take the chewable dispersible tablet, you may either swallow it whole with a glass of water, or chew it first and then swallow it. You may also place the tablet into 1 teaspoon of water or diluted fruit juice and allow it to disperse in the liquid for about 1 minute. Gently swirl the liquid and then swallow all of the mixture right away. Do not save it for later use.

Do not stop using Lamictal suddenly, even if you feel fine. Stopping suddenly may cause increased seizures. Follow your doctor's instructions about tapering your dose.

Wear a medical alert tag or carry an ID card stating that you take Lamictal. Any medical care provider who treats you should know that you take seizure medication.

Use this medicine regularly to get the most benefit. Get your prescription refilled before you run out of medicine completely.

Store at room temperature away from light and moisture.

## What happens if I miss a dose?

Take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not take extra medicine to make up the missed dose.

## What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include blurred vision, problems with coordination, increased seizures, feeling light-headed, or fainting.

## What should I avoid while taking Lamictal?

Lamictal may cause blurred vision or impair your thinking or reactions. Be careful if you drive or do anything that requires you to be alert and able to see clearly.

[Apply For Disability](#)



† [disabilityapprovalguide.com](http://disabilityapprovalguide.com)

Find Out If You May Qualify For Up To  
\$2,663/mo in Disability Benefits

[2015 Best Skin  
Tighteners](#)



## 5) Foods To Never Eat

### Lamictal side effects

Get emergency medical help if you have any **signs of an allergic reaction to Lamictal**: hives; fever; swollen glands; painful sores in or around your eyes or mouth; difficulty breathing; swelling of your face, lips, tongue, or throat.

Seek emergency medical attention if you have **signs of a life-threatening skin rash**: fever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain, followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling.

If you have to stop taking Lamictal because of a serious skin rash, you may not be able to take it again in the future.

Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, depression, anxiety, or if you feel agitated, hostile, restless, hyperactive (mentally or physically), or have thoughts about suicide or hurting yourself.

Call your doctor at once if you have:

- the first sign of any skin rash, no matter how mild;
- changes in your menstrual periods;
- low blood cell counts - flu-like symptoms, swollen gums, mouth sores, skin sores, rapid heart rate, pale skin, easy bruising, unusual bleeding, feeling light-headed;
- possible signs of brain inflammation - fever, chills, headache, neck stiffness, increased sensitivity to light, purple spots on the skin, confusion, drowsiness; or
- signs of inflammation in your body - swollen glands, severe tingling or numbness, muscle weakness, upper stomach pain, jaundice (yellowing of the skin or eyes), chest pain, new or worsening cough with fever, trouble breathing.

Common Lamictal side effects may include:

- headache, dizziness;
- blurred vision, double vision;
- tremor, loss of coordination;
- dry mouth, nausea, vomiting, stomach pain;
- fever, sore throat, runny nose;
- drowsiness, tired feeling;
- back pain; or
- sleep problems (insomnia).

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**See also:** [Side effects](#) (in more detail)

### What other drugs will affect Lamictal?






Other drugs may interact with lamotrigine, including prescription and over-the-counter medicines, vitamins, and herbal products.

Tell your doctor about all other medicines you use, especially:

- hormonal forms of birth control (pills, injections, implants, or skin patches);
- carbamazepine (Carbatrol, Equetro, Tegretol);
- divalproex (Depakote);
- oxcarbazepine (Trileptal);
- phenobarbital (Solfoton);
- primidone (Mysoline);
- phenytoin (Dilantin);

- rifampin (Rifadin, Rifater, Rifamate, Rimactane); or
- valproic acid (Depakene, Stavzor).

This list is not complete and other drugs may interact with Lamictal. Tell your doctor about all medications you use. Do not start a new medication without telling your doctor.

 Google  Facebook  Twitter  Print  Email  Add to Mednotes

Next Page → [Side Effects](#)

## More about Lamictal (lamotrigine)

- [Side Effects](#)
- [During Pregnancy](#) or [Breastfeeding](#)
- [Dosage Information](#)
- [Drug Images](#)
- [Drug Interactions](#)
- [Support Group](#)
- [Pricing & Coupons](#)
- [273 Reviews](#) - [Add your own review/rating](#)
- [Generic Availability](#)

### Consumer resources

- [Lamictal](#)
- [Lamictal Orally Disintegrating Tablets](#)
- [Lamictal chewable dispersible tablets](#)
- [Lamictal](#) (Advanced Reading)

### Professional resources

- [Lamictal](#) (AHFS Monograph)
- [Lamictal](#) (FDA)

### Other formulations

- [Lamictal XR](#)
- [Lamictal ODT](#)
- [Lamictal CD](#)

### Related treatment guides

- [Bipolar Disorder](#)
- [Schizoaffective Disorder](#)
- [Seizure Prevention](#)

## Where can I get more information?

- Your pharmacist can provide more information about Lamictal.

## Prostate Cancer Stages

See the Progression of Stages for Prostate Cancer. Get Expert Info.



Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Disclaimer: Every effort has been made to ensure that the information provided by Cerner Multum, Inc. ('Multum') is accurate, up-to-date, and complete, but no guarantee is made to that effect. Drug information contained herein may be time sensitive. Multum information has been compiled for use by healthcare practitioners and consumers in the United States and therefore Multum does not warrant that uses outside of the United States are appropriate, unless specifically indicated otherwise. Multum's drug information does not endorse drugs, diagnose patients or recommend therapy. Multum's drug information is an informational resource designed to assist licensed healthcare practitioners in caring for their patients and/or to serve

consumers viewing this service as a supplement to, and not a substitute for, the expertise, skill, knowledge and judgment of healthcare practitioners. The absence of a warning for a given drug or drug combination in no way should be construed to indicate that the drug or drug combination is safe, effective or appropriate for any given patient. Multum does not assume any responsibility for any aspect of healthcare administered with the aid of information Multum provides. The information contained herein is not intended to cover all possible uses, directions, precautions, warnings, drug interactions, allergic reactions, or adverse effects. If you have questions about the drugs you are taking, check with your doctor, nurse or pharmacist.

Copyright 1996-2015 Cerner Multum, Inc. Version: 17.01. Revision Date: 2015-01-22, 9:49:36 AM.



## Support

[Advertise With Us](#)  
[Frequently Asked Questions](#)  
[Ask a Question](#)  
[Content Submissions](#)  
[Mobile Apps](#)  
[About Drugs.com](#)  
[Contact Us](#)  
[Sitemap](#)

Follow:     

## About Drugs.com

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment.

Data sources include Micromedex® (updated Feb 4th, 2015), Cerner Multum™ (updated Feb 17th, 2015), Wolters Kluwer™ (updated Apr 6th, 2015) and others. To view content sources and attributions, refer to our [editorial policy](#).

Copyright © 2000-2015 Drugs.com. All rights reserved.

## Your Privacy



We comply with the HONcode standard for trustworthy health information.

[Verify here](#)



Ad Choices

[Terms of Use](#) · [Privacy Policy](#)