

National Institutes of Health / U.S. National Library of Medicine



[Home](#) → [Medical Encyclopedia](#) → Diskectomy

URL of this page: //medlineplus.gov/ency/article/007250.htm

Diskectomy

Diskectomy is surgery to remove all or part of the cushion that helps support part of your spinal column. These cushions are called disks, and they separate your spinal bones (vertebrae).

Description

A surgeon may perform disk removal (diskectomy) in these different ways.

- **Microdiskectomy:** When you have a microdiskectomy, the surgeon does not need to do much surgery on the bones, joints, ligaments, or muscles of your spine.
- Diskectomy in the lower part of your back (lumbar spine) may be part of a larger surgery that also includes a laminectomy, foraminotomy, or spinal fusion.
- Diskectomy in your neck (cervical spine) is most often done along with laminectomy, foraminotomy, or fusion.

Microdiskectomy is done in a hospital or outpatient surgical center. You will be given spinal anesthesia (to numb your spine area) or general anesthesia (asleep and pain-free).

- The surgeon makes a small (1 to 1-1/2 inch) incision (cut) on your back and moves the back muscles away from your spine. The surgeon uses a special microscope to see the problem disk or disks and nerves during surgery.
- The nerve root is located and gently moved away.
- The surgeon removes the injured disk tissue and pieces of the disk.
- The back muscles are returned to place.
- The incision is closed with stitches or staples.
- The surgery takes about 1 to 2 hours.

Diskectomy and laminotomy are usually done in the hospital, using general anesthesia (asleep

and pain-free).

- The surgeon makes a larger cut on your back over the spine.
- Muscles and tissue are gently moved to expose your spine.
- A small part of the lamina bone (part of the vertebrae that surrounds the spinal column and nerves) is cut away. The opening may be as large as the ligament that runs along your spine.
- A small hole is cut in the disk that is causing your symptoms. Material from inside the disk is removed. Other fragments of the disk may also be removed.

Why the Procedure is Performed

When one of your disks moves out of place (herniates), the soft gel inside pushes through the wall of the disk. The disk may then place pressure on the spinal cord and nerves that are coming out of your spinal column.

Many of the symptoms caused by a herniated disk get better or go away over time without surgery. Most people with low back or neck pain, numbness, or even mild weakness are often first treated with anti-inflammatory medicines, physical therapy, and exercise.

Only a few people with a herniated disk need surgery.

Your doctor may recommend a discectomy if you have a herniated disk and:

- Leg pain or numbness that is very bad or is not going away, making it hard to do daily tasks
- Severe weakness in muscles of your lower leg or buttocks
- Pain that spreads into your buttocks or legs

If you are having problems with your bowels or bladder, or the pain is so bad that strong pain drugs do not help, you will probably have surgery right away.

Risks

Risks for anesthesia and surgery in general are:

- Reactions to medicines
- Breathing problems
- Bleeding, blood clots, infection

Risks for this surgery are:

- Damage to the nerves that come out of the spine, causing weakness or pain that does not go away
- Your back pain does not get better, or pain comes back later
- Pain after surgery, if all the disk fragments are not removed
- Spinal fluid may leak
- The disk may bulge out again

Before the Procedure

Tell your doctor or nurse what medicines you are taking, even medicines, supplements or herbs you bought without a prescription.

During the days before the surgery:

- Prepare your home for when you come back from the hospital.
- If you are a smoker, you need to stop. Your recovery will be slower and possibly not as good if you continue to smoke. Ask your doctor for help.
- 2 weeks before surgery, you may be asked to stop taking medicines that make it harder for your blood to clot. These include aspirin, ibuprofen (Advil, Motrin), naproxen (Aleve, Naprosyn), and other medicines like these.
- If you have diabetes, heart disease, or other medical problems, your surgeon will ask you to see the doctors who treat you for those conditions.
- Talk with your doctor if you have been drinking a lot of alcohol.
- Ask your doctor which medicines you should still take on the day of the surgery.
- Always let your doctor know about any cold, flu, fever, herpes breakout, or other illnesses you may have.
- You may want to visit the physical therapist to learn some exercises to do before surgery and to practice using crutches.

On the day of the surgery:

- Follow instructions about when to stop eating and drinking.
- Take the medicines your doctor told you to take with a small sip of water.
- Bring your cane, walker, or wheelchair if you have one already. Also bring shoes with flat, nonskid soles.

- Follow instructions about when to arrive at the hospital. Arrive on time.

After the Procedure

Your doctor or nurse will ask you to get up and walk around as soon as your anesthesia wears off. Most people go home the day of surgery. Do NOT drive yourself home.

Follow instructions about how to care for yourself at home.

Outlook (Prognosis)

Most people have pain relief and can move better after surgery. Numbness and tingling should get better or disappear. Your pain, numbness, or weakness may not get better or go away if you had nerve damage before surgery, or if you have symptoms caused by other spinal conditions.

Further changes may occur in your spine over time and new symptoms may occur.

Talk with your doctor about how to prevent future back problems.

Alternative Names

Spinal microdiscectomy; Microdecompression; Laminotomy; Disk removal; Spine surgery - discectomy; Discectomy

References

Ehni BL, Satyan K. Lumbar discectomy. In: Benzel EC, ed. *Spine Surgery*. 3rd ed. Philadelphia, PA: Elsevier Saunders; 2012:chap 78.

Gardocki RJ, Park AL. Lower back pain and disorders of intervertebral discs. In: Canale ST, Beaty JH, eds. *Campbell's Operative Orthopaedics*. 12th ed. Philadelphia, PA: Elsevier Mosby; 2013:chap 42.

Wilson AS, Samartzis D, Shen FH. Anterior cervical discectomy and fusion. In: Shen FH, Samartzis D, Fessler RG, eds. *Textbook of the Cervical Spine*. Philadelphia, PA: Elsevier Saunders; 2015:chap 30.

Review Date 5/9/2015

Updated by: C. Benjamin Ma, MD, Assistant Professor, Chief, Sports Medicine and Shoulder Service, UCSF Department of Orthopaedic Surgery, San Francisco, CA. Also reviewed by David Zieve, MD, MHA, Isla Ogilvie, PhD, and the A.D.A.M. Editorial team.



A.D.A.M., Inc. is accredited by URAC, also known as the American Accreditation HealthCare Commission (www.uran.org). URAC's [accreditation program](#) is an independent audit to verify that A.D.A.M. follows rigorous standards of quality and accountability. A.D.A.M. is among the first to achieve this important distinction for online health information and services. Learn more about A.D.A.M.'s [editorial policy](#), [editorial process](#) and [privacy policy](#). A.D.A.M. is also a founding member of Hi-Ethics and subscribes to the principles of the Health on the Net Foundation (www.hon.ch).

The information provided herein should not be used during any medical emergency or for the diagnosis or treatment of any medical condition. A licensed physician should be consulted for diagnosis and treatment of any and all medical conditions. Call 911 for all medical emergencies. Links to other sites are provided for information only -- they do not constitute endorsements of those other sites. Copyright 1997-2016, A.D.A.M., Inc. Duplication for commercial use must be authorized in writing by ADAM Health Solutions.



U.S. National Library of Medicine 8600 Rockville Pike, Bethesda, MD 20894

U.S. Department of Health and Human Services National Institutes of Health

Page last updated: 05 October 2016