

[U.S. Department of Health & Human Services](#)

[NIH National Institutes of Health](#)

[Contact Us](#)

[Get Email Alerts](#)

Font Size



[Accessible Search Form](#)

[NHLBI Entire Site](#)



Classification of Overweight and Obesity by BMI, Waist Circumference, and Associated Disease Risks

	BMI (kg/m ²)	Obesity Class	Disease Risk* Relative to Normal Weight and Waist Circumference	
			Men 102 cm (40 in) or less Women 88 cm (35 in) or less	Men > 102 cm (40 in) Women > 88 cm (35 in)
Underweight	< 18.5		-	-
Normal	18.5–24.9		-	-
Overweight	25.0–29.9		Increased	High
Obesity	30.0–34.9	I	High	Very High
	35.0–39.9	II	Very High	Very High
Extreme Obesity	40.0 +	III	Extremely High	Extremely High

* Disease risk for type 2 diabetes, hypertension, and CVD.
 + Increased waist circumference also can be a marker for increased risk, even in persons of normal weight.

[Home](#)

[Assessing Your Weight and Health Risk](#)

[Control Your Weight](#)

[Eat Right](#)

[Be Physically Active](#)

[Healthy Weight Tools](#)

- [BMI Calculator](#)
- [Menu Plans](#)
- [Portion Distortion](#)

[Key Recommendations](#)

[Healthy Weight Resources for Parents/Families \(We Can!\)](#)

[Health Professional Resources](#)

[Skip footer links and go to content](#)

[Twitter](#) [Facebook](#) [YouTube](#) [Google+](#)

[SITE INDEX](#) | [ACCESSIBILITY](#) | [PRIVACY STATEMENT](#) | [FOIA](#) | [No Fear Act](#) | [OIG](#) | [CONTACT US](#)
[National Institutes of Health](#) [Department of Health and Human Services](#) [USA.gov](#)