

## Global Assessment of Functioning (GAF) Scale

(From DSM-IV-TR, p. 34.)

Consider psychological, social, and occupational functioning on a hypothetical continuum of mental health-illness. Do not include impairment in functioning due to physical (or environmental) limitations.

Code	(Note: Use intermediate codes when appropriate, e.g., 45, 68, 72.)
100   91	<b>Superior functioning in a wide range of activities, life's problems never seem to get out of hand, is sought out by others because of his or her many positive qualities. No symptoms.</b>
90   81	<b>Absent or minimal symptoms</b> (e.g., mild anxiety before an exam), <b>good functioning in all areas, interested and involved in a wide range of activities. socially effective, generally satisfied with life, no more than everyday problems or concerns</b> (e.g. an occasional argument with family members).
80   71	<b>If symptoms are present, they are transient and expectable reactions to psychosocial stressors</b> (e.g., difficulty concentrating after family argument); <b>no more than slight impairment in social, occupational or school functioning</b> (e.g., temporarily failing behind in schoolwork).
70   61	<b>Some mild symptoms</b> (e.g. depressed mood and mild insomnia) <b>OR some difficulty in social, occupational, or school functioning</b> (e.g., occasional truancy, or theft within the household), <b>but generally functioning pretty well, has some meaningful interpersonal relationships.</b>
60   51	<b>Moderate symptoms</b> (e.g., flat affect and circumstantial speech, occasional panic attacks) <b>OR moderate difficulty in social, occupational, or school functioning</b> (e.g., few friends, conflicts with peers or co-workers).
50   41	<b>Serious symptoms</b> (e.g., suicidal ideation, severe obsessional rituals, frequent shoplifting) <b>OR any serious impairment in social, occupational, or school functioning</b> (e.g., no friends, unable to keep a job).
40   31	<b>Some impairment in reality testing or communication</b> (e.g., speech is at times illogical, obscure, or irrelevant) <b>OR major impairment in several areas, such as work or school, family relations, judgment, thinking, or mood</b> (e.g., depressed man avoids friends, neglects family, and is unable to work; child frequently beats up younger children, is defiant at home, and is failing at school).
30   21	<b>Behavior is considerably influenced by delusions or hallucinations</b> <b>OR serious impairment in communication or judgment</b> (e.g., sometimes incoherent, acts grossly inappropriately, suicidal preoccupation) <b>OR inability to function in almost all areas</b> (e.g., stays in bed all day; no job, home, or friends).
20   11	<b>Some danger of hurting self or others</b> (e.g., suicide attempts without clear expectation of death; frequently violent; manic excitement) <b>OR occasionally fails to maintain minimal personal hygiene</b> (e.g., smears feces) <b>OR gross impairment in communication</b> (e.g., largely incoherent or mute).
10   1	<b>Persistent danger of severely hurting self or others</b> (e.g., recurrent violence) <b>OR persistent inability to maintain minimal personal hygiene</b> <b>OR serious suicidal act with clear expectation of death.</b>
0	Inadequate information.