



Find a Doctor | Appointments | Login to MyChart

ABOUT HEALTH PATIENT CARE RESEARCH EDUCATION

FOLLOW [social media icons]

Health Library

to the diagnosis, treatment and prevention of conditions from A to Z.

PRINT THIS PAGE [print icon]


Home > Health Library

Medial Epicondylitis (Golfer's and Baseball Elbow)

<< Back to Diseases and Conditions

What is medial epicondylitis?

Medial epicondylitis, also known as golfer's elbow, baseball elbow, suitcase elbow, or forehand tennis elbow, is characterized by pain from the elbow to the wrist on the inside (medial side) of the elbow. The pain is caused by damage to the tendons that bend the wrist toward the palm. A tendon is a tough cord of tissue that connects muscles to bones.



Click Image to Enlarge

What causes medial epicondylitis?

Medial epicondylitis is caused by the excessive force used to bend the wrist toward the palm, such as swinging a golf club or pitching a baseball. Other possible causes of medial epicondylitis include the following:

- Serving with great force in tennis or using a spin serve
- Weak shoulder and wrist muscles

Experience Our Care

Find a Doctor Who Specializes in...

- Ankle Injuries
- Baseball Injuries
- Hip Injuries
- Knee Injuries
- Lateral Collateral Ligament Tears (LCL) Injury

See More +

Find a physician at another Johns Hopkins Member Hospital:

- Howard County General
- Sibley Memorial Hospital
- Suburban Hospital

Connect with a



Health Information

Search entire library by keyword

OR

Choose by letter to browse topics

A B C D E F
G H I J K L M
N O P Q R S
T U V W X Y
Z

(A-Z listing includes diseases, conditions, tests and procedures)

Diseases and Conditions

Tests and Procedures

Recipes

- [Nutrition Information](#)
- [Prevention Guidelines](#)

- Using a too tightly strung, too short, and/or too heavy tennis racket
- Throwing a javelin
- Carrying a heavy suitcase
- Chopping wood with an ax
- Operating a chain saw
- Frequent use of other hand tools on a continuous basis

- Treatment Center**
- Orthopaedic Surgery
 - Sports Medicine
 - Sports Medicine - JHCP
- Find Additional Treatment Centers at:
- Howard County General
 - Sibley Memorial Hospital
 - Suburban Hospital

Main Menu
Elbow


Medial Epicondylitis (Golfer's Elbow)

Play Video

Print Brochure

Share

Last Reviewed 10/08/14



Languages
Disclaimer
Enlarge

Maryland Patients


Find a location near you.

Traveling for Care?

Whether you're crossing the country or the globe, we make it easy to access world-class care at Johns Hopkins.

Outside of Maryland (toll free)
410-464-6713
 Request an Appointment Medical Concierge Services

International Patients
+1-410-502-7683
 Request an Appointment Medical Concierge Services



What are the symptoms of medial epicondylitis?

The following are the most common symptoms of medial epicondylitis. However, each individual may experience symptoms differently.

The most common symptom of medial epicondylitis is pain along the palm side of the forearm, from the elbow to the wrist, on the same side as the little finger. The pain can be felt when bending the wrist toward the palm against resistance, or when squeezing a rubber ball.

The symptoms of medial epicondylitis may resemble other medical problems or conditions. Always consult your doctor for

a diagnosis.

How is medial epicondylitis diagnosed?

The diagnosis of medial epicondylitis usually can be made based on a physical examination. The doctor may rest the arm on a table, palm side up, and ask the patient to raise the hand by bending the wrist against resistance. If a person has medial epicondylitis, pain usually is felt in the inner aspect of the elbow.

Treatment for medial epicondylitis

Specific treatment for medial epicondylitis will be determined by your doctor based on:

- Your age, overall health, and medical history
- Extent of the condition
- Your tolerance for specific medications, procedures, and therapies
- Expectation for the course of the condition
- Your opinion or preference

Treatment for medial epicondylitis includes stopping the activity that produces the symptoms. It is important to avoid the movement that caused the condition in the first place.

Treatment may include:

- Ice pack application (to reduce inflammation)
- Strengthening exercises
- Anti-inflammatory medications
- Bracing
- Corticosteroid injections
- Surgery (rarely necessary)

Discover Our Research

Death Risk For Marathoners Remains Low During Or Soon After Race

Low-carb, Higher-fat Diets Add No Arterial Health Risks To Obese People Seeking To Lose Weight

Expand Your Knowledge

- Sign up for an e-Newsletter
- Attend a Health Seminar



Related Articles

Tailor-made Treatment For Acl Tears

Spring Into Shape!

We've put together a team of professionals to help you break...

Building Rapport Is The Name Of The Game

As a team physician for the Baltimore Orioles, John Wilckens...

Related Videos

Sudden Cardiac Death

Sudden cardiac death kills more than 3,000 young people each year,



many of...

FOLLOW JOHNS HOPKINS MEDICINE



Contact Information

View our [phone directory](#) or find a [patient care location](#).

Privacy Policy

[Notice of Privacy Practices](#)
(Patients & Health Plan Members)

[Privacy Policy and Disclaimer](#)

© The Johns Hopkins University, The Johns Hopkins Hospital, and Johns Hopkins Health System. All rights reserved.

[Legal Disclaimer](#)