



3 Herbs that Beat Anxiety

Doctors Reveal 1 Weird Compound to Calm Anxiety that May Surprise You



ADVERTISEMENT

Home › Drugs A to Z › Risperidone

Print Share

Risperidone



Generic Name: risperidone (oral) (ris PER i done)

Brand Names: *RisperDAL*, *RisperDAL M-Tab*

Bipolar I Disorder

Nationally Recognized Expert Care Call and Learn More

Now



Overview

Side Effects

Dosage

Interactions

Professional

More

What is risperidone?

Risperidone is a antipsychotic medicine. It works by changing the effects of chemicals in the brain.

Risperidone is used to treat [schizophrenia](#) and symptoms of [bipolar disorder](#) (manic depression). It is also used in autistic children to treat symptoms of irritability.

Risperidone may also be used for purposes not listed in this medication guide.

ADVERTISEMENT



07/28/2015



Important information

Risperidone is not approved for use in psychotic conditions related to dementia. It may increase the risk of death in older adults with dementia-related conditions.

Do not give risperidone to a child without a doctor's advice.

While you are taking risperidone, you may be more sensitive to temperature extremes such as very hot or cold conditions. Avoid getting too cold, or becoming overheated or dehydrated. Drink plenty of fluids, especially in hot weather and during exercise. It is easier to become dangerously overheated and dehydrated while you are taking this medication.

Risperidone may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be alert. Drinking alcohol can increase certain side effects of risperidone.

Stop using this medicine and call your doctor at once if you have fever, stiff muscles, confusion, sweating, fast or uneven heartbeats, restless muscle movements in your face or neck, tremor (uncontrolled shaking), trouble swallowing, feeling light-headed, or fainting.

Before taking this medicine

You should not use risperidone if you are allergic to it.

Risperidone is not approved for use in psychotic conditions related to dementia. Risperidone may increase the risk of death in older adults with dementia-related conditions.

To make sure this medicine is safe for you, tell your doctor if you have:

Slideshow: Can Prescription Drugs Lead to Weight Gain?



- heart disease, high blood pressure, heart rhythm problems;
- coronary artery disease or recent heart attack;
- diabetes (or risk factors such as obesity or family history of diabetes);
- a history of low white blood cell (WBC) counts;
- liver or kidney disease;
- a history of seizures;
- a history of breast cancer;
- low bone mineral density;
- trouble swallowing;
- Parkinson's disease;
- if you are dehydrated; or
- if you also take blood pressure medicine.

Some people with mental illness have thoughts about suicide. Your doctor will need to check your progress at regular visits while you are using risperidone. Your family or other caregivers should also be alert to changes in your mood or symptoms.

The risperidone orally disintegrating tablet may contain phenylalanine. Talk to your doctor before using orally disintegrating tablets if you have phenylketonuria (PKU).

FDA pregnancy category C. It is not known whether risperidone will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant while taking risperidone or within 12 weeks after you stop taking this medicine.

See also: [Pregnancy and breastfeeding warnings](#) (in more detail)

Taking antipsychotic medication during the last 3 months of pregnancy may cause problems in the newborn, such as withdrawal symptoms, breathing problems, feeding problems, fussiness, tremors, and limp or stiff muscles. However, you may have withdrawal symptoms or other problems if you stop taking your medicine during pregnancy. **If you become pregnant while taking this medicine, do not stop taking it without your doctor's advice.**

Risperidone can pass into breast milk and may harm a nursing baby. **Do not breast-feed while taking this medicine and for at least 12 weeks after your treatment ends.**

Do not give this medicine to a child without a doctor's advice.





How should I take risperidone?

Take risperidone exactly as prescribed by your doctor. Follow all directions on your prescription label. Do not take this medicine in larger or smaller amounts or for longer than recommended.

Risperidone can be taken with or without food.

To take the orally disintegrating tablet (Risperdal M-Tabs):

- Keep the tablet in its blister pack until you are ready to take it. Open the package and peel back the foil. Do not push a tablet through the foil or you may damage the tablet.
- Use dry hands to remove the tablet and place it in your mouth.
- Do not swallow the tablet whole. Allow it to dissolve in your mouth without chewing. If desired, you may drink liquid to help swallow the dissolved tablet.

Measure liquid medicine with the dosing syringe provided, or with a special dose-measuring spoon or medicine cup. If you do not have a dose-measuring device, ask your pharmacist for one.

Use risperidone regularly to get the most benefit. Get your prescription refilled before you run out of medicine completely.

Do not mix the **liquid medicine** with cola or tea.

It may take up to several weeks before your symptoms improve. Keep using the medication as directed and tell your doctor if your symptoms do not improve.

Store at room temperature away from moisture, heat, and light. Do not liquid medicine to freeze.

What happens if I miss a dose?

Take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not take extra medicine to make up the missed dose.

07/28/2015

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include severe drowsiness, fast heart rate, feeling light-headed, fainting, and restless muscle movements in your eyes, tongue, jaw, or neck.

What should I avoid while taking risperidone?

Risperidone may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be alert.

Avoid getting up too fast from a sitting or lying position, or you may feel dizzy. Get up slowly and steady yourself to prevent a fall.

Drinking alcohol can increase certain side effects of risperidone.

While you are taking this medicine, you may be more sensitive to temperature extremes such as very hot or cold conditions. Avoid getting too cold, or becoming overheated or dehydrated. Drink plenty of fluids, especially in hot weather and during exercise.

Risperidone side effects

Get emergency medical help if you have any of these **signs of an allergic reaction to risperidone**: hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Call your doctor at once if you have:

- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement);

- breast swelling or tenderness (in men or women), nipple discharge, impotence, lack of interest in sex, missed menstrual periods;
- severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out;
- low white blood cells--sudden weakness or ill feeling, fever, chills, sore throat, mouth sores, red or swollen gums, trouble swallowing, skin sores, cold or flu symptoms, cough, trouble breathing;
- low levels of platelets in your blood--easy bruising, unusual bleeding (nose, mouth, vagina, or rectum), purple or red pinpoint spots under your skin;
- high blood sugar--increased thirst, increased urination, hunger, dry mouth, fruity breath odor, drowsiness, dry skin, blurred vision, weight loss; or
- penis erection that is painful or lasts 4 hours or longer.

Common risperidone side effects may include:

- headache;
- dizziness, drowsiness, tired feeling;
- tremors, twitching or uncontrollable muscle movements;
- agitation, anxiety, restless feeling;
- depressed mood;
- dry mouth, upset stomach, constipation;
- weight gain; or
- pain in your arms or legs.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

See also: [Side effects](#) (in more detail)

What other drugs will affect risperidone?

Taking this medicine with other drugs that make you sleepy can worsen this effect. Ask your doctor before taking risperidone with a sleeping pill, narcotic pain medicine, muscle relaxer, or medicine for anxiety, depression, or seizures.

Other drugs may interact with risperidone, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell each of your health care providers about all medicines you use now and any medicine you start or stop using.

Next → [Side Effects](#)

[Add to My Med List](#)

More about risperidone

- [Side Effects](#)
- [During Pregnancy or Breastfeeding](#)
- [Dosage Information](#)
- [Drug Images](#)
- [Drug Interactions](#)
- [Support Group](#)
- [Pricing & Coupons](#)
- [En Espanol](#)
- [197 Reviews - Add your own review/rating](#)

Consumer resources

- [Risperidone](#)
- [Risperidone orally disintegrating tablets](#)
- [Risperidone solution](#)
- [Risperidone tablets](#)
- [Risperidone \(Advanced Reading\)](#)
- [Risperidone Intramuscular \(Advanced Reading\)](#)
- *Other brands: [Risperdal](#), [Risperdal Consta](#), [Risperdal M-Tab](#)*

Professional resources

- [Risperidone \(AHFS Monograph\)](#)
- [More \(4\) »](#)

Related treatment guides

- [Bipolar Disorder](#)
- [Schizoaffective Disorder](#)
- [Borderline Personality Disorder](#)
- [Asperger Syndrome](#)
- [Autism](#)
- [More \(12\) »](#)

Where can I get more information?

- Your pharmacist can provide more information about risperidone.

3 Herbs that Beat Anxiety

Doctors Reveal 1 Weird Compound to Calm Anxiety that May Surprise You

View



Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Disclaimer: Every effort has been made to ensure that the information provided by Cerner Multum, Inc. ('Multum') is accurate, up-to-date, and complete, but no guarantee is made to that effect. Drug information contained herein may be time sensitive. Multum information has been compiled for use by healthcare practitioners and consumers in the United States and therefore Multum does not warrant that uses outside of the United States are appropriate, unless specifically indicated otherwise. Multum's drug information does not endorse drugs, diagnose patients or recommend therapy. Multum's drug information is an informational resource designed to assist licensed healthcare practitioners in caring for their patients and/or to serve consumers viewing this service as a supplement to, and not a substitute for, the expertise, skill, knowledge and judgment of healthcare practitioners. The absence of a warning for a given drug or drug combination in no way should be construed to indicate that the drug or drug combination is safe, effective or appropriate for any given patient. Multum does not assume any responsibility for any aspect of healthcare administered with the aid of information Multum provides. The information contained herein is not intended to cover all possible uses, directions, precautions, warnings, drug interactions, allergic reactions, or adverse effects. If you have questions about the drugs you are taking, check with your doctor, nurse or pharmacist.

Copyright 1996-2015 Cerner Multum, Inc. Version: 20.02. Revision Date: 2014-06-16, 8:51:54 AM.

DRUG STATUS



Availability
Prescription only



Pregnancy Category
Risk cannot be ruled out



CSA Schedule
Not a controlled drug





Approval History
Drug history at FDA



Drug Class

[Atypical antipsychotics](#)

Related Drugs

Bipolar Disorder

[Seroquel](#), [lithium](#),
[clonazepam](#), [Abilify](#),
[Klonopin](#), [Lamictal](#),
More...

Schizoaffective Disorder

[Seroquel](#), [Prozac](#),
[lithium](#), [Abilify](#),
[fluoxetine](#), [Lamictal](#),
More...

Schizophrenia

[Seroquel](#), [Abilify](#),
[quetiapine](#),
[olanzapine](#), [Zyprexa](#),
[Latuda](#), More...

Autism

[Abilify](#), [venlafaxine](#),
[Effexor](#), [Risperdal](#),
[Geodon](#), [aripiprazole](#),
More...

[2 more conditions...](#)

ADVERTISEMENT

The Depression Treatment Device

Fisher Wallace Stimulator
212 688 8100
fisherwallace.com

20

Use for 20 minutes, once or twice a day.

Risperidone Rating

[197 User Reviews](#)

6.7/10



Risperidone Images



Risperidone 1 mg



[View all 157 images](#)

Get Updates by Email

Sign up for daily, weekly or monthly emails to receive news updates to your inbox!

News and Warnings related to this drug

FDA Medwatch Alerts for all Medications

News Roundup **Daily**

Related News and Articles

Psychiatric Woes Often Drive Euthanasia Requests in Belgium

July 28, 2015

Health Tip: Protect Your Heart From Stress and Depression

July 28, 2015

Health Tip: Offer Support to Seniors

July 28, 2015



Harvard Health Publications

Related: [Bipolar Disorder \(Manic Depressive Illness or Manic Depression\)](#)



ADVERTISEMENT

Risperidone

 [Pronunciation](#)

Drugs.com Mobile Apps

The easiest way to lookup drug information, identify pills, check interactions and set up your own personal medication records. Available for Android and iOS devices.

[Explore Apps](#)



SUPPORT

[Help Center](#)
[Frequent Questions](#)
[Sitemap](#)
[Contact Us](#)

ABOUT

[About Drugs.com](#)
[Advertising Policy](#)
[Content Submissions](#)
[Drugs.com Blog](#)

TERMS & PRIVACY


[Editorial Policy](#)
[Privacy Policy](#)
[Terms of Use](#)
[Attribution & Citations](#)

CONNECT



[Subscribe](#) to receive email notifications whenever new articles are published.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex® (updated July 13th, 2015), Cerner Multum™ (updated July 2nd, 2015), Wolters Kluwer™ (updated July 2nd, 2015) and others. To view content sources and attributions, please refer to our [editorial policy](#).



We comply with the HONcode standard for trustworthy health information - [verify here](#)

Copyright © 2000-2015 Drugs.com. All rights reserved.