

 **Results** **Safety Profile** **Experience** **Support**[Home](#) > [Conditions](#) > [Pain](#) > [Flector Patch](#)[Print](#) [Share](#)

Flector Patch



Generic Name: diclofenac transdermal (dye KLOE fen ak)

Brand Names: *Flector Patch*

Relapsed Multiple Myeloma - Treatment Option

Get More Prescribing and Dosing Information at the Official Physician Website. Go to Prescription treatment website



Overview

Side Effects

Dosage

Interactions

Professional

More

What is Flector?

Flector Patches contain diclofenac, a nonsteroidal anti-inflammatory drug (NSAID). Diclofenac works by reducing substances in the body that cause pain and inflammation.

Flector is used to treat pain caused by minor [sprains](#), strains, or bruising.

Flector may also be used for other purposes not listed in this medication guide.

ADVERTISEMENT



Important information

Diclofenac can increase your risk of fatal heart attack or stroke, especially if you use it long term or take high doses, or if you have heart disease. Do not use Flector just before or after heart bypass surgery (coronary artery bypass graft, or CABG).

Get emergency medical help if you have chest pain, weakness, shortness of breath, slurred speech, or problems with vision or balance while using Flector Patches.

Before using Flector, tell your doctor if you are allergic to any drugs, or if you have heart disease, congestive heart failure, high blood pressure, a history of heart attack or stroke, a history of stomach ulcer or bleeding, liver or kidney disease, a blood clotting disorder, asthma or polyps in your nose, or if you smoke.

While the risk of absorbing diclofenac into your bloodstream is low, an NSAID may cause life-threatening heart or circulation problems such as heart attack or stroke, especially if you use it long term.

Flector may also cause serious effects on the stomach or intestines, including bleeding or perforation (forming of a hole). These conditions can be fatal and can occur without warning while you are using Flector, especially in older adults.

Call your doctor at once if you have symptoms of stomach bleeding such as black, bloody, or tarry stools, or coughing up blood or vomit that looks like coffee grounds.

Before taking this medicine

The diclofenac in Flector patches can increase your risk of fatal heart attack or stroke, especially if you use it long term or take high doses, or if you have heart disease. Even people without heart disease or risk factors could have a stroke or heart attack while using Flector patches.

Do not use Flector just before or after heart bypass surgery (coronary artery bypass graft, or CABG).

Diclofenac may also cause stomach or intestinal bleeding, which can be fatal. These conditions can occur without warning while you are using Flector, especially in older adults.

3 Fatigue-Causing Foods

Boost Your Energy Level Every Day By Never Eating These 3 Foods

VitalReds



SLIDESHOW

[Tramadol: 9 Things You Should Know](#)

You should not use Flector if you are allergic to diclofenac (Voltaren, Cataflam, Cambia, and others), or if you have ever had an asthma attack or severe allergic reaction after taking aspirin or an NSAID.

To make sure Flector is safe for you, tell your doctor if you have:

- heart disease, high blood pressure, high cholesterol, diabetes, or if you smoke;
- a history of heart attack, stroke, or blood clot;
- a history of stomach ulcers or bleeding;
- asthma;
- liver or kidney disease;
- fluid retention.

Using Flector during the last 3 months of pregnancy may harm the unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

It is not known whether diclofenac transdermal passes into breast milk or if it could harm a nursing baby. You should not breast-feed while using Flector.

Flector is not approved for use by anyone younger than 18 years old.

ADVERTISEMENT

Ads by



How should I use Flector Patches?

Use Flector Patches exactly as prescribed by your doctor. Follow all directions on your prescription label. Do not use this medicine in larger amounts or for longer than recommended. Using extra skin patches will not make the medicine more effective, and could cause dangerous side effects.

Apply the skin patch directly to the area of pain. The skin patch can be worn for up to 12 hours and then removed. Apply a new patch at that time if pain continues.

Do not apply Flector on an open skin wound, or on areas of eczema, infection, skin rash, or burn injury.

Wash your hands after applying or removing a Flector skin patch.

If the patch does not stick well, you may place medical tape around the edges. You may also use a mesh netting sleeve to hold a patch in place on your skin. Do not cover the patch with a bandage or other covering that does not allow air to pass through.

After removing a skin patch fold it in half, sticky side in, and throw it away in a place where children or pets cannot get to it. Keep both used and unused skin patches out of the reach of children or pets.

If you use Flector long-term, you may need frequent medical tests.

The Flector patch may burn your skin if you wear the patch during an MRI (magnetic resonance imaging). Remove the patch before undergoing such a test.

Store Flector at room temperature away from moisture and heat. Reseal the storage envelope each time you remove a patch from it.

Read all patient information, medication guides, and instruction sheets provided to you. Ask your doctor or pharmacist if you have any questions.

See also: [Dosage Information \(in more detail\)](#)

What happens if I miss a dose?

Apply a skin patch as soon as you remember, and wear it for 12 hours before applying a new one. Do not use extra patches to make up the missed dose. Do not wear a Flector skin patch for longer than 12 hours.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while using Flector?

Do not wear a Flector skin patch while taking a bath or shower or while swimming.

Avoid getting a skin patch near your eyes, nose, or mouth. If this does happen, rinse with water. Call your doctor if you have eye irritation that lasts longer than 1 hour.

Avoid drinking alcohol. It may increase your risk of stomach bleeding.

Avoid taking aspirin or other NSAIDs.

Ask a doctor or pharmacist before using any cold, allergy, or pain medication. Many medicines available over the counter contain aspirin or other medicines similar to diclofenac. Taking certain products together can cause you to get too much of this type of medication. Check the label to see if a medicine contains aspirin, ibuprofen, ketoprofen, or naproxen.

Ads by

Flector side effects

Get emergency medical help if you have signs of an allergic reaction to Flector: sneezing, runny or stuffy nose; wheezing or trouble breathing; hives; swelling of your face, lips, tongue, or throat.

Get emergency medical help if you have signs of a heart attack or stroke: chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath.

Stop using Flector patches and call your doctor at once if you have:

- the first sign of any skin rash, no matter how mild;
- shortness of breath (even with mild exertion);
- swelling or rapid weight gain;
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds;
- liver problems--nausea, upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes);
- kidney problems--little or no urinating, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath;
- low red blood cells (anemia)--pale skin, feeling light-headed or short of breath, rapid heart rate, trouble concentrating; or
- severe skin reaction--fever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling.

Common Flector side effects may include:

- heartburn, gas, stomach pain, nausea, vomiting;
- diarrhea, constipation; or
- mild itching, burning, redness, or other skin irritation where the patch was worn.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

See also: [Side effects \(in more detail\)](#)

What other drugs will affect Flector?

Ask your doctor before using Flector if you take an antidepressant such as citalopram, escitalopram, fluoxetine (Prozac), fluvoxamine, paroxetine, sertraline (Zoloft), trazodone, or vilazodone. Taking any of these medicines with an NSAID may cause you to bruise or bleed easily.

Tell your doctor about all your current medicines and any you start or stop using, especially:

- cyclosporine;
- lithium;
- methotrexate;
- a blood thinner (warfarin, Coumadin, Jantoven);
- heart or blood pressure medication, including a diuretic or "water pill"; or
- steroid medicine (prednisone and others).

This list is not complete. Other drugs may interact with diclofenac, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible interactions are listed in this medication guide.

Next → [Side Effects](#)

Celecoxib - \$35.75

GoodRx can save you up to 80% on your Celecoxib prescription. Go to goodrx.com



ADVERTISEMENT



Add to My Med List

More about Flector Patch (diclofenac topical)

- [Side Effects](#)
- [During Pregnancy](#) or [Breastfeeding](#)
- [Dosage Information](#)
- [Drug Interactions](#)
- [Support Group](#)

- [Pricing & Coupons](#)
- [En Español](#)
- [52 Reviews – Add your own review/rating](#)
- [Drug class: topical non-steroidal anti-inflammatories](#)

Consumer resources

- [Flector patch](#)
- [Flector Topical](#) (Advanced Reading)

Other brands: [Voltaren Gel](#), [Pennsaid](#), [Solaraze](#), [Klofensaid II](#)

Professional resources

- [Flector](#) (FDA)
- [Diclofenac Sodium topical](#) (AHFS Monograph)

Related treatment guides

- [Pain](#)

Where can I get more information?

- Your pharmacist can provide more information about Flector.
- Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Flector only for the indication prescribed.
- Disclaimer: Every effort has been made to ensure that the information provided is accurate, up-to-date, and complete, but no guarantee is made to that effect. Multum's drug information is an informational resource designed to assist licensed healthcare practitioners in caring for their patients and/or to serve consumers viewing this service as a supplement to, and not a substitute for, the expertise, skill, knowledge and judgment of healthcare practitioners. The information contained herein is not intended to cover all possible uses, directions, precautions, warnings, drug interactions, allergic reactions, or adverse effects. If you have questions about the drugs you are taking, check with your doctor, nurse or pharmacist.

Copyright 1996-2017 Cerner Multum, Inc. Version: 4.01. Revision Date: 2015-09-02, 12:11:11 PM.

DRUG STATUS



Availability
Prescription only



Pregnancy Category
Risk depends on usage



CSA Schedule
Not a controlled drug





Approval History
Drug history at FDA



Drug Class

[Topical non-steroidal anti-inflammatories](#)

ADVERTISEMENT



Angel MedFlight
WORLDWIDE AIR AMBULANCE



**SUPERIOR
SAFETY**

**EXCEPTIONAL
CARE**

855.215.8460

Related Drugs

Pain

[tramadol](#), [acetaminophen](#), [Tylenol](#), [oxycodone](#), [naproxen](#), [aspirin](#), [ibuprofen](#), [amitriptyline](#), [clonidine](#), [Percocet](#), [diclofenac topical](#), [hydrocodone](#), [Norco](#), [hydroxyzine](#), [Celebrex](#), [morphine](#), [fentanyl](#), [Dilaudid](#), [lidocaine topical](#), [Vicodin](#), [Voltaren](#), [methadone](#), [OxyContin](#), [More...](#)

Flector Patch Rating

[52 User Reviews](#)

7.8/10

Related News and Articles

[4 in 10 People Will Suffer Arthritic Hands Over Lifetime](#)

May 4, 2017

[April 29 Is National Prescription Drug Take Back Day](#)

April 27, 2017

[Inspirion Delivery Sciences Receives FDA Approval for RoxyBond \(oxycodone hydrochloride\) Immediate Release Tablets with Abuse-Deterrent Label Claims](#)

April 26, 2017



Harvard Health Publications

Related: [Pain](#)

ADVERTISEMENT

Drugs.com Mobile Apps

The easiest way to lookup drug information, identify pills, check interactions and set up your own personal medication records. Available for Android and iOS devices.

[Explore Apps](#)

SUPPORT

[Help Center](#)[Frequent Questions](#)[Sitemap](#)[Contact Us](#)

ABOUT

[About Drugs.com](#)[Advertising Policy](#)[Content Submissions](#)[Drugs.com Blog](#)

TERMS & PRIVACY

[Editorial Policy](#)[Privacy Policy](#)[Terms of Use](#)[Attribution & Citations](#)

[Subscribe](#) to receive email notifications whenever new articles are published.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex® (updated May 1st, 2017), Cerner Multum™ (updated May 3rd, 2017), Wolters Kluwer™ (updated May 2nd, 2017) and others. To view content sources and attributions, please refer to our [editorial policy](#).



We comply with the HONcode standard for trustworthy health information - [verify here](#)

Copyright © 2000-2017 Drugs.com. All rights reserved.