

What's important to you in a treatment for moderate to severe Rheumatoid Arthritis (RA)?

[Home](#) > [Conditions](#) > [Muscle Spasm](#) > [Skelaxin](#)

[Print](#) [Share](#)

Skelaxin



Generic Name: [metaxalone](#) (me TAX a lone)

Brand Names: *Skelaxin*

Relapsed Multiple Myeloma - Treatment Option

Get More Prescribing and Dosing Information at the Official Physician Website. Go to Prescription treatment website



[Overview](#)

[Side Effects](#)

[Dosage](#)

[Interactions](#)

[Professional](#)

[More](#)

What is Skelaxin?

Skelaxin (metaxalone) is a muscle relaxant. It works by blocking nerve impulses (or pain sensations) in the brain.

Skelaxin is used together with rest and physical therapy to treat skeletal muscle conditions such as pain or injury.

Skelaxin may also be used for purposes not listed in this medication guide.

ADVERTISEMENT





Important information

You should not use Skelaxin if you have anemia (low red blood cells), or severe kidney or liver disease. You may need a dose adjustment or special tests to safely take Skelaxin.

Skelaxin can cause side effects that may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be awake and alert. Avoid drinking alcohol. It can increase some of the side effects of metaxalone.

Before taking this medicine

You should not use Skelaxin if you are allergic to metaxalone, or if you have:

- anemia (low red blood cells);
- severe kidney disease; or
- severe liver disease.

To make sure Skelaxin is safe for you, tell your doctor if you have liver or kidney disease.

It is not known whether this medicine will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

It is not known whether metaxalone passes into breast milk or if it could harm a nursing baby. You should not breast-feed while using this medicine.

Skelaxin is not approved for use by anyone younger than 12 years old.


ADVERTISEMENT

Ads by

3

Harmful Foods

**3 harmful foods
that are the root
of all your
stomach
problems**



How should I take Skelaxin?

Take Skelaxin exactly as it was prescribed for you. Follow all directions on your prescription label. Do not take this medicine in larger or smaller amounts or for longer than recommended.

This medicine can cause unusual results with certain medical tests. Tell any doctor who treats you that you are using this medicine.

Skelaxin is only part of a complete treatment program that may also include rest, physical therapy, or other pain relief measures. Follow your doctor's instructions.

Store at room temperature away from moisture and heat.

Skelaxin dosing information

Usual Adult Dose for Muscle Spasm:

800 mg orally 3 to 4 times a day.

Usual Pediatric Dose for Muscle Spasm:

>= 12 years: 800 mg orally 3 to 4 times a day.

See also: [Dosage Information \(in more detail\)](#)

What happens if I miss a dose?

Take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not take extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. An overdose of metaxalone can be fatal.

What should I avoid while taking Skelaxin?

Do not drink alcohol. Dangerous side effects or death can occur when alcohol is combined with Skelaxin. Check

your food and medicine labels to be sure these products do not contain alcohol.

This medicine may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be alert.

Dizziness or drowsiness may be more likely in older adults.



Skelaxin side effects

Get emergency medical help if you have signs of an allergic reaction to Skelaxin: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Stop using Skelaxin and call your doctor at once if you have:

- weak or shallow breathing;
- a light-headed feeling, like you might pass out;
- pale or yellowed skin, dark colored urine, fever, confusion or weakness; or
- upper stomach pain, loss of appetite, jaundice (yellowing of the skin or eyes).

Common Skelaxin side effects may include:

- dizziness, drowsiness;
- nausea, vomiting, upset stomach;
- headache; or
- feeling nervous or irritable;

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

See also: [Side effects \(in more detail\)](#)

What other drugs will affect Skelaxin?

Taking Skelaxin with other drugs that make you sleepy or slow your breathing can cause dangerous or life-threatening side effects. Ask your doctor before taking a sleeping pill, narcotic pain medicine, prescription cough medicine, a muscle relaxer, or medicine for anxiety, depression, or seizures.

Other drugs may interact with metaxalone, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell each of your health care providers about all medicines you use now and any medicine you start or stop using.

Next → [Side Effects](#)

3 Fatigue-Causing Foods - Never eat these 3 foods

Boost your energy level every day by never eating these 3 foods Go to haveenergyatanyage.com



ADVERTISEMENT

 Add to My Med List

More about Skelaxin (metaxalone)

- [Side Effects](#)
 - [During Pregnancy](#)
 - [Dosage Information](#)
 - [Drug Images](#)
 - [Drug Interactions](#)
 - [Support Group](#)
 - [Pricing & Coupons](#)
 - [En Español](#)
 - [76 Reviews – Add your own review/rating](#)
 - [Generic Availability](#)
 - [Drug class: skeletal muscle relaxants](#)
- Consumer resources
- [Skelaxin](#)
 - [Skelaxin \(Advanced Reading\)](#)

- Professional resources
 - [Skelaxin](#) (AHFS Monograph)
 - [Skelaxin](#) (FDA)
- Related treatment guides
 - [Muscle Spasm](#)

Where can I get more information?

- Your pharmacist can provide more information about Skelaxin.
- Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Skelaxin only for the indication prescribed.
- Disclaimer: Every effort has been made to ensure that the information provided by Cerner Multum, Inc. ('Multum') is accurate, up-to-date, and complete, but no guarantee is made to that effect. Drug information contained herein may be time sensitive. Multum information has been compiled for use by healthcare practitioners and consumers in the United States and therefore Multum does not warrant that uses outside of the United States are appropriate, unless specifically indicated otherwise. Multum's drug information does not endorse drugs, diagnose patients or recommend therapy. Multum's drug information is an informational resource designed to assist licensed healthcare practitioners in caring for their patients and/or to serve consumers viewing this service as a supplement to, and not a substitute for, the expertise, skill, knowledge and judgment of healthcare practitioners. The absence of a warning for a given drug or drug combination in no way should be construed to indicate that the drug or drug combination is safe, effective or appropriate for any given patient. Multum does not assume any responsibility for any aspect of healthcare administered with the aid of information Multum provides. The information contained herein is not intended to cover all possible uses, directions, precautions, warnings, drug interactions, allergic reactions, or adverse effects. If you have questions about the drugs you are taking, check with your doctor, nurse or pharmacist.

Copyright 1996-2017 Cerner Multum, Inc. Version: 6.02. Revision Date: 2016-09-19, 1:18:48 PM.

DRUG STATUS

-  Availability
Prescription only
-  Pregnancy Category
Not classified
-  CSA Schedule
Not a controlled drug
-  Approval History
FDA approved 1962

Manufacturer

[King Pharmaceuticals, Inc.](#)

Drug Class

[Skeletal muscle relaxants](#)





WORLDWIDE AIR AMBULANCE

SUPERIOR SAFETY

EXCEPTIONAL CARE

855.215.8460

Related Drugs

Muscle Spasm

[cyclobenzaprine](#), [Soma](#), [baclofen](#), [Flexeril](#), [tizanidine](#), [diazepam](#), [Valium](#), [methocarbamol](#), [Robaxin](#), [Zanaflex](#), [carisoprodol](#), [metaxalone](#), [orphenadrine](#), [Norflex](#), [chlorzoxazone](#), [Amrix](#), [Lorzone](#), [Lioresal](#), [Parafon Forte DSC](#), [Robaxin-750](#), [Diastat](#), [Diastat AcuDial](#), [Norgesic](#), [Fexmid](#), [More...](#)

Skelaxin Rating

[76 User Reviews](#)

7.3/10

Skelaxin Images



Skelaxin (metaxalone) 800 mg

[View larger images](#)

Related News and Articles

[Brain Chip Helps Paralyzed 'Type' With Their Mind](#)

February 21, 2017

[Scientists Create Clothing With 'Knitted' Muscle Power](#)

January 26, 2017

[Forward-Thinking Tips for Back Pain](#)

January 18, 2017

ADVERTISEMENT

Drugs.com Mobile Apps

The easiest way to lookup drug information, identify pills, check interactions and set up your own personal medication records. Available for Android and iOS devices.



[Explore Apps](#)

SUPPORT

[Help Center](#)[Frequent Questions](#)[Sitemap](#)[Contact Us](#)

ABOUT

[About Drugs.com](#)[Advertising Policy](#)[Content Submissions](#)[Drugs.com Blog](#)

TERMS & PRIVACY

[Editorial Policy](#)[Privacy Policy](#)[Terms of Use](#)[Attribution & Citations](#)

[Subscribe](#) to receive email notifications whenever new articles are published.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex® (updated May 1st, 2017), Cerner Multum™ (updated May 3rd, 2017), Wolters Kluwer™ (updated May 2nd, 2017) and others. To view content sources and attributions, please refer to our [editorial policy](#).



We comply with the HONcode standard for trustworthy health information - [verify here](#)

Copyright © 2000-2017 Drugs.com. All rights reserved.