



[Home](#) > [Conditions](#) > [Muscle Spasm](#) > [Flexeril](#)

[Print](#) [Share](#)

Flexeril



Generic Name: [cyclobenzaprine](#) (syə kloə BEN zə preən)

Brand Names: *Amrix, Fexmid, Flexeril*

Relapsed Multiple Myeloma - Treatment Option

Get More Prescribing and Dosing Information at the Official Physician Website. Go to Prescription treatment website



[Overview](#)

[Side Effects](#)

[Dosage](#)

[Interactions](#)

[Patient Tips](#)

[Professional](#)

[More](#)

What is Flexeril?

Flexeril (cyclobenzaprine) is a muscle relaxant. It works by blocking nerve impulses (or pain sensations) that are sent to your brain.

Flexeril is used together with rest and physical therapy to treat skeletal muscle conditions such as pain or injury.

ADVERTISEMENT



Flexeril may also be used for other purposes not listed in this medication guide.

Important information

Do not take Flexeril if you have used an MAO inhibitor such as isocarboxazid (Marplan), tranylcypromine (Parnate), phenelzine (Nardil), or selegiline (Eldepryl, Emsam) within the past 14 days. Serious, life-threatening side effects can occur if you take Flexeril before the MAO inhibitor has cleared from your body.

You should not take Flexeril if you have recently had a heart attack, or if you have a heart rhythm disorder, congestive heart failure, heart block, or an overactive thyroid.

Flexeril can cause side effects that may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be awake and alert. Avoid drinking alcohol, which can increase some of the side effects of Flexeril.

Before taking this medicine


Do not take Flexeril if you have used an MAO inhibitor such as isocarboxazid (Marplan), tranylcypromine (Parnate), phenelzine (Nardil), or selegiline (Eldepryl, Emsam) within the past 14 days. Serious, life-threatening side effects can occur if you take Flexeril before the MAO inhibitor has cleared from your body. Do not use Flexeril if you have recently had a heart attack, or if you have:

- a heart rhythm disorder;
- congestive heart failure;
- heart block; or
- an overactive thyroid.

Before using Flexeril, tell your doctor if you are allergic to any drugs, or if you have:

- problems with urination;
- enlarged prostate;
- glaucoma; or
- liver disease.

If you have any of these conditions, you may need a dose adjustment or special tests to safely take Flexeril.



The Real Source of Your Fatigue

GUNDRY MD™

[\[Watch Now \]](#)




SLIDESHOW

[Flexeril \(cyclobenzaprine\): How to Safely Use This Muscle Relaxant](#)

FDA pregnancy category B. Flexeril is not expected to be harmful to an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant during treatment. It is not known whether cyclobenzaprine passes into breast milk or if it could harm a nursing baby. Do not use this medication without telling your doctor if you are breast-feeding a baby.

See also: [Pregnancy and breastfeeding warnings \(in more detail\)](#)

Older adults may be more sensitive to the side effects of Flexeril.



How should I take Flexeril?

Take this medication exactly as it was prescribed for you. Do not take Flexeril in larger amounts, or take it for longer than recommended by your doctor. Follow the directions on your prescription label.

Take this medicine with a full glass of water.

Flexeril is only part of a complete program of treatment that may also include rest, physical therapy, or other pain relief measures. Follow your doctor's instructions.

Store Flexeril at room temperature away from moisture, heat, and light.

See also: [Dosage Information \(in more detail\)](#)

What happens if I miss a dose?

Take the missed dose as soon as you remember. If it is almost time for your next dose, skip the missed dose and take the medicine at the next regularly scheduled time. Do not take extra medicine to make up the missed dose.


What happens if I overdose?

Seek emergency medical attention if you think you have used too much of this medicine. An overdose of Flexeril can be fatal.

Overdose symptoms may include drowsiness, fast heartbeat, tremors or shaking, slurred speech, confusion, nausea, vomiting, hallucinations (seeing things), chest pain, or seizure (convulsions).

What should I avoid?

Flexeril can cause side effects that may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be awake and alert. Avoid drinking alcohol, which can increase some of the side effects of Flexeril. Cold or allergy medicine, narcotic pain medicine, sleeping pills, and medicine for seizures, depression or anxiety can add to sleepiness caused by Flexeril. Tell your doctor if you regularly use any of these medicines, or any other muscle relaxer.



Flexeril side effects

Get emergency medical help if you have any of these signs of an allergic reaction to Flexeril: hives; difficulty breathing; swelling of your face, lips, tongue, or throat. Stop using Flexeril and call your doctor at once if you have any of these serious side effects:

- fast, pounding, or uneven heartbeats;
- chest pain or heavy feeling, pain spreading to the arm or shoulder, nausea, sweating, general ill feeling;
- sudden numbness or weakness, especially on one side of the body;
- sudden headache, confusion, problems with vision, speech, or balance;
- feeling light-headed, fainting;
- confusion, weakness, lack of coordination;
- nausea, stomach pain, low fever, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes);
- seizure (convulsions);
- unusual thoughts or behavior, hallucinations (seeing things); or
- easy bruising or bleeding, unusual weakness.

Less serious Flexeril side effects may include:

- dry mouth or throat;
- blurred vision;
- drowsiness, dizziness, tired feeling;
- loss of appetite, stomach pain, nausea;
- diarrhea, constipation, gas; or
- muscle weakness.

This is not a complete list of side effects and others may occur. Tell your doctor about any unusual or bothersome side effect. You may report side effects to FDA at 1-800-FDA-1088.

See also: [Side effects \(in more detail\)](#)

What other drugs will affect Flexeril?

Many drugs can interact with Flexeril. Below is just a partial list. Tell your doctor if you are using:

- atropine (Donnatal, and others), benztropine (Cogentin), dimenhydrinate (Dramamine), methscopolamine (Pamine), or scopolamine (Transderm-Scop);
- a bronchodilator such as ipratropium (Atrovent) or tiotropium (Spiriva);
- glycopyrrolate (Robinul);
- guanethidine (Ismelin);
- mepenzolate (Cantil);
- tramadol (Ultram);
- bladder or urinary medications such as darifenacin (Enablex), flavoxate (Urispas), oxybutynin (Ditropan, Oxytrol), tolterodine (Detrol), or solifenacin (Vesicare); or
- irritable bowel medications such as dicyclomine (Bentyl), hyoscyamine (Anaspaz, Cystospaz, Levsin, and others), or propantheline (Pro-Banthine).

This list is not complete and there may be other drugs that can interact with Flexeril. Tell your doctor about all your prescription and over-the-counter medications, vitamins, minerals, herbal products, and drugs prescribed by other doctors. Do not start a new medication without telling your doctor.

Next → [Side Effects](#)

Flush the bloat

Cut a bit of belly bloat each day by avoiding these 3 foods [Go to nucific.com](#)



[Add to My Med List](#)

More about Flexeril (cyclobenzaprine)

- [Side Effects](#)
- [During Pregnancy](#) or [Breastfeeding](#)
- [Dosage Information](#)
- [Drug Images](#)
- [Drug Interactions](#)
- [Support Group](#)
- [157 Reviews](#) – [Add your own review/rating](#)
- [Drug class: skeletal muscle relaxants](#)

Consumer resources

- [Flexeril](#)
- [Flexeril](#) (Advanced Reading)

Other brands: [Amrix](#), [Fexmid](#)

Professional resources

- [Flexeril](#) (AHFS Monograph)
- [Flexeril](#) (FDA)

Related treatment guides

- [Muscle Spasm](#)
- [Sciatica](#)

Where can I get more information?

- Your pharmacist can provide more information about Flexeril.
- Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Flexeril only for the indication prescribed.
- Disclaimer: Every effort has been made to ensure that the information provided is accurate, up-to-date, and complete, but no guarantee is made to that effect. Drug information contained herein may be time sensitive. Multum's drug information is an informational resource designed to assist licensed healthcare practitioners in caring for their patients and/or to serve consumers viewing this service as a supplement to, and not a substitute for, the expertise, skill, knowledge and judgment of healthcare practitioners. The absence of a warning for a given drug or drug combination in no way should be construed to indicate that the drug

or drug combination is safe, effective or appropriate for any given patient. The information contained herein is not intended to cover all possible uses, directions, precautions, warnings, drug interactions, allergic reactions, or adverse effects.

Copyright 1996-2017 Cerner Multum, Inc. Version: 3.11. Revision Date: 4/12/2009 4:38:46 PM.

DRUG STATUS



Availability
Prescription only



Pregnancy Category
No proven risk in humans



CSA Schedule
Not a controlled drug



Approval History
FDA approved 1977




Diagnosed with
ankylosing spondylitis?

[Learn about biologics](#) >

Drug Class

[Skeletal muscle relaxants](#)

ADVERTISEMENT



Related Drugs

Muscle Spasm

[cyclobenzaprine](#), [Soma](#), [baclofen](#), [tizanidine](#), [diazepam](#), [Valium](#), [methocarbamol](#), [Robaxin](#), [Zanaflex](#), [carisoprodol](#), [metaxalone](#), [Skelaxin](#), [More...](#)

Sciatica

[acetaminophen](#), [Tylenol](#), [naproxen](#), [cyclobenzaprine](#), [aspirin](#), [ibuprofen](#), [diclofenac](#), [Voltaren](#), [Paracetamol](#), [Motrin](#), [indomethacin](#), [Aleve](#), [More...](#)


Flexeril Rating

[157 User Reviews](#)

6.5/10



Flexeril Images



Flexeril (cyclobenzaprine) 10 mg

[View all 4 images](#)

Related News and Articles

['Mindfulness' Probably Won't Cure Your Back Pain: Study](#)

April 25, 2017

[Healing Hands: Massage May Ease Chronic Back Pain](#)

April 14, 2017

[Chiropractors Not Magicians When It Comes to Chronic Back Pain](#)

April 11, 2017

Diagnosed with
ankylosing spondylitis?

Learn about biologics >

ADVERTISEMENT

Drugs.com Mobile Apps

The easiest way to lookup drug information, identify pills, check interactions and set up your own personal medication records. Available for Android and iOS devices.

Explore Apps

SUPPORT

[Help Center](#)

[Frequent Questions](#)

[Sitemap](#)

[Contact Us](#)

ABOUT

[About Drugs.com](#)

[Advertising Policy](#)

[Content Submissions](#)

[Drugs.com Blog](#)

TERMS & PRIVACY

[Editorial Policy](#)

[Privacy Policy](#)

[Terms of Use](#)

[Attribution & Citations](#)



[Subscribe](#) to receive email notifications whenever new articles are published.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex® (updated May 1st, 2017), Cerner Multum™ (updated May 3rd, 2017), Wolters Kluwer™ (updated May 2nd, 2017) and others. To view content sources and attributions, please refer to our [editorial policy](#).



We comply with the HONcode standard for trustworthy health information - [verify here](#)

Copyright © 2000-2017 Drugs.com. All rights reserved.