

What's important to you in a treatment for moderate to severe Rheumatoid Arthritis (RA)?

[Home](#) > [Conditions](#) > [Pain](#) > [Norco](#)

[Print](#) [Share](#)

Norco



Generic Name: [acetaminophen and hydrocodone](#) (a SEET a MIN oh fen and hye droe KOE done)

Brand Names: *Hycet, Lorcet, Lortab 10/325, Lortab 5/325, Lortab 7.5/325, Lortab Elixir, Norco, Verdrocet, Vicodin, Xodol*

3 Fatigue-Causing Foods - 3 foods that cause fatigue.

Discover 3 foods that cause fatigue & 1 natural energy-boosting remedy! Go to [haveenergyatanyage.com](#)



[Overview](#)

[Side Effects](#)

[Dosage](#)

[Interactions](#)

[Patient Tips](#)

[Professional](#)

[More](#)

What is Norco?

Norco contains a combination of acetaminophen and hydrocodone. Hydrocodone is an opioid pain medication. An opioid is sometimes called a narcotic. Acetaminophen is a less potent pain reliever that increases the effects of hydrocodone.

Norco is used to relieve moderate to moderately severe [pain](#).

ADVERTISEMENT

Ads by

Norco may also be used for purposes not listed in this medication guide.

Important information

Hydrocodone can slow or stop your breathing. Never use Norco in larger amounts, or for longer than prescribed. Narcotic pain medicine may be habit-forming, even at regular doses. Never share Norco with another person, especially someone with a history of drug abuse or addiction. Keep the medication in a place where others cannot get to it.

MISUSE OF NARCOTIC MEDICINE CAN CAUSE ADDICTION, OVERDOSE, OR DEATH, especially in a child or other person using the medicine without a prescription.

Do not use Norco if you have used a MAO inhibitor in the past 14 days, such as isocarboxazid, linezolid, methylene blue injection, phenelzine, rasagiline, selegiline, or tranylcypromine.

An overdose of acetaminophen can damage your liver or cause death. Call your doctor at once if you have nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, or jaundice (yellowing of your skin or eyes).

Stop taking Norco and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling.

Before taking this medicine


You should not use Norco if you are allergic to acetaminophen (Tylenol) or hydrocodone, or if you have recently used alcohol, sedatives, tranquilizers, or other narcotic medications.

Do not use Norco if you have taken a MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, phenelzine, rasagiline, selegiline, and tranylcypromine.

Some medicines can interact with hydrocodone and cause a serious condition called serotonin syndrome. Be sure your doctor knows if you also take medicine for depression, mental illness, Parkinson's disease, migraine headaches, serious infections, or prevention of nausea and vomiting. Ask your doctor before making any changes in how or when you take your medications.

To make sure Norco is safe for you, tell your doctor if you have:

- liver disease, cirrhosis, or if you drink more than 3 alcoholic beverages per day;
- a history of alcoholism or drug addiction;
- diarrhea, inflammatory bowel disease, bowel obstruction, severe constipation;
- kidney disease;
- low blood pressure, or if you are dehydrated;



- a history of head injury, brain tumor, or stroke;
- asthma, COPD, sleep apnea, or other breathing disorders; or
- if you use a sedative like Valium (diazepam, alprazolam, lorazepam, Ativan, Klonopin, Restoril, Tranxene, Versed, Xanax, and others).

Norco is more likely to cause breathing problems in older adults and people who are severely ill, malnourished, or otherwise debilitated.

If you use narcotic medicine while you are pregnant, your baby could become dependent on the drug. This can cause life-threatening withdrawal symptoms in the baby after it is born. Babies born dependent on habit-forming medicine may need medical treatment for several weeks. Tell your doctor if you are pregnant or plan to become pregnant.

Acetaminophen and hydrocodone can pass into breast milk and may harm a nursing baby. You should not breast-feed while using Norco.



How should I take Norco?

Take Norco exactly as prescribed. Follow all directions on your prescription label. Never take this medicine in larger amounts, or for longer than prescribed. An overdose can damage your liver or cause death. Tell your doctor if the medicine seems to stop working as well in relieving your pain.

Hydrocodone may be habit-forming, even at regular doses. Never share this medicine with another person, especially someone with a history of drug abuse or addiction. MISUSE OF NARCOTIC MEDICINE CAN CAUSE ADDICTION, OVERDOSE, OR DEATH, especially in a child or other person using the medicine without a prescription. Selling or giving away Norco is against the law.

If you need surgery or medical tests, tell the doctor ahead of time that you are using this medicine. You may need to stop using the medicine for a short time.

Do not stop using this medicine suddenly after long-term use, or you could have unpleasant withdrawal symptoms. Ask your doctor how to safely stop using Norco.

Store Norco at room temperature away from moisture and heat. Keep track of the amount of medicine used from each new bottle. Hydrocodone is a drug of abuse and you should be aware if anyone is using your medicine improperly or without a prescription.

Always check your bottle to make sure you have received the correct pills (same brand and type) of medicine prescribed by your doctor.

Norco dosing information

Usual Adult Dose of Norco for Pain:

Tablets:

Recommended dose:

Acetaminophen-hydrocodone 300 mg-5 mg: 1 to 2 tablets orally every 4 to 6 hours as needed

Maximum dose:

Acetaminophen-hydrocodone 300 mg-5 mg: 8 tablets daily

Comments:

Dosage should be adjusted according to severity of the pain and the response of the patient. Tolerance to hydrocodone can develop with continued use and the incidence of side effects is dose related.

Approved indication: For the relief of moderate to moderately severe pain

See also: [Dosage Information \(in more detail\)](#)

What happens if I miss a dose?

Since Norco is taken as needed, you may not be on a dosing schedule. If you are taking the medication regularly, take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not use extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. An overdose of acetaminophen and hydrocodone can be fatal.

The first signs of an acetaminophen overdose include loss of appetite, nausea, vomiting, stomach pain, sweating, and confusion or weakness. Later symptoms may include pain in your upper stomach, dark urine, and yellowing of your skin or the whites of your eyes.

Overdose symptoms may also include extreme drowsiness, pinpoint pupils, cold and clammy skin, muscle weakness, fainting, weak pulse, slow heart rate, coma, blue lips, shallow breathing, or no breathing


What should I avoid while taking Norco?

This medication may impair your thinking or reactions. Avoid driving or operating machinery until you know how

Norco will affect you. Dizziness or severe drowsiness can cause falls or other accidents.

Ask a doctor or pharmacist before using any other cold, allergy, pain, or sleep medication. Acetaminophen (sometimes abbreviated as APAP) is contained in many combination medicines. Taking certain products together can cause you to get too much acetaminophen which can lead to a fatal overdose. Check the label to see if a medicine contains acetaminophen or APAP.

Avoid drinking alcohol. It may increase your risk of liver damage while taking acetaminophen.



Norco side effects

Get emergency medical help if you have signs of an allergic reaction to Norco: hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen.

Call your doctor at once if you have:

- shallow breathing, slow heartbeat;
- a light-headed feeling, like you might pass out;
- confusion, unusual thoughts or behavior;
- seizure (convulsions);
- easy bruising or bleeding;
- infertility, missed menstrual periods;
- impotence, sexual problems, loss of interest in sex;
- liver problems--nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes); or

- low cortisol levels-- nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness.

Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea.

Common Norco side effects include:

- drowsiness, headache;
- upset stomach, constipation;
- blurred vision; or
- dry mouth.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

See also: [Side effects \(in more detail\)](#)

What other drugs will affect Norco?

Narcotic (opioid) medication can interact with many other drugs and cause dangerous side effects or death. Be sure your doctor knows if you also use:

- other narcotic medications - opioid pain medicine or prescription cough medicine;
- drugs that make you sleepy or slow your breathing - a sleeping pill, muscle relaxer, sedative, tranquilizer, or antipsychotic medicine; or
- drugs that affect serotonin levels in your body - medicine for depression, Parkinson's disease, migraine headaches, serious infections, or prevention of nausea and vomiting.

This list is not complete. Other drugs may interact with acetaminophen and hydrocodone, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible interactions are listed in this medication guide.

Next → [Side Effects](#)

5 Prostate Cancer Signs

Dr. Brownstein Reveals the 5 Signals when Prostate Cancer is Coming. Go to w3.brownsteinhealth.com



ADVERTISEMENT



Add to My Med List

More about Norco (acetaminophen / hydrocodone)

- [Side Effects](#)
- [During Pregnancy](#)
- [Dosage Information](#)
- [Drug Images](#)
- [Drug Interactions](#)
- [Support Group](#)
- [Pricing & Coupons](#)
- [En Español](#)
- [210 Reviews – Add your own review/rating](#)
- [Drug class: narcotic analgesic combinations](#)

Consumer resources

- [Norco](#) (Advanced Reading)

Other brands: [Vicodin](#), [Lortab](#), [Lorcet](#), [Hycet](#), [More...](#) (9)

Professional resources

- [Norco](#) (FDA)

Related treatment guides

- [Back Pain](#)
- [Rheumatoid Arthritis](#)
- [Pain](#)

Where can I get more information?

- Your pharmacist can provide more information about Norco.
- Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Norco only for the indication prescribed.
- Disclaimer: Every effort has been made to ensure that the information provided is accurate, up-to-date, and complete, but no guarantee is made to that effect. Multum's drug information is an informational resource designed to assist licensed healthcare practitioners in caring for their patients and/or to serve consumers viewing this service as a supplement to, and not a substitute for, the expertise, skill, knowledge and judgment of healthcare practitioners. The information contained herein is not intended to cover all possible uses, directions, precautions, warnings, drug interactions, allergic reactions, or adverse effects. If you have questions about the drugs you are taking, check with your doctor, nurse or pharmacist.

Copyright 1996-2017 Cerner Multum, Inc. Version: 14.11. Revision Date: 2016-09-29, 5:18:57 PM.

DRUG STATUS



Availability
Prescription only



Pregnancy Category
Risk cannot be ruled out



CSA Schedule
High potential for abuse



Approval History
Drug history at FDA



Diagnosed with
ankylosing spondylitis?

Learn about biologics >

Manufacturer

[Watson Pharmaceuticals, Inc.](#)

Drug Class

[Narcotic analgesic combinations](#)

ADVERTISEMENT





Related Drugs

Back Pain

[tramadol](#), [naproxen](#), [Cymbalta](#), [aspirin](#), [ibuprofen](#), [diclofenac](#), [duloxetine](#), [Vicodin](#), [Voltaren](#), [Motrin](#), [Ultram](#), [More...](#)

Pain

[tramadol](#), [acetaminophen](#), [Tylenol](#), [oxycodone](#), [naproxen](#), [aspirin](#), [ibuprofen](#), [amitriptyline](#), [clonidine](#), [Percocet](#), [hydrocodone](#), [More...](#)

Norco Rating

[210 User Reviews](#)

8.0/10



Norco Images

Norco (acetaminophen / hydrocodone) 325 mg / 10 mg

[View all 7 images](#)

Related News and Articles

[4 in 10 People Will Suffer Arthritic Hands Over Lifetime](#)

May 4, 2017

[Babies Born Addicted to Opioids Often Struggle With Learning](#)

May 3, 2017

[Opioid Use by Iraq, Afghanistan War Vets Mirrors Rest of U.S.: Study](#)

April 27, 2017

Diagnosed with
ankylosing spondylitis?
Learn about biologics >



Harvard Health Publications

Related: [Back Pain](#)



MAYO CLINIC

Related: [Back pain](#)



Drugs.com Mobile Apps

The easiest way to lookup drug information, identify pills, check interactions and set up your own personal medication records. Available for Android and iOS devices.

Explore Apps

SUPPORT

[Help Center](#)[Frequent Questions](#)[Sitemap](#)[Contact Us](#)

ABOUT

[About Drugs.com](#)[Advertising Policy](#)[Content Submissions](#)[Drugs.com Blog](#)

TERMS & PRIVACY

[Editorial Policy](#)[Privacy Policy](#)[Terms of Use](#)[Attribution & Citations](#)

[Subscribe](#) to receive email notifications whenever new articles are published.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex® (updated May 1st, 2017), Cerner Multum™ (updated May 3rd, 2017), Wolters Kluwer™ (updated May 2nd, 2017) and others. To view content sources and attributions, please refer to our [editorial policy](#).



We comply with the HONcode standard for trustworthy health information - [verify here](#)

Copyright © 2000-2017 Drugs.com. All rights reserved.