



## Coronavirus Disease 2019 (COVID-19)

### People with Moderate to Severe Asthma

This information is based on what we currently know about the spread and severity of coronavirus disease 2019 (COVID-19).

#### Risk of Severe Illness from COVID-19

People with moderate to severe asthma may be at higher risk of getting very sick from COVID-19. COVID-19 can affect your respiratory tract (nose, throat, lungs), cause an asthma attack, and possibly lead to pneumonia and acute respiratory disease.

#### Treatment

There is currently no specific treatment for or vaccine to prevent COVID-19. The best way to prevent illness is to avoid being exposed to this virus.

#### Prepare for COVID-19

- [Stock up on supplies](#).
- Take [everyday precautions](#) to keep space between yourself and others.
- When you go out in public, keep away from others who are sick.
- Clean your hands often by washing with soap and water or using an alcohol-based hand sanitizer.
- [Avoid crowds](#) and people who are sick.
- Avoid [cruise travel](#) and non-essential air travel.
- During a COVID-19 outbreak in your community, [stay home](#) as much as possible to further reduce your risk of being exposed.
- If someone in your home is sick, have them stay away from the rest of the household to reduce the risk of spreading the virus in your home.
- Avoid sharing personal household items such as cups and towels.

#### Follow your Asthma Action Plan

- Keep your asthma under control by following [your asthma action plan](#).
- Continue your current medications, including any inhalers with steroids in them (“steroids” is another word for corticosteroids).
- Don’t stop any medications or change your asthma treatment plan without talking to your healthcare provider.
- Discuss any concerns about your treatment with your healthcare provider.
- Talk to your healthcare provider, insurer, and pharmacist about creating an emergency supply of prescription medications, such as asthma inhalers. Make sure that you have 30 days of non-prescription medications and [supplies](#) on hand too in case you need to stay home for a long time.

Visited on 06/16/2020

- [Know how to use your inhaler.](#)
- Avoid your [asthma triggers](#).
- As more cases of COVID-19 are discovered and our communities take action to combat the spread of disease, it is natural for some people to feel concerned or stressed. Strong emotions can trigger an asthma attack. [Take steps to help yourself cope with stress and anxiety.](#)

For information on administering asthma medication in schools, visit [K-12 Schools and Child Care Programs](#).

## Clean and disinfect things you or your family touch frequently

- If possible, have someone who doesn't have asthma do the cleaning and disinfecting. When they use cleaning and disinfecting products, have them:
  - Make sure that people with asthma are not in the room.
  - Minimize use of disinfectants that can cause an asthma attack.
  - Open windows or doors and use a fan that blows air outdoors.
  - [Clean and disinfect](#) surfaces like phones, remotes, tables, doorknobs, light switches, countertops, handles, desks, keyboards, toilets, faucets, and sinks daily.
  - Always follow the instructions on the product label.
  - Spray or pour spray products onto a cleaning cloth or paper towel instead of spraying the product directly onto the cleaning surface (if the product label allows).

## If you have symptoms

Contact your health care provider to ask about your symptoms.

Page last reviewed: April 2, 2020

Content source: [National Center for Immunization and Respiratory Diseases \(NCIRD\), Division of Viral Diseases](#)