

Visited 04/21/2015



Synthroid

Generic Name: levothyroxine (LEE voe thye ROX een)

Brand Names: *Levothroid, Levoxyl, Synthroid, Tirosint, Unithroid*

What is Synthroid?

Synthroid (levothyroxine) is a replacement for a hormone normally produced by your thyroid gland to regulate the body's energy and metabolism. Levothyroxine is given when the thyroid does not produce enough of this hormone on its own.

Synthroid treats hypothyroidism (low thyroid hormone). Synthroid is also used to treat or prevent goiter (enlarged thyroid gland), which can be caused by hormone imbalances, radiation treatment, surgery, or cancer.

Synthroid may also be used for purposes not listed in this medication guide.

Important information

You may not be able to take Synthroid if you have certain medical conditions. Tell your doctor if you have an untreated or uncontrolled adrenal gland disorder, a thyroid disorder called thyrotoxicosis, or if you have any recent or current symptoms of a heart attack.

Synthroid should not be used to treat obesity or weight problems. Dangerous side effects or death can occur from the misuse of Synthroid, especially if you are taking any other weight-loss medications or appetite suppressants.

Before taking this medicine

Synthroid should not be used to treat obesity or weight problems. Dangerous side effects or death can occur from the misuse of levothyroxine, especially if you are taking any other weight-loss medications or appetite suppressants.

Since thyroid hormone occurs naturally in the body, almost anyone can take Synthroid. However, you may not be able to take this medication if you have certain medical conditions.

To make sure Synthroid is safe for you, tell your doctor if you have:

- a thyroid disorder called thyrotoxicosis;
- heart disease, coronary artery disease, or a history of blood clots;
- diabetes (insulin or oral [diabetes medication](#)  doses may need to be changed when you start taking levothyroxine);
- anemia (lack of red blood cells);
- osteoporosis, or low bone mineral density;
- problems with your pituitary gland;

Visited 04/21/2015

- any food or drug [allergies](#) 
- an untreated or uncontrolled adrenal gland disorder; or
- if you have recently had a heart attack, or are having any symptoms of a heart attack (chest pain or heavy feeling, pain spreading to the jaw or shoulder, nausea, sweating, general ill feeling).

Tell your doctor if you have recently received radiation therapy with iodine (such as I-131).

FDA pregnancy category A. Synthroid is not expected to harm an unborn [baby](#) . If you become pregnant while taking levothyroxine, do not stop taking the medicine without your doctor's advice. Having low thyroid hormone levels [during pregnancy](#)  could harm both mother and baby. Your dose needs may be different during pregnancy.

See also: Pregnancy and breastfeeding warnings (in more detail)

Levothyroxine can pass into breast milk, but it is not expected to be harmful to a nursing baby. Do not use this medication without telling your doctor if you are breast-feeding a baby. Your dose needs may be different while you are nursing.

How should I take Synthroid?

Take Synthroid exactly as prescribed by your doctor. Follow all directions on your prescription label. Your doctor may occasionally change your dose to make sure you get the best results. Do not take this medicine in larger or smaller amounts or for longer than recommended.

Do not share this [medication](#)  with another person, even if they have the same symptoms you have.

Synthroid works best if you take it on an empty stomach, at least 30 minutes before breakfast. Follow your doctor's dosing instructions and try to take the medicine at the same time each day.

It is very important to take Synthroid with a full glass (8 ounces) of water. The levothyroxine tablet can dissolve very quickly and swell in the throat, possibly causing choking or gagging.

While using Synthroid, you may need frequent medical tests.

Tell any doctor or dentist who treats you that you are using Synthroid.

Store at room temperature away from moisture and heat.

It may take several weeks before your body starts to respond to Synthroid. Keep using this medicine even if you feel well. You may need to use this medicine for the rest of your life to replace the thyroid hormone your body cannot produce.

What happens if I miss a dose?

Take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not take extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include headache, leg cramps, tremors, feeling nervous or irritable, chest pain, shortness of breath, and fast or pounding heartbeats.

Visited 04/21/2015

What should I avoid?

Certain medicines can make Synthroid less effective if taken at the same time. If you use any of the following drugs, avoid taking them within 4 hours before or 4 hours after you take Synthroid:

- calcium carbonate (Alka-Mints, Calcium Oyster Shell, Caltrate, Os-Cal, Oyster Shell Calcium, Roloids Soft Chew, Tums, and others);
- cholestyramine, colestipol;
- ferrous sulfate iron supplement;
- sucralfate;
- sodium polystyrene sulfonate (Kalexate, Kayexalate, Kionex); or
- antacids that contain aluminum or magnesium - Acid Gone, Gaviscon, Maalox, Milk of Magnesia, Mintox, Mylanta, Pepcid Complete, and others).

Avoid the following food products, which can make your body absorb less Synthroid: infant soy formula, cotton seed meal, walnuts, and high-fiber foods.

Synthroid side effects

Get emergency medical help if you have any of these signs of an allergic reaction to Synthroid: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Call your doctor at once if you have a serious side effect such as:

- fast or irregular heart rate;
- fever, hot flashes, sweating;
- sleep problems (insomnia);
- changes in your menstrual periods; or
- vomiting, diarrhea, appetite changes, weight changes.

Common Synthroid side effects may include mild hair loss.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

See also: Side effects (in more detail)

What other drugs will affect Synthroid?

Many other medicines can be affected by your thyroid hormone levels. Other medicine may also increase or decrease the effects of Synthroid.

Many drugs can interact with levothyroxine and not all possible interactions are listed in this medication guide.

Tell your doctor about all medications you use, start using, or stop using during your treatment with Synthroid. This includes prescription, over-the-counter, vitamin, and herbal products. Give a list of all your medicines to any healthcare provider who treats you.

Where can I get more information?

Visited 04/21/2015

- Your pharmacist can provide more information about Synthroid.

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Synthroid only for the indication prescribed.

Disclaimer: Every effort has been made to ensure that the information provided by Cerner Multum, Inc. ('Multum') is accurate, up-to-date, and complete, but no guarantee is made to that effect. Drug information contained herein may be time sensitive. Multum information has been compiled for use by healthcare practitioners and consumers in the United States and therefore Multum does not warrant that uses outside of the United States are appropriate, unless specifically indicated otherwise. Multum's drug information does not endorse drugs, diagnose patients or recommend therapy. Multum's drug information is an informational resource designed to assist licensed healthcare practitioners in caring for their patients and/or to serve consumers viewing this service as a supplement to, and not a substitute for, the expertise, skill, knowledge and judgment of healthcare practitioners. The absence of a warning for a given drug or drug combination in no way should be construed to indicate that the drug or drug combination is safe, effective or appropriate for any given patient. Multum does not assume any responsibility for any aspect of healthcare administered with the aid of information Multum provides. The information contained herein is not intended to cover all possible uses, directions, precautions, warnings, drug interactions, allergic reactions, or adverse effects. If you have questions about the drugs you are taking, check with your doctor, nurse or pharmacist.

Copyright 1996-2015 Cerner Multum, Inc. Version: 11.03. Revision Date: 2014-09-14, 6:08:51 PM.