
Asthma & COPD Treatment | SPIRIVA® (tiotropium bromide)

Visited 05/05/2020



INDICATIONS for SPIRIVA RESPIMAT (tiotropium bromide) Inhalation Spray and SPIRIVA HANDIHALER (tiotropium bromide inhalation powder)

SPIRIVA RESPIMAT, 2.5 mcg, and SPIRIVA HANDIHALER are long-term, once-daily, prescription maintenance medicines used to control symptoms of chronic obstructive pulmonary disease (COPD) by relaxing your airways and keeping them open. COPD includes chronic bronchitis and emphysema. SPIRIVA RESPIMAT and SPIRIVA HANDIHALER also reduce the likelihood of flare-ups (COPD exacerbations).

SPIRIVA RESPIMAT, 1.25 mcg, is a long-term, once-daily, prescription maintenance treatment of asthma for people 6 years and older.

SPIRIVA is not a treatment for sudden symptoms of asthma or COPD.

r any ingradiant in these medicines

IMPORTANT SAFETY INFORMATION for SPIRIVA RESPIMAT (tiotropium bromide) Inhalation Spray and SPIRIVA HANDIHALER (tiotropium bromide inhalation powder)

Do not use SPIRIVA RESPIMAT or SPIRIVA HANDIHALER if you are allergic to tiotropium, ipratropium, atropine or similar drugs,

IMPORTANT SAFETY INFORMATION for SPIRIVA RESPIMAT (tiotropium bromide) and SPIRIVA HANDIHALER (tiotropium bromide inhalation powder)

Do not use SPIRIVA RESPIMAT or SPIRIVA HANDIHALER if you are allergic to tiotropium, ipratropium, atropine or similar drugs, or any ingredient in these medicines.

SPIRIVA RESPIMAT or SPIRIVA HANDIHALER are not rescue medicines and should not be used for treating sudden breathing problems. Your doctor may give you other medicine to use for sudden breathing problems.

Visited 05/05/2020

SPIRIVA RESPIMAT or SPIRIVA HANDIHALER can cause your breathing to suddenly get worse (bronchospasm). If this happens, use your rescue inhaler, stop taking SPIRIVA, and call your doctor right away or seek emergency medical care.

SPIRIVA RESPIMAT or SPIRIVA HANDIHALER can increase the pressure in your eyes (acute narrow-angle glaucoma) which can cause the following symptoms: eye pain, blurred vision, seeing halos or colored images along with red eyes. If you have any of these symptoms, stop taking your medicine and call your doctor right away.

Dizziness and blurred vision may occur with SPIRIVA RESPIMAT or SPIRIVA HANDIHALER. If you experience these symptoms, use caution when engaging in activities such as driving a car, or operating appliances or machinery.

SPIRIVA RESPIMAT or SPIRIVA HANDIHALER can cause new or worsened urinary retention. Symptoms of blockage in your bladder and/or enlarged prostate may include difficulty passing urine and/or painful urination. If you have any of these symptoms, stop taking your medicine and call your doctor right away.

The most common side effects reported with SPIRIVA RESPIMAT in patients with COPD include sore throat, cough, dry mouth, and sinus infection.

The most common side effects with SPIRIVA RESPIMAT in adult patients with asthma were sore throat, headache, bronchitis, and sinus infection. The side effect profile for adolescent and pediatric patients was comparable to that observed in adult patients with asthma.

The most common side effects reported with SPIRIVA HANDIHALER in patients with COPD include upper respiratory tract infection, dry mouth, sinus infection, sore throat, non-specific chest pain, urinary tract infection, indigestion, runny nose, constipation, increased heart rate, and blurred vision.

Do not swallow SPIRIVA capsules. The contents of the capsule should only be inhaled through your mouth using the HANDIHALER device.

Do not spray SPIRIVA RESPIMAT into your eyes, as this may cause blurring of vision and pupil dilation.

Tell your doctor about all your medical conditions including kidney problems, glaucoma, enlarged prostate, problems passing urine, or blockage in your bladder.

Tell your doctor all the medicines you take, including eye drops. Ask your doctor if you are taking any anticholinergic medicines because taking them together with SPIRIVA can increase side effects. Do not use SPIRIVA RESPIMAT and SPIRIVA HANDIHALER together.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit <u>www.FDA.gov/medwatch</u> or call 1-800-FDA-1088.

Please see full <u>Prescribing Information</u> for SPIRIVA RESPIMAT, including <u>Instructions for Use</u>, and full <u>Prescribing Information</u> for SPIRIVA HANDIHALER, including <u>Patient Information</u> and <u>Instructions for Use</u>.

CL-SVR-0048 2.15.2017

IMPORTANT SAFETY INFORMATION for SPIRIVA RESPIMAT (tiotropium bromide) and SPIRIVA HANDIHALER (tiotropium bromide inhalation powder)

Do not use SPIRIVA RESPIMAT or SPIRIVA HANDIHALER if you are allergic to tiotropium, ipratropium, atropine or similar drugs, or any ingredient in these medicines.

SPIRIVA RESPIMAT or SPIRIVA HANDIHALER are not rescue medicines and should not be used for treating sudden breathing problems. Your doctor may give you other medicine to use for sudden breathing problems.

Visited 05/05/2020

This information is for U.S. residents only. Products discussed herein may have different labeling in different countries. Use of this site is subject to the Internet Site Legal Notices and Disclaimers and Privacy Notice.

Copyright © 2019 Boehringer Ingelheim Pharmaceuticals, Inc. All rights reserved. (11/19)

Partnership for Prescription Assistance

If you or someone you know needs help paying for medicine, click here.

IMPORTANT SAFETY INFORMATION for SPIRIVA RESPIMAT (tiotropium bromide) and SPIRIVA HANDIHALER (tiotropium bromide inhalation powder)

Do not use SPIRIVA RESPIMAT or SPIRIVA HANDIHALER if you are allergic to tiotropium, ipratropium, atropine or similar drugs, or any ingredient in these medicines.

SPIRIVA RESPIMAT or SPIRIVA HANDIHALER are not rescue medicines and should not be used for treating sudden breathing problems. Your doctor may give you other medicine to use for sudden breathing problems.