
MAYO CLINIC UPD						Request an AppointmentLog in to PatientFind a DoctorAccountFind a JobTranslated ContentMake a DonationTranslated Content	
Search Mayo Clinic				Submit Query			
PATIENT CARE & HEALTH INFO DEPARTMENTS & CENTERS PRODUCTS & SERVICES GIVING TO MAYO CLINIC		ITERS	RESEARCH	EDUCATION	FOR MEDICAL PRO	FESSIONALS	
Appointments at Mayo Clinic Mayo Clinic offers appointments in Arizona, Florida and Minnesota and at Request Appointment Mayo Clinic Health System locations. Request Appointment							

Ejection fraction: What does it measure?

Products and services



Free E-newsletter Subseribe to Housecell Our weekly general interest e-newsletter keeps you up to date on a wide variety of health topics.

Sign up now

What does the term "ejection fraction" mean? What does it measure?

Answers from Martha Grogan, M.D.

Ejection fraction is a measurement of the percentage of blood leaving your heart each time it contracts.

During each heartbeat cycle, the heart contracts and relaxes. When your heart contracts, it ejects blood from the two pumping chambers (ventricles). When your heart relaxes, the ventricles refill with blood. No matter how forceful the contraction, it doesn't empty all of the blood out of a ventricle. The term "ejection fraction" refers to the percentage of blood that's pumped out of a filled ventricle with each heartbeat.


The left ventricle is the heart's main pumping chamber, so ejection fraction is usually measured only in the left ventricle (LV). An LV ejection fraction of 55 percent or higher is considered normal. An LV ejection fraction of 50 percent or lower is considered reduced. Experts vary in their opinion about an ejection fraction between 50 and 55 percent, and some would consider this a "borderline" range.

Keep in mind that ejection fraction is just one measure of heart function. Even with a normal ejection fraction, overall heart function may not be normal. Talk with your doctor if you have concerns about your heart.

The ejection fraction may decrease if:

- You have weakness of your heart muscle, such as dilated cardiomyopathy
- · A heart attack has damaged your heart
- · You have problems with your heart's valves
- · You have had long-standing, uncontrolled high blood pressure

Ejection fraction can be measured with imaging techniques, including:



Advertising and Policy

Mayo Clinic is a not-for-profit organization. Proceeds from website advertising help support our mission. Mayo Clinic does not endorse non-Mayo products and services.

Advertising & Sponsorship

Policy Opportunities

Print

- · Cardiac catheterization. During cardiac catheterization, a thin, plastic tube (catheter) is inserted into a vein in the arm or leg and then moved to the heart - most likely the left ventricle. Using images taken during the catheterization, your doctor can see how your blood pumps through your heart.
- Magnetic resonance imaging (MRI). During an MRI scan, a magnetic field and radio waves are used to create crosssectional images of specific parts of the body. When an MRI is used to study the heart, it's known as a cardiovascular MRI.
- Computerized tomography (CT). During a CT scan, a special X-ray technique is used to create cross-sectional images of specific parts of the body. When a CT scan is used to study the heart, it's known as a cardiac CT.
- Nuclear medicine scan. During a nuclear scan, trace amounts of radioactive material - such as thallium - are injected into the bloodstream. Special cameras then detect the radioactive material in your blood as it flows through the heart and lungs.

With

Martha Grogan, M.D.

Share

Feb. 20, 2013

References

1. Bonow RO, et al. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 9th ed. Philadelphia, Pa.: Saunders Elsevier; 2012. http://www.mdconsult.com/books/about.do?eid=4-u1.0-B978-1-4377-0398-6..C2009-0-59734-6--TOP&isbn=978-1-4377-0398-6&about=true&uniqId=236798031-10. Accessed Jan. 18, 2013.

Tweet

- 2. Ejection fraction heart failure measurement. American Heart Association. http://www.heart.org/HEARTORG/Conditions/HeartFailure/SymptomsDiagnosisofHe33151ipatientovears of life -Fraction-Heart-Failure-Measurement_UCM_306339_Article.jsp. Accessed Jan. 18, 2013.
- 3. Common tests for heart failure. American Heart Association. http://www.heart.org/HEARTORG/Conditions/HeartFailure/SymptomsDiagnosisofHeartFailure/Common -Tests-for-Heart-Failure_UCM_306334_Article.jsp. Accessed Jan. 30, 2013.
- 4. Grogan M (expert opinion). Mayo Clinic, Rochester, Minn. Feb. 4, 2013.

See more Expert Answers

See also

Sudden cardiac arrest Sudden death in young people Holiday Heart Heart failure Cardiac asthma: What causes it?

Products and services

Mayo Clinic Store

Check out these best-sellers and special offers on books and newsletters from Mayo Clinic

Try Mayo Clinic Health Letter FREE!

Improve heart health in just 10 minutes a dav

Get a better night's sleep with this threestep action plan NEW! Mayo Clinic on Better Hearing and Balance

The Mayo Clinic Diet - Eat well, Enjoy life. Lose weight.

Ads by Google

Ask The ICD

www.asktheicd.com

Get Answers To Common Defibrillator Questions. Learn More Here!

Mouse/Rat Blood Pressure

kentscientific.com/BloodPressure

Validated, Non-invasive, Tail-cuff. 99% Correlation With Telemetry.

Senior Balance Exercises


thebalancemanual.com Seniors Can Improve Balance And Prevent Falls With Simple Exercises

Total Artificial Heart

www.syncardia.com

Over 1,200 implants account for more than

http://www.mayoclinic.org/ejection-fraction/expert-answers/faq-20058286



 $\ensuremath{\textcircled{\sc 0}}$ 1998-2014 Mayo Foundation for Medical Education and Research. All rights reserved.

Captured 2/24/2014