

Severe Allergies:
What You Need to Know

[watch video](#)



Browse all medications [A](#) [B](#) [C](#) [D](#) [E](#) [F](#) [G](#) [H](#) [I](#) [J](#) [K](#) [L](#) [M](#) [N](#) [O](#) [P](#) [Q](#) [R](#) [S](#) [T](#) [U](#) [V](#) [W](#) [X](#) [Y](#) [Z](#) [Advanced Search](#) [Phonetic Search](#)



Drugs A-Z ▾

Pill Identifier

Interactions Checker

News ▾

Health Professionals ▾

Q & A ▾

Mednotes

Apps

Home → Conditions → Muscle Spasm → Cyclobenzaprine

Print Share

[Sign In](#) or [Register](#)

Cyclobenzaprine

Generic Name: cyclobenzaprine (sye kloee BEN za preen)

Brand Names: *Amrix, Comfort Pac with Cyclobenzaprine, Fexmid, Flexeril*

Overview

Side Effects

Dosage

Interactions

For Professionals

More ▾

Bulging Disc Problems

Diagnosis & Treatment of Spine Disorders. Minimally Invasive



What is cyclobenzaprine?

Cyclobenzaprine is a muscle relaxant. It works by blocking nerve impulses (or pain sensations) that are sent to your brain.

Cyclobenzaprine is used together with rest and physical therapy to treat skeletal muscle conditions such as pain or [injury](#).

Cyclobenzaprine may also be used for purposes not listed in this medication [guide](#).

Important information

You should not use cyclobenzaprine if you are allergic to it, or if you have a thyroid disorder, heart block, congestive heart failure, a heart rhythm disorder, or you have recently had a heart attack.

Do not use cyclobenzaprine if you have taken an MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, phenelzine, rasagiline, selegiline, and tranylcypromine.

Slideshow: 2014 Update - First Time Brand-to-Generic Switches



Before taking this medicine

Do not use cyclobenzaprine if you have taken an MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, phenelzine, rasagiline, selegiline, and tranylcypromine.

Related Information



Availability
Prescription only



Pregnancy Category
No proven risk in humans



CSA Schedule
Not a controlled drug



Approval History
Drug history at FDA

SEVERE ALLERGIES?
LEARN ABOUT EMERGENCY
EPINEPHRINE.

[Watch >](#)

Reviews

[226 User Reviews](#)

Average User Rating

6.9 [Rate it!](#)

Drug Class

[Skeletal muscle relaxants](#)

Related Drugs

Muscle Spasm

[Soma](#)
[Flexeril](#)
[baclofen](#)
[diazepam](#)
[Valium](#)
[tizanidine](#)
[methocarbamol](#)
[Robaxin](#)
[Zanaflex](#)
[carisoprodol](#)
[metaxalone](#)
[Skelaxin](#)
[More...](#)

Sciatica

[naproxen](#)
[acetaminophen](#)
[Tylenol](#)
[aspirin](#)
[ibuprofen](#)
[Flexeril](#)
[diclofenac](#)
[Voltaren](#)
[indomethacin](#)
[Paracetamol](#)
[Aleve](#)

ADVERTISEMENT

ADVERTISEMENT

You should not use cyclobenzaprine if you are allergic to it, or if you have:

- a heart rhythm disorder, or you have recently had a heart attack;
- congestive heart failure;
- heart block; or
- a thyroid disorder.

To make sure cyclobenzaprine is safe for you, tell your doctor if you have:

- liver disease;
- glaucoma;
- enlarged prostate; or
- problems with urination.

Older adults may be more sensitive to the side effects of this medication.

FDA pregnancy category B. Cyclobenzaprine is not expected to harm an unborn [baby](#). Tell your doctor if you are [pregnant](#) or plan to become pregnant during treatment.

See also: [Pregnancy and breastfeeding warnings](#) (in more detail)

It is not known whether cyclobenzaprine passes into breast milk or if it could harm a [nursing](#) baby. Tell your doctor if you are breast-feeding a baby.

Rebif Official Site



[rebif.com](#)

Is Rebif® (interferon beta-1a) Right For You?
Come Learn More Now!

Sciatica Exercises?

Stop Leg Cramps in 1 Min.

How should I take cyclobenzaprine?

Cyclobenzaprine is usually taken once daily for only 2 or 3 weeks. Follow all directions on your prescription label. Do not take this medicine in larger or smaller amounts or for longer than recommended.

Take the medicine at the same time each day.

Do not crush, chew, break, or open an extended-release capsule. Swallow it whole.

You may have unpleasant withdrawal symptoms when you stop taking cyclobenzaprine after long-term use. Ask your doctor how to avoid withdrawal symptoms when you stop using this medicine.

Cyclobenzaprine is only part of a complete program of treatment that may also include rest, physical therapy, or other pain relief [measures](#). Follow your doctor's instructions.

Store at room temperature away from moisture, heat, and light.

What happens if I miss a dose?

Take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not take extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. An overdose of cyclobenzaprine can be fatal.

[Motrin](#)

[More...](#)

Cyclobenzaprine Images



Cyclobenzaprine 10 mg



Cyclobenzaprine 10 mg

[View all 65 images](#)

Get Updates by Email

Email me:

News and Warnings related to this drug

FDA Medwatch Alerts for all Medications

News Roundup Daily

Related News and Articles

[Avanir Pharmaceuticals Announces Preliminary Feedback from the FDA on AVP-825 for the Acute Treatment of Migraine](#) November 7, 2014

Avanir Pharmaceuticals, Inc. (NASDAQ: AVNR) today announced that the U.S. Food and Drug Administration (FDA) has issued preliminary written feedback to its New...

[Avanir Pharmaceuticals Announces Publication of Pivotal Phase III Results from AVP-825 Acute Migraine Study](#) October 30, 2014

Avanir Pharmaceuticals, Inc. (NASDAQ: AVNR) today announced the publication of results from TARGET, a pivotal phase III study evaluating the efficacy and...

[Medication Errors Occur Every 8 Minutes in U.S. Children](#) October 20, 2014

A child receives the wrong medication or the wrong dosage every eight minutes in the United States, according to a recent study. Nearly 700,000 children under...

GET THE FACTS ABOUT EPILEPSY

[Watch the video](#) >

What should I avoid?

This medication may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be [alert](#).

Drinking alcohol can increase certain side effects of cyclobenzaprine.

Antiplatelet Treatment

Find Information On Reducing
The Risk Of Thrombotic CV
Events.



Cyclobenzaprine side effects

Get emergency medical help if you have any of these signs of an allergic reaction to cyclobenzaprine: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Stop using this medicine and call your doctor at once if you have:

- severe drowsiness, fast heart rate;
- tremors or shaking;
- pounding heartbeats or fluttering in your chest; or
- agitation, hallucinations, [fever](#), overactive reflexes, nausea, vomiting, diarrhea, loss of coordination, fainting.

Common cyclobenzaprine side effects may include:

- headache, dizziness;
- drowsiness, tired feeling;
- trouble concentrating;
- blurred vision, dry mouth or throat, altered sense of taste; or
- nausea, upset stomach, constipation.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

See also: [Side effects](#) (in more detail)

Cyclobenzaprine dosing information

Usual Adult Dose for Muscle Spasm:

Initial dose: 5 mg orally 3 times a day.

Based on individual patient response, the dose may be increased to either 7.5 mg or 10 mg 3 times a day.

Alternate doses include the 15 mg extended release capsule or 30 mg extended release capsule orally once a day.

Usual Geriatric Dose for Muscle Spasm:

Because most muscle relaxants and antispasmodic drugs can cause anticholinergic adverse [events](#), sedation, and weakness, and because their effectiveness at doses tolerated by [elderly](#) people is questionable, cyclobenzaprine meets the Beers criteria as a medication that is potentially inappropriate for use in older adults.

The manufacturer suggests an initial dose in the elderly of 5 mg, titrating slowly upward based on individual patient response.

Extended release formulations are not recommended for use in elderly patients.

Usual Pediatric Dose for Muscle Spasm:

Pediatric patients 15 years of age and older:






Initial dose: 5 mg orally 3 times a day.

Based on individual patient response, the dose may be increased to either 7.5 mg or 10 mg 3 times a day.

What other drugs will affect cyclobenzaprine?

Taking cyclobenzaprine with other drugs that make you sleepy or slow your breathing can cause dangerous or life-threatening side effects. Ask your doctor before taking cyclobenzaprine with a [sleeping](#) pill, narcotic pain medicine, muscle relaxer, or medicine for anxiety, depression, or seizures.

This list is not complete. Other drugs may interact with cyclobenzaprine, including prescription and over-the-counter medicines, [vitamins](#), and herbal products. Not all possible interactions are listed in this medication guide.

 [Google](#)  [Facebook](#)  [Twitter](#)  [Print](#)  [Email](#)  [Add to Mednotes](#)

Next Page → [Side Effects](#)

More about cyclobenzaprine

- [Side Effects](#)
- [During Pregnancy](#) or [Breastfeeding](#)
- [Dosage Information](#)
- [Drug Images](#)
- [Drug Interactions](#)
- [Support Group](#)
- [Pricing & Coupons](#)
- [En Espanol](#)
- [226 Reviews](#) - [Add your own review/rating](#)

Consumer resources

- [Cyclobenzaprine](#)
- [Cyclobenzaprine extended-release capsules](#)
- [Cyclobenzaprine](#) (Advanced Reading)
- *Other brands:* [Flexeril](#), [Amrix](#), [Fexmid](#)

Professional resources

- [Cyclobenzaprine](#) (AHFS Monograph)
- [More \(3\) »](#)

Related treatment guides

- [Muscle Spasm](#)
- [Sciatica](#)
- [Fibromyalgia](#)
- [Chronic Myofascial Pain](#)
- [Temporomandibular Joint Disorder](#)
- [More \(1\) »](#)

Where can I get more information?

- Your pharmacist can provide more information about cyclobenzaprine.

1 Trick to Fibromyalgia

See How to Relieve Fibromyalgia With This Fast and Easy

Trick



Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medicine only for the indication prescribed.

Disclaimer: Every effort has been made to ensure that the information provided by Cerner Multum, Inc. ('Multum') is accurate, up-to-date, and complete, but no guarantee is made to that effect. Drug information contained herein may be time sensitive. Multum information has been compiled for use by healthcare practitioners and consumers in the United States and therefore Multum does not warrant that uses outside of the United States are appropriate, unless specifically indicated otherwise. Multum's drug information does not endorse drugs, diagnose patients or recommend therapy. Multum's drug information is an informational resource designed to assist licensed healthcare practitioners in caring for their patients and/or to serve consumers viewing this service as a supplement to, and not a substitute for, the expertise, skill, knowledge and judgment of healthcare practitioners. The absence of a warning for a given drug or drug combination in no way should be construed to indicate that the drug or drug combination is safe, effective or appropriate for any given patient. Multum does not assume any responsibility for any aspect of healthcare administered with the aid of information Multum provides. The information contained herein is not intended to cover all possible uses, directions, precautions, warnings, drug interactions, allergic reactions, or adverse effects. If you have questions about the drugs you are taking, check with your doctor, nurse or pharmacist.

Copyright 1996-2014 Cerner Multum, Inc. Version: 4.01. Revision Date: 2013-06-16, 11:37:41 PM.



Support

- [Advertise With Us](#)
- [Frequently Asked Questions](#)
- [Ask a Question](#)
- [Content Submissions](#)
- [Mobile Apps](#)
- [About Drugs.com](#)
- [Contact Us](#)
- [Sitemap](#)

Follow:

About Drugs.com

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment.

Data sources include Micromedex® (updated Oct 30th, 2014), Cerner Multum™ (updated Nov 17th, 2014), Wolters Kluwer™ (updated Nov 3rd, 2014) and others. To view content sources and attributions, refer to our [editorial policy](#).

Copyright © 2000-2014 Drugs.com. All rights reserved.

Your Privacy



We comply with the HONcode standard for trustworthy health information.

[Verify here](#)



Ad Choices

[Terms of Use](#) · [Privacy Policy](#)