



Drugs A-Z ▾

Pill Identifier

Interactions Checker

News ▾

Health Professionals ▾

Q & A ▾

Mednotes

Apps

Home → Drugs A to Z → Lidoderm patch → MedFacts

Print
 Share

[Sign In](#) or [Register](#)

Lidoderm patch

Generic Name: [lidocaine](#) (LYE-doe-kane)

Brand Name: Lidoderm

 Overview **Side Effects** Dosage Interactions More ▾

2015 Best Skin Tighteners

Products To Tighten Loose, Sagging Face Skin, Without Plastic Surgery



Lidoderm patch is used for:

Relieving pain associated with herpes zoster (shingles).

Lidoderm patch is a local anesthetic. It works by stopping nerves from transmitting painful impulses to the brain.

Do NOT use Lidoderm patch if:

- you are allergic to any ingredient in Lidoderm patch or other similar medicines (eg, amide-type medicines)


Contact your doctor or health care provider right away if any of these apply to you.

Before using Lidoderm patch:

Some medical conditions may interact with Lidoderm patch. Tell your doctor or pharmacist if you have any medical conditions, especially if any of the following apply to you:

- if you are pregnant, planning to become pregnant, or are breast-feeding
- if you are taking any prescription or nonprescription medicine, herbal preparation, or dietary supplement
- if you have allergies to medicines, foods, or other substances
- if you have broken, damaged, or inflamed skin at the application site
- if you have heart, liver, or kidney problems

Slideshow: Fighting the Fight: Fibromyalgia Explained



Related Information


Availability
 Rx and/or OTC

Pregnancy Category
 No proven risk in humans

CSA Schedule
 Not a controlled drug

Approval History
 Drug history at FDA

Harvard Health Publications
Related: [Pain](#)

Reviews

22 User Reviews

Average User Rating

7.9 [Rate it!](#)

Drug Class

[Topical anesthetics](#)

Related Drugs

Pain

[tramadol](#), [oxycodone](#), [naproxen](#), [acetaminophen](#), [Tylenol](#), [More...](#)

Hemorrhoids

[lidocaine topical](#), [hydrocortisone topical](#), [phenylephrine topical](#), [Anusol](#), [Anusol-HC](#), [benzocaine topical](#), [More...](#)

Anal Itching

[lidocaine topical](#), [hydrocortisone topical](#), [phenylephrine topical](#), [Anusol-HC](#), [benzocaine topical](#), [Anucort-HC](#), [More...](#)

Anesthesia

[fentanyl](#), [lidocaine topical](#), [hyoscyamine](#), [ketamine](#), [propofol](#), [More...](#)
[4 more conditions...](#)

ADVERTISEMENT

ADVERTISEMENT

- if you have had a severe allergic reaction (eg, severe rash, hives, difficulty breathing, dizziness) to any anesthetic medicine (eg, benzocaine)
- if you have very poor health
- if you are using another medicine that contains an anesthetic

Some MEDICINES MAY INTERACT with Lidoderm patch. Tell your health care provider if you are taking any other medicines, especially any of the following:

- Beta-blockers (eg, propranolol) or cimetidine because they may increase the risk of Lidoderm patch's side effects
- Antiarrhythmics (eg, amiodarone, mexiletine, tocainide) or local anesthetics (eg, benzocaine) because the risk of their side effects may be increased by Lidoderm patch

This may not be a complete list of all interactions that may occur. Ask your health care provider if Lidoderm patch may interact with other medicines that you take. Check with your health care provider before you start, stop, or change the dose of any medicine.

Reduce Nerve Pain Now



neuropathytreatmentgroup.com

Don't Let Neuropathy Symptoms Slow You Down. Get A Free Trial Now!

If you're 50+ read this



Bulging Disc Pain
Relief



How to use Lidoderm patch:

Use Lidoderm patch as directed by your doctor. Check the label on the medicine for exact dosing instructions.

- Lidoderm patch is only to be used on intact, clean, dry skin. Do not apply Lidoderm patch to broken, damaged, or inflamed skin.
- Wash your hands before and immediately after using Lidoderm patch.
- Remove the liner on the patch. Apply the patch so that it covers the painful area. If the painful area is larger than the patch, apply the patch over the most painful area. Patches may also be cut into smaller sizes before removing the liner if desired. Be sure to discard the unused pieces of cut patches out of the reach of children and away from pets.
- If irritation or burning occurs when you apply the patch, remove it. Do not reapply any patches until the irritation stops.
- Patches may be worn for up to 12 hours within a 24-hour period as directed by your doctor. Clothing may be worn over the area where the patch is applied.
- After removing a patch, fold it in half with the sticky sides together. Discard the patch out of reach of children and away from pets. Do not apply another patch to the same area of skin for 12 hours.
- If you miss a dose of Lidoderm patch, use the dose when you remember. Continue to use it as directed by your doctor. Do not use Lidoderm patch for more than 12 hours in a 24-hour period. Do not use 2 doses sooner than 12 hours apart.

Ask your health care provider any questions you may have about how to use Lidoderm patch.

Important safety information:

- Lidoderm patch is for external use only. Avoid contact with your eyes, nose, and mouth. If you get Lidoderm patch in your eyes, rinse immediately with cool water.
- Lidoderm patch may cause drowsiness, dizziness, or blurred vision. These effects may be worse if you take it with alcohol or certain medicines. Use Lidoderm patch with caution. Do not drive or perform other possibly unsafe tasks until you know how you react to it.
- Tell your doctor or dentist that you take Lidoderm patch before you receive any medical or dental care, emergency care, or surgery.

Diagnosed with
ankylosing spondylitis?

[Learn about biologics](#) >

- Lidoderm patch will cause a numbing effect at the application site. Do not scratch, rub, or expose the area to extreme hot or cold temperatures until the numbness is gone.
- Do NOT use more medicine, apply more often, or use for longer than prescribed. Your condition will not improve faster, but the risk of side effects may be increased.
- Lidoderm patch may cause harm if it is chewed or swallowed. If you may have taken it by mouth, contact your [poison control center](#) or emergency room right away.
- Do not get the patch wet. It may not stick. Do not bathe, swim, or shower while you are wearing the patch.
- Do not place sources of heat (eg, heating pad, electric blanket) over the patch; they may increase the risk of Lidoderm patch's side effects.
- Use Lidoderm patch with caution in the ELDERLY; they may be more sensitive to its effects.
- Lidoderm patch should be used with extreme caution in CHILDREN; safety and effectiveness in children have not been confirmed. CHILDREN may be more sensitive to the effects of Lidoderm patch.
- PREGNANCY and BREAST-FEEDING: If you become pregnant, contact your doctor. You will need to discuss the benefits and risks of using Lidoderm patch while you are pregnant. Lidoderm patch is found in breast milk. If you are or will be breast-feeding while you use Lidoderm patch, check with your doctor. Discuss any possible risks to your baby.

Possible side effects of Lidoderm patch:

All medicines may cause side effects, but many people have no, or minor, side effects. Check with your doctor if any of these most COMMON side effects persist or become bothersome:

Mild redness, swelling, burning, blisters, bruising, irritation, change in skin color, or numbness at the application site.

Seek medical attention right away if any of these SEVERE side effects occur:

Severe allergic reactions (rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue; unusual hoarseness); chest, jaw, or left arm pain; confusion; dizziness or light-headedness; fainting; fast, slow, or irregular heartbeat; feeling of being unusually warm, cold, or numb; mood or mental changes; nervousness; numbness of an arm or leg; ringing in the ears or hearing changes; seizures; severe irritation, burning, or tingling at the patch site; shallow or slow breathing; shortness of breath; sudden, severe headache; tremors or twitching; unusual drowsiness; vision changes; vomiting.

This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your health care provider. Call your doctor for medical advice about side effects. To report side effects to the appropriate agency, please read the [Guide to Reporting Problems to FDA](#).

SUBOXONE®
Sublingual Film



[suboxone.com](#)

(buprenorphine and naloxone) CIII. Find a doctor.

[Withdrawal Ease - Home](#)

[Windows® 7 Driver Updates](#)

If OVERDOSE is suspected:

Contact 1-800-222-1222 (the American Association of Poison Control Centers), your local [poison control center](#), or emergency room immediately.

Proper storage of Lidoderm patch:

Store Lidoderm patch at 77 degrees F (25 degrees C). Brief storage between 59 and 86 degrees F (15 and 30 degrees C) is permitted. Store away from heat, moisture, and light. Do not store in the bathroom. Keep the child-resistant envelope sealed at all times when not in use. Keep Lidoderm patch out of the reach of children and away from pets.

General information:

- If you have any questions about Lidoderm patch, please talk with your doctor, pharmacist, or other health care provider.
- Lidoderm patch is to be used only by the patient for whom it is prescribed. Do not share it with other people.
- If your symptoms do not improve or if they become worse, check with your doctor.
- Check with your pharmacist about how to dispose of unused medicine.

This information should not be used to decide whether or not to take Lidoderm patch or any other medicine. Only your health care provider has the knowledge and training to decide which medicines are right for you. This information does not endorse any medicine as safe, effective, or approved for treating any patient or health condition. This is only a brief summary of general information about Lidoderm patch. It does NOT include all information about the possible uses, directions, warnings, precautions, interactions, adverse effects, or risks that may apply to Lidoderm patch. This information is not specific medical advice and does not replace information you receive from your health care provider. You must talk with your healthcare provider for complete information about the risks and benefits of using Lidoderm patch.

Issue Date: April 1, 2015

Database Edition 15.2.1.001

Copyright © 2015 Clinical Drug Information, LLC

Sciatica Exercises?

Stop: What You MUST know Before attempting to Treat your Sciatica



Disclaimer: This information should not be used to decide whether or not to take this medicine or any other medicine. Only your health care provider has the knowledge and training to decide which medicines are right for you. This information does not endorse any medicine as safe, effective, or approved for treating any patient or health condition. This is only a brief summary of general information about this medicine. It does NOT include all information about the possible uses, directions, warnings, precautions, interactions, adverse effects, or risks that may apply to this medicine. This information is not specific medical advice and does not replace information you receive from your health care provider. You must talk with your healthcare provider for complete information about the risks and benefits of using this medicine.

Google Facebook Twitter Print Email Add to Mednotes

Next Page → [Side Effects](#)

More about Lidoderm (lidocaine topical)

- [Side Effects](#)
- [During Pregnancy](#) or [Breastfeeding](#)
- [Dosage Information](#)
- [Drug Interactions](#)
- [Support Group](#)
- [Pricing & Coupons](#)
- [En Espanol](#)
- [22 Reviews](#) - [Add your own review/rating](#)
- [Generic Availability](#)

Consumer resources

- [Lidoderm](#)
- [Lidoderm Topical application](#) (Advanced Reading)
- *Other brands:* [Lidocaine Viscous](#), [Xylocaine Jelly](#), [RectiCare](#), [Xylocaine Viscous](#), [More \(27\)](#) »

Professional resources

- [Lidocaine Topical Oral Solution 4%](#) (FDA)

Related treatment guides

- [Pain](#)
- [Burns, External](#)
- [Anal Itching](#)
- [Hemorrhoids](#)
- [Anesthesia](#)
- [More \(3\) »](#)



Support

[Advertise With Us](#)
[Frequently Asked Questions](#)
[Ask a Question](#)
[Content Submissions](#)
[Mobile Apps](#)
[About Drugs.com](#)
[Contact Us](#)
[Sitemap](#)

Follow:     

About Drugs.com

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment.

Data sources include Micromedex® (updated Feb 4th, 2015), Cerner Multum™ (updated Feb 17th, 2015), Wolters Kluwer™ (updated Apr 6th, 2015) and others. To view content sources and attributions, refer to our [editorial policy](#).

Copyright © 2000-2015 Drugs.com. All rights reserved.

Your Privacy



We comply with the HONcode standard for trustworthy health information.
[Verify here](#)



Ad Choices

[Terms of Use](#) · [Privacy Policy](#)