

If you're an adult who
can't straighten your fingers

[Home](#) > [Drugs A to Z](#) > [Meloxicam](#)

[Print](#)  [Share](#)

Meloxicam



Generic Name: meloxicam (mel OKS i kam)

Brand Names: *Mobic*, *Vivlodex*

"48 Hours & Pain Was Gone"

Use This 1 Inflammation Trick, How Thousands Fight Inflammation in 1-Day!
Go to amazon.com/Inflammation



Overview

[Side Effects](#)

[Dosage](#)

[Interactions](#)

[Patient Tips](#)

[Professional](#)

[More](#) 

What is meloxicam?

Meloxicam is a nonsteroidal anti-inflammatory drug (NSAID). It works by reducing hormones that cause inflammation and pain in the body.

Meloxicam is used to treat pain or inflammation caused by [rheumatoid arthritis](#) and [osteoarthritis](#) in adults.

ADVERTISEMENT

If you're an adult who
can't straighten

Meloxicam is also used to treat juvenile rheumatoid arthritis in children who are at least 2 years old.

Important information

Meloxicam can increase your risk of fatal heart attack or stroke, especially if you use it long term or take high doses, or if you have heart disease. Do not use this medicine just before or after heart bypass surgery (coronary artery bypass graft, or CABG).

Get emergency medical help if you have chest pain, weakness, shortness of breath, slurred speech, or problems with vision or balance.

Meloxicam may also cause stomach or intestinal bleeding, which can be fatal. These conditions can occur without warning while you are using meloxicam, especially in older adults.

Call your doctor at once if you have symptoms of stomach bleeding such as black, bloody, or tarry stools, or coughing up blood or vomit that looks like coffee grounds.

Avoid drinking alcohol. It may increase your risk of stomach bleeding.

Ask a doctor or pharmacist before using any other cold, allergy, or pain medicine. Medicines similar to meloxicam are contained in many combination medicines. Check the label to see if a medicine contains an NSAID (non-steroidal anti-inflammatory drug) such as aspirin, ibuprofen, ketoprofen, or naproxen.

Meloxicam can increase your risk of fatal heart attack or stroke, especially if you use it long term or take high doses, or if you have heart disease. Even people without heart disease or risk factors could have a stroke or heart attack while taking this medicine.

Do not use this medicine just before or after heart bypass surgery (coronary artery bypass graft, or CABG).

Meloxicam may also cause stomach or intestinal bleeding, which can be fatal. These conditions can occur without warning while you are using meloxicam, especially in older adults.

You should not use meloxicam if you are allergic to it, or if you have ever had an asthma attack or severe allergic reaction after taking aspirin or an NSAID.

To make sure meloxicam is safe for you, tell your doctor if you have:

- heart disease, high blood pressure, high cholesterol, diabetes, or if you smoke;
- a history of heart attack, stroke, or blood clot;
- a history of stomach ulcers or bleeding;
- asthma;



SLIDESHOW

[Flexeril \(cyclobenzaprine\): How to Safely Use This Muscle Relaxant](#)

- kidney disease (or if you are on dialysis);
- liver disease; or
- fluid retention.

Taking meloxicam during the last 3 months of pregnancy may harm the unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

Meloxicam may cause a delay in ovulation (the release of an egg from an ovary). You should not take this medicine if you are undergoing fertility treatment, or are otherwise trying to get pregnant.

Meloxicam can pass into breast milk and may harm a nursing baby. You should not breast-feed while using this medicine.

Meloxicam is not approved for use by anyone younger than 2 years old.

How should I take meloxicam?

Take meloxicam exactly as prescribed by your doctor. Follow all directions on your prescription label. Your doctor may occasionally change your dose to make sure you get the best results. Do not take this medicine in larger amounts or for longer than recommended. Use the lowest dose that is effective in treating your condition.

You may take meloxicam with or without food.

Shake the oral suspension (liquid) well just before you measure a dose. Measure liquid medicine with the dosing syringe provided, or with a special dose-measuring spoon or medicine cup. If you do not have a dose-measuring device, ask your pharmacist for one.

If a child is taking this medication, tell your doctor if the child has any changes in weight. Meloxicam doses are based on weight in children.

If you use this medicine long-term, you may need frequent medical tests.

Store at room temperature, away from moisture and heat. Keep the bottle tightly closed when not in use.

Read all patient information, medication guides, and instruction sheets provided to you. Ask your doctor or pharmacist if you have any questions.



Simple Way to Tighten Skin



Surprisingly Simple Tip Removes Wrinkles & Eye Bags Without Surgery

Science of Dermaceuticals

See also: [Dosage Information \(in more detail\)](#)

What happens if I miss a dose?

Take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not take extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking meloxicam?

Avoid drinking alcohol. It may increase your risk of stomach bleeding.

Avoid taking aspirin while you are taking meloxicam.

Ask a doctor or pharmacist before using any cold, allergy, or pain medication. Many medicines available over the counter contain aspirin or other medicines similar to meloxicam. Taking certain products together can cause you to get too much of this type of medication. Check the label to see if a medicine contains aspirin, ibuprofen, ketoprofen, or naproxen.

ADVERTISEMENT

3 Fatigue-Causing Foods - Never eat these 3 foods

Boost your energy level every day by never eating these 3 foods

haveenergyatanyage.com



Stop Leg Cramps in 1 Min.

Proven Old Amish Remedy Stops Night Time Leg or Foot Cramps in 1 Minute

stopslegcramps.com





Meloxicam side effects

Get emergency medical help if you have signs of an allergic reaction to meloxicam: sneezing, runny or stuffy nose; wheezing or trouble breathing; hives; swelling of your face, lips, tongue, or throat.

Get emergency medical help if you have signs of a heart attack or stroke: chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath.

Stop using meloxicam and call your doctor at once if you have:

- the first sign of any skin rash, no matter how mild;
- shortness of breath (even with mild exertion);
- swelling or rapid weight gain;
- signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds;
- liver problems - nausea, upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes);
- kidney problems - little or no urinating, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath;
- low red blood cells (anemia) - pale skin, feeling light-headed or short of breath, rapid heart rate, trouble concentrating; or
- severe skin reaction - fever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling.

Common meloxicam side effects may include:

- upset stomach, nausea, vomiting, heartburn;
- diarrhea, constipation, gas;
- dizziness; or
- cold symptoms, flu symptoms.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

See also: [Side effects \(in more detail\)](#)

Meloxicam dosing information

Usual Adult Dose of Meloxicam for Osteoarthritis:

Initial dose: 7.5 mg orally once daily
Maintenance dose: 7.5 mg orally once daily
Maximum dose: 15 mg orally daily

Usual Adult Dose of Meloxicam for Rheumatoid Arthritis:

Initial dose: 7.5 mg orally once daily
Maintenance dose: 7.5 mg orally once daily
Maximum dose: 15 mg orally daily

Usual Pediatric Dose for Juvenile Rheumatoid Arthritis:

Greater than or equal to 2 years: 0.125 mg/kg orally once daily

Maximum dose: 7.5 mg orally daily

There was no additional benefit demonstrated by increasing the dose above 0.125 mg/kg once daily in clinical trials.

What other drugs will affect meloxicam?

Ask your doctor before using meloxicam if you take an antidepressant such as citalopram, escitalopram, fluoxetine (Prozac), fluvoxamine, paroxetine, sertraline (Zoloft), trazodone, or vilazodone. Taking any of these medicines with an NSAID may cause you to bruise or bleed easily.

Tell your doctor about all your current medicines and any you start or stop using, especially:

- cyclosporine;
- lithium;
- methotrexate;
- sodium polystyrene sulfonate (Kayexalate);
- a blood thinner (warfarin, Coumadin, Jantoven);
- heart or blood pressure medication, including a diuretic or "water pill"; or
- steroid medicine (such as prednisone).

This list is not complete. Other drugs may interact with meloxicam, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible interactions are listed in this medication guide.

Next → [Side Effects](#)

"48 Hours & Pain Was Gone"

Use This 1 Inflammation Trick, How Thousands Fight Inflammation in 1-Day!
Go to amazon.com/Inflammation




☐ Add to My Med List

More about meloxicam

- [Side Effects](#)
- [During Pregnancy](#) or [Breastfeeding](#)
- [Dosage Information](#)
- [Drug Images](#)
- [Drug Interactions](#)
- [Support Group](#)
- [Pricing & Coupons](#)
- [En Español](#)
- [299 Reviews](#) – [Add your own review/rating](#)
- [Drug class: nonsteroidal anti-inflammatory agents](#)

☐ Consumer resources

- [Meloxicam](#)
- [Meloxicam suspension](#)
- [Meloxicam](#) (Advanced Reading)

Other brands: [Mobic](#), [Vivlodex](#)

☐ Professional resources

- [Meloxicam](#) (AHFS Monograph)
- [More...](#) (3)

☐ Related treatment guides

- [Inflammatory Conditions](#)
- [Osteoarthritis](#)
- [Fibromyalgia](#)
- [Tendonitis](#)
- [Rheumatoid Arthritis](#)
- [Juvenile Rheumatoid Arthritis](#)

Where can I get more information?

- Your pharmacist can provide more information about meloxicam.
- Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use meloxicam only for the indication prescribed.
- Disclaimer: Every effort has been made to ensure that the information provided by Cerner Multum, Inc. ('Multum') is accurate, up-to-date, and complete, but no guarantee is made to that effect. Drug information contained herein may be time sensitive. Multum information has been compiled for use by healthcare practitioners and consumers in the United States and therefore Multum does not warrant that uses outside of the United States are appropriate, unless specifically indicated otherwise. Multum's drug information does not endorse drugs, diagnose patients or recommend therapy. Multum's drug information is an informational resource designed to assist licensed healthcare practitioners in caring for their patients and/or to serve consumers viewing this service as a supplement to, and not a substitute for, the expertise, skill, knowledge and judgment of healthcare practitioners. The absence of a warning for a given drug or drug combination in no way should be construed to indicate that the drug or drug combination is safe, effective or appropriate for any given patient. Multum does not assume any responsibility for any aspect of healthcare administered with the aid of information Multum provides. The information contained herein is not intended to cover all possible uses, directions, precautions, warnings, drug interactions, allergic reactions, or adverse effects. If you have questions about the drugs you are taking, check with your doctor, nurse or pharmacist.

Copyright 1996-2017 Cerner Multum, Inc. Version: 12.03. Revision Date: 2015-12-04, 11:20:16 AM.

DRUG STATUS

- 

Availability

Prescription only
- 

Pregnancy Category

Risk depends on usage

☐
- 

CSA Schedule

Not a controlled drug

☐
- 

Approval History

Drug history at FDA

☐



Drug Class

Nonsteroidal anti-inflammatory agents



Related Drugs

Osteoarthritis
[prednisone](#), [naproxen](#), [Cymbalta](#), [aspirin](#), [ibuprofen](#), [duloxetine](#), [diclofenac](#), [More...](#)



Rheumatoid Arthritis
[Humira](#), [Plaquenil](#), [hydroxychloroquine](#), [Enbrel](#), [Remicade](#), [sulfasalazine](#), [leflunomide](#), [Rituxan](#), [More...](#)

Juvenile Rheumatoid Arthritis
[prednisone](#), [naproxen](#), [aspirin](#), [ibuprofen](#), [Celebrex](#), [Mobic](#), [dexamethasone](#), [Aleve](#), [More...](#)



[View all 45 images](#)

Related News and Articles

[FDA Throws Cold Water on Whole Body Cryotherapy](#)
March 3, 2017

[Can Depression Up Odds for Arthritis Linked to Psoriasis?](#)
February 24, 2017

[ACL Tears on the Rise Among Kids, Especially Girls](#)
February 22, 2017

ADVERTISEMENT

Drugs.com Mobile Apps

The easiest way to lookup drug information, identify pills,

check interactions and set up your own personal medication records. Available for Android and iOS devices.

Explore Apps

SUPPORT

[Help Center](#)

[Frequent Questions](#)

[Sitemap](#)

[Contact Us](#)

ABOUT

[About Drugs.com](#)

[Advertising Policy](#)

[Content Submissions](#)

[Drugs.com Blog](#)

TERMS & PRIVACY

[Editorial Policy](#)

[Privacy Policy](#)

[Terms of Use](#)

[Attribution & Citations](#)



[Subscribe](#) to receive email notifications whenever new articles are published.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex® (updated Mar 2nd, 2017), Cerner Multum™ (updated Mar 2nd, 2017), Wolters Kluwer™ (updated Mar 1st, 2017) and others. To view content sources and attributions, please refer to our [editorial policy](#).



Ad Choices



We comply with the HONcode standard for trustworthy health information - [verify here](#)

Copyright © 2000-2017 Drugs.com. All rights reserved.