
Visited on 09/01/2017



Hees

Uses of This Medicine

Hypoglycemic (About this - PubMed Health)

Glipizide is used to treat high blood sugar levels caused by a type of diabetes mellitus (sugar diabetes) called type 2 diabetes. In type 2 diabetes, your body does not work properly to store excess sugar and the sugar remains in your bloodstream.

Chronic high blood sugar can lead to serious health problems in the future.

Proper diet is the first step in managing type 2 diabetes, but often medicines are needed to help your body. Glipizide belongs to a class of drugs called sulfonylureas. It stimulates the release of insulin from the pancreas, directing your body to store blood sugar. This helps lower blood sugar and restore the way you use food to make energy.

This medicine is available only with your doctor's prescription

Other uses (PubMed Health)

How To Use

Tablet, Long Acting Tablet

Take your medicine as directed. Your dose may need to be changed several times to find what works best for you.

Take the **regular tablet** about 30 minutes before you eat a meal. Take the **extended-release tablet** with breakfast, unless your doctor tells you differently.

Swallow the extended-release tablet whole. Do not crush, break, or chew it.

Read and follow the patient instructions that come with this medicine. Talk to your doctor or pharmacist if you have any questions.

Missed dose: Take a dose as soon as you remember. If it is almost time for your next dose, wait until then and take a regular dose. Do not take extra medicine to make up for a missed dose.

Store the medicine in a closed container at room temperature, away from heat, moisture, and direct light.

Drugs and Foods to Avoid

Ask your doctor or pharmacist before using any other medicine, including over-the-counter medicines, <u>vitamins</u>, and herbal products.

Some medicines can affect how glipizide works. Tell your doctor if you are using any of the following:


- Aspirin, chloramphenicol, fluconazole, isoniazid, miconazole, niacin, phenytoin, probenecid
- Birth control pill
- Beta blocker medicine
- Blood thinner (including warfarin)
- Diuretic (water pill)
- MAO inhibitor
- Medicine to treat an infection (including ciprofloxacin, levofloxacin, moxifloxacin)
- NSAID pain or arthritis medicine (including celecoxib, diclofenac, ibuprofen, naproxen)
- Phenothiazine medicine
- · Steroid medicine
- Sulfa drug
- <u>Thyroid</u> medicine

If you are also taking <u>colesevelam</u>, take it at least 4 hours after you take <u>glipizide</u>.

When Not To Use

This medicine is not right for everyone. Do not use it if you had an allergic reaction to glipizide.

Warnings





PubMed Health

See more..

Approved drug uses

More information about a drug

Visited on 09/01/2017

problems, stomach or bowel problems, adrenal or pituitary gland problems, or G6PD deficiency. Tell your doctor if you drink alcohol

This medicine may cause the following problems:

- · Higher risk of heart or blood vessel problems
- · Low blood sugar

Part of the extended-release tablet may pass in your stool. This is normal

Your doctor will do lab tests at regular visits to check on the effects of this medicine. Keep all appointments.

Keep all medicine out of the reach of children. Never share your medicine with anyone.

Possible side effects

Summary More details

Call your doctor right away if you notice any of these side effects:

Allergic reaction: Itching or hives, swelling in your face or hands, swelling or tingling in your mouth or throat, chest tightness, trouble breathing

Blurred vision, changes in vision

Fast or pounding heartbeat, lightheadedness, dizziness

Headache or confusion

Sweating, trembling, shakiness, increased hunger

If you notice these less serious side effects, talk with your doctor:

Mild nausea, diarrhea, stomach upset

If you notice other side effects that you think are caused by this medicine, tell your doctor.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

More side effects of this drug

Brand names include

Glucotrol, Glucotrol XL

There may be other brand names for this medicine.

More detailed version of this drug page



Truven Health Analytics, an IBM company, Disclaimer

Portions Copyright © 2017. Truven Health Analytics™, an IBM company. Information is for End User's use only and may not be sold, redistributed or otherwise used for commercial purposes. Use of this content is subject to Truven Health Analytics' full disclaimer, available above







PubMed Health Home | About PubMed Health | Copyright | Disclaimer | 📳 See us on Facebook | 📡 Follow us | 🎏 Add us | Contact Us

You are here: NCBI > Literature > PubMed Health

Support Cente

GETTING STARTED POPULAR FEATURED RESOURCES NCBI Education Chemicals & Bioassays PubMed Genetic Testing Registry Bookshelf PubMed Health NCBI Help Manual Data & Software PubMed Central NCBI Handbook Training & Tutorials Domains & Structures PubMed Health Reference Sequences Genes & Expression Gene Expression Omnibus Genetics & Medicine Nucleotide Genomes & Maps Genome Human Genome SNP Homology Mouse Genome Literature Gene Influenza Virus Sequence Analysis PubChem Sequence Read Archive Taxonomy

National Center for Biotechnology Information, U.S. National Library of Medicine 8600 Rockville Pike, Bethesda MD, 20894 USA







NCBI INFORMATION

About NCBI

NCBI FTP Site

Research at NCBI

NCBI News & Blog

NCBI on Facebook

NCBI on Twitter

NCBI on YouTube



