



Search

Browse all medications

ABCDEFGHIJKLMNOPQRSTUVWXYZ

Advanced Search

Phonetic Search

- Home

Drugs A to Z

Pill Identifier

Interactions Checker

News & Alerts

Health Professionals

Community

- Mednotes
- Drugs A to Z

Drugs by Condition

Drugs by Class

Side Effects

Dosage

Pregnancy

Natural Products

Vet Drugs

Symptom Checker

Harvard Health Topics

Twitter

Facebook

LinkedIn

Home > Drugs A to Z > C > Co > Compazine Consumer Information

Print Save or Share

Compazine

Generic Name: [prochlorperazine](#) (oral) (pro klor PER a zeen)
Brand Names: *Compazine*

- Ads by Google
- Free prescription assist

[www.yourrxdiscounts.com](#)

Paying out of pocket? Save up to 75% on your prescription drugs
- Lower Blood Pressure Fast

[www.l-arginineplus.com](#)

Nobel Prize Winning Breakthrough Can L-arginine Plus Help You?
- Psychiatric Drug Facts

[CCHR.org](#)

Do You Know The Long Term Effects Of Psychiatric Drugs? Free Info Kit

What is oral prochlorperazine?

Prochlorperazine is an anti-psychotic medication in a group of drugs called phenothiazines (FEEN-oh-THYE-a-zeens). It works by changing the actions of chemicals in your brain.

Prochlorperazine is used to treat psychotic disorders such as schizophrenia. It is also used to treat anxiety, and to control severe nausea and vomiting.

Prochlorperazine may also be used for other purposes not listed in this medication guide.

ADVERTISEMENT

Save up to 65% on your Prescriptions.

plus get a FREE Pharmacy Card

Click Here

SIMPLE SAVINGS CARD™

What is the most important information I should know about oral prochlorperazine?

Stop using this medication and call your doctor at once if you have twitching or uncontrollable movements of your eyes, lips, tongue, face, arms, or legs. These could be early signs of dangerous side effects. Prochlorperazine is not for use in psychotic conditions related to dementia. Prochlorperazine may cause heart failure, sudden death, or pneumonia in older adults with dementia-related conditions. Do not use prochlorperazine if you have brain damage, bone marrow depression, or are also using large amounts of alcohol or medicines that make you sleepy. Do not use if you are allergic to prochlorperazine or other phenothiazines. Before you take prochlorperazine, tell your doctor if you have glaucoma, heart disease or high blood pressure, liver or kidney disease, severe asthma or breathing problems, a history of

Sign In or Register

Forgot password?

username

password

Sign In

Related Compazine Information

10+ years

Approval History

FDA approved 1956

N/A

CSA Schedule

Not a controlled drug

OTC Rx

Availability

Rx and/or OTC

This document has been reviewed by one or more clinical specialists. See our [editorial policy](#) for more information.

Reviews

Average User Rating

8 User Reviews

6.7 Rate it!

- Related Pages
- Detailed Side Effects

Drug Interactions

Drug Images

Pregnancy & Breastfeeding

Dosage Information

Support Group Q & A

Drug Class

Phenothiazine antiemetics

Phenothiazine antipsychotics

ADVERTISEMENT

Tailorsville

FREE SHIPPING + FAST DELIVERY

Newton

FREE

Save up to \$100

- Related Drugs
- Nausea/ Vomiting

[promethazine](#)

[lorazepam](#)

[Ativan](#)

[hydroxyzine](#)

[diphenhydramine](#)

[Benadryl](#)

[Phenergan](#)

[Zofran](#)

[More...](#)
- Anxiety

[Xanax](#)

[Lexapro](#)

[Cymbalta](#)

[alprazolam](#)

[lorazepam](#)

[atenolol](#)

[Valium](#)

[Ativan](#)

[More...](#)

seizures, adrenal gland tumor, Parkinson's disease, enlarged prostate or urination problems, an infectious disease (such as chickenpox, measles, flu, or central nervous system infection), past or present breast cancer, low levels of calcium in your blood, or if you have ever had a serious side effect while using prochlorperazine or similar medicines.

Before taking prochlorperazine, tell your doctor about all other medications you use.

What should I discuss with my healthcare provider before taking oral prochlorperazine?

Prochlorperazine is not for use in psychotic conditions related to dementia. Prochlorperazine may cause heart failure, sudden death, or pneumonia in older adults with dementia-related conditions. Do not use prochlorperazine if you have brain damage, bone marrow depression, or are also using large amounts of alcohol or medicines that make you sleepy. Do not use if you are allergic to prochlorperazine or other phenothiazines such as chlorpromazine (Thorazine), fluphenazine (Permitil), perphenazine (Trilafon), promethazine (Adgan, Pentazine, Phenergan), thioridazine (Mellaril), or trifluoperazine (Stelazine). If you have certain conditions, you may need a dose adjustment or special tests to safely use this medication. Before you take prochlorperazine, tell your doctor if you have:

- glaucoma;
- heart disease or high blood pressure;
- liver or kidney disease;
- severe asthma, emphysema, or other breathing problem;
- a history of seizures;
- adrenal gland tumor (pheochromocytoma);
- Parkinson's disease;
- an enlarged prostate or urination problems;
- an infectious disease such as chickenpox, measles, stomach flu, or an infection of the central nervous system;
- past or present breast cancer;
- low levels of calcium in your blood (hypocalcemia); or
- if you have ever had a serious side effect while using prochlorperazine or another phenothiazine.

Tell your doctor if you will be exposed to extreme heat or cold, or to insecticide poisons while you are taking prochlorperazine.

It is not known whether prochlorperazine will harm an unborn baby. Prochlorperazine may cause side effects in a newborn if the mother takes the medication during pregnancy. Do not take this medication without first talking to your doctor if you are pregnant. Tell your doctor if you become pregnant while taking prochlorperazine. Prochlorperazine can pass into breast milk and may harm a nursing baby. Do not use this medication without telling your doctor if you are breast-feeding a baby. Older adults may be more likely to have side effects from this medication. Prochlorperazine is not for use in children younger than 2 years old or weighing less than 20 pounds. Talk with your doctor before giving this medication to a child who has been ill with a fever or flu symptoms.

Ads by Google

[SUBOXONE® Sublingual Film](#)
(buprenorphine and naloxone) CIII: Is here: learn how it's different
[SUBOXONE.com](#)

[Depression Treatment](#)
Find Treatments For Depression That Will Get You Back On Your Feet Now.
[Topicologist.com](#)

[Try OMNARIS® For Less](#)
Pay as low as \$11 for each of your 12 OMNARIS® (ciclesonide) Rx's.
[www.OMNARIS.com](#)



Psychosis
[Haldol](#)
[prochlorperazine](#)
[haloperidol](#)
[Thorazine](#)
[chlorpromazine](#)
[perphenazine](#)
[Prolixin](#)
[Trilafon](#)
[More...](#)

FDA Consumer Updates

- **FDA Intends to Remove Unapproved Drugs from Market**
- **Beware of Fraudulent Weight-Loss 'Dietary Supplements'**
- **A Guide to Safe Use of Pain Medicine**
- **Primatene Mist With Chlorofluorocarbons No Longer Available After Dec. 31, 2011**

[More FDA Updates...](#)

Drug Images



[View larger images](#)

Get Updates by Email

Email me about **Compazine:**

News and Warnings related to this drug

Also send the Drugs.com monthly newsletter
FDA Medwatch Alerts for all medications

Related News and Articles

Antipsychotic Drugs Raise Heart Risks, Experts Warn
18 Feb 2011
Doctors must not forget about the physical health of people with serious mental health disorders who take antipsychotic drugs, experts warn. The authors of ...

Schizophrenia Drugs May Spur Subtle Brain Tissue Loss

How should I take oral prochlorperazine?

Take this medication exactly as prescribed by your doctor. Do not take it in larger amounts or for longer than recommended. Follow the directions on your prescription label. Your doctor may occasionally change your dose to make sure you get the best results from this medication.

This medication can cause you to have unusual results with certain medical tests. Tell any doctor who treats you that you are using prochlorperazine.

If you need to have an x-ray or CT scan of your spinal column using a dye that is injected into a vein, you may need to temporarily stop taking prochlorperazine. Be sure the doctor knows ahead of time that you are using this medication.

Do not stop using prochlorperazine suddenly after long-term use, or you could have unpleasant withdrawal symptoms such as nausea, vomiting, dizziness, or feeling shaky. Talk to your doctor about how to avoid withdrawal symptoms when you stop using prochlorperazine. Store prochlorperazine at room temperature away from moisture, heat, and light.

See also: [Compazine dosage](#) (in more detail)

What happens if I miss a dose?

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention if you think you have used too much of this medicine. Overdose symptoms may include dry [mouth](#), constipation, bloating or stomach cramps, extreme drowsiness or feeling restless and agitated, changes in heart rate, fever, and fainting.

What should I avoid while taking oral prochlorperazine?

Prochlorperazine can cause side effects that may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be awake and alert. Avoid getting up too fast from a sitting or lying position, or you may feel dizzy. Get up slowly and steady yourself to prevent a fall. Avoid drinking alcohol. It can increase some of the side effects of prochlorperazine. Avoid exposure to sunlight or tanning beds. Prochlorperazine can make your skin more sensitive to sunlight, and a sunburn may result. Wear sunscreen (SPF 15 or higher) and protective clothing if you must be outdoors.

Oral prochlorperazine side effects

Get emergency medical help if you have any of these signs of an allergic reaction: hives; difficulty breathing; swelling of your face, lips, tongue, or throat. Stop using prochlorperazine and call your doctor at once if you have a serious side effect such as:

- twitching or uncontrollable movements of your eyes, lips, tongue, face, arms, or legs;
- tremor (uncontrolled shaking), drooling, trouble swallowing, problems with balance or walking;
- feeling restless, jittery, or agitated;
- high fever, stiff muscles, confusion, sweating, fast or uneven heartbeats, rapid breathing;
- feeling like you might pass out;
- seizure (black-out or convulsions);
- decreased night vision, tunnel vision, watery eyes, increased sensitivity to light;
- nausea and stomach pain, skin rash, and jaundice (yellowing of the skin or eyes);
- pale skin, easy bruising or bleeding, fever, sore throat, flu symptoms;
- urinating less than usual or not at all;

10 Feb 2011

MONDAY, Feb. 7 -- The use of antipsychotic drugs to treat schizophrenia is associated with the loss of a small but measurable amount of brain tissue, a new ...

Antipsychotics, Conventional and Atypical FDA Medwatch Alert

16 Jun 2008

FDA notified healthcare professionals that both conventional and atypical antipsychotics are associated with an increased risk of mortality in elderly ...

Related Questions & Answers

- [Can Compazine be safely used to halt a prolonged case of hiccups?](#)
- [Is Compazine a narcotic?](#)
- [Is compazine a narcotic?](#)
- [Can Compazine be used when using phenytoin?](#)
- [I recently went to my dr with migrains. He said he was perscribing imitrex. My persription reads?](#)

 [More Compazine questions](#)

ADVERTISEMENT

Lower Your Blood Pressure - The All Natural Way
The only non-drug medical device clinically proven to lower blood pressure [www.resperate.com](#)

\$350,000 in Life Insurance
Protect Your Family Today. Coverage as low as \$13.04/month. [Free Quote Here](#) [BestQuoteMarket.com](#)

Weight Loss Medicine
Immediate weight loss - reduces fat absorbtion - less fat means less pounds [www.pharmcom.com](#)

- joint pain or swelling with fever, swollen glands, muscle aches, chest pain, vomiting, unusual thoughts or behavior, and patchy skin color; or
- slow heart rate, weak pulse, fainting, slow breathing (breathing may stop).

Less serious side effects may include:

- dizziness, drowsiness, anxiety;
- sleep problems (insomnia), strange dreams;
- dry mouth, stuffy nose;
- blurred vision;
- constipation;
- breast swelling or discharge;
- a missed menstrual period;
- weight gain, swelling in your hands or feet;
- impotence, trouble having an orgasm;
- mild itching or skin rash; or
- headache.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

[Ads by Google](#)

[Phentramine HCL 37.5 mg](#)

Free Shipping options Low price \$41 No script - No Hidden Fees!

[pghp.com/phentramine](#)

[#1 Addiction Rehab Center](#)

Adult, Addiction Rehab & Detox All Private Rooms, Insurance OK

[TransformationsTreatment.com](#)

[Managing Side Effects](#)

There is hope. Chat privately with our oncology info experts today.

[CancerCenter.com/CareThatNeverQuits](#)

What other drugs will affect oral prochlorperazine?

Cold or allergy medicine, sedatives, narcotic pain medicine, sleeping pills, muscle relaxers, and medicine for seizures, depression or anxiety can interact with prochlorperazine and cause medical problems or increase side effects. Tell your doctor if you regularly use any of these medicines, or any other anti-psychotic medications.

Also tell your doctor if you are taking any of the following medicines:

- atropine (Atreza, Sal-Tropine);
- lithium (Eskalith, Lithobid);
- a diuretic (water pill);
- an antibiotic;
- birth control pills or hormone replacement estrogens;
- blood pressure medication;
- a blood thinner such as warfarin (Coumadin);
- certain asthma medications or bronchodilators;
- drugs to treat a prostate disorder, such as alfuzosin (Uroxatral), doxazosin (Cardura), prazosin (Minipress), terazosin (Hytrin), tamsulosin (Flomax);
- incontinence medications;
- insulin or diabetes medications you take by mouth;
- medication for nausea, vomiting, or motion sickness;
- medications to treat or prevent malaria;

- medications used for general anesthesia;
- medicines used to prevent organ transplant rejection;
 - numbing medicine such as lidocaine or Novocain;
 - a stimulant or ADHD medication;
 - ulcer or [irritable bowel](#) medications; or
 - medicines to treat Parkinson's disease, restless leg syndrome, or pituitary gland tumor (prolactinoma).

This list is not complete and there are many other medicines that can interact with prochlorperazine. Tell your doctor about all your prescription and over-the-counter medications, vitamins, minerals, herbal products, and drugs prescribed by other doctors. Do not start a new medication without telling your doctor. Keep a list with you of all the medicines you use and show this list to any doctor or other healthcare provider who treats you.

More Compazine resources

- [Compazine Side Effects](#) (in more detail)
- [Compazine Dosage](#)
- [Compazine Use in Pregnancy & Breastfeeding](#)
- [Drug Images](#)
- [Compazine Drug Interactions](#)
- [Compazine Support Group](#)
- [8 Reviews for Compazine](#) - [Add your own review/rating](#)

- Med Facts

[Compazine MedFacts Consumer Leaflet](#) (Wolters Kluwer)
- Professional

[Compazine Prescribing Information](#) (FDA)
- Professional

[Prochlorperazine Prescribing Information](#) (FDA)
- Professional

[Prochlorperazine Professional Patient Advice](#) (Wolters Kluwer)
- Professional

[Prochlorperazine Monograph](#) (AHFS DI)
- Med Facts

[Compro Suppositories MedFacts Consumer Leaflet](#) (Wolters Kluwer)
- Professional

[Compro Prescribing Information](#) (FDA)

Compare Compazine with other medications

- [Anxiety](#)
- [Hiccups](#)
- [Nausea/Vomiting](#)
- [Psychosis](#)

Where can I get more information?

- Your pharmacist can provide more information about oral prochlorperazine.

 [Printable Version](#)

 [Email](#)

[Recommend](#)

 [Tweet](#)

 [Save or Share](#)

Ads by Google

Drug Safety Information

Get Specialized Resources At The Official Pfizer Professional Site.
www.PfizerPro.com/Relpax

Bladder Neuromodulation

Innovative treatment option for OAB when medication doesn't work.
www.everyday-freedom.com

- Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.
- Every effort has been made to ensure that the information provided by Cerner Multum, Inc. ('Multum') is accurate, up-to-date, and complete, but no guarantee is made to that effect. Drug information contained herein may be time sensitive. Multum information has been compiled for use by healthcare practitioners and consumers in the United States and therefore Multum does not warrant that uses outside of the United States

are appropriate, unless specifically indicated otherwise. Multum's drug information does not endorse drugs, diagnose patients or recommend therapy. Multum's drug information is an informational resource designed to assist licensed healthcare practitioners in caring for their patients and/or to serve consumers viewing this service as a supplement to, and not a substitute for, the expertise, skill, knowledge and judgment of healthcare practitioners. The absence of a warning for a given drug or drug combination in no way should be construed to indicate that the drug or drug combination is safe, effective or appropriate for any given patient. Multum does not assume any responsibility for any aspect of healthcare administered with the aid of information Multum provides. The information contained herein is not intended to cover all possible uses, directions, precautions, warnings, drug interactions, allergic reactions, or adverse effects. If you have questions about the drugs you are taking, check with your doctor, nurse or pharmacist.

Copyright 1996-2006 Cerner Multum, Inc. Version: 7.04. Revision Date: 12/15/2010 5:01:39 PM.

Q enter a search term

Search

Ads by Google [Drug Rehab](#) [Drug Treatment](#) [Holistic Drug](#) [Drug Clinic](#)

Support	About Drugs.com	Your Privacy
Advertise With Us Frequently Asked Questions Pharmaceutical Companies Widgets for your Website Content Submissions About Drugs.com Contact Us Sitemap Follow us on:  	Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines & natural products. This material is provided for educational purposes only and is not to be used for medical advice, diagnosis or treatment. Data sources include Micromedex™ (updated Apr 2nd, 2011), Cerner Multum™ (updated Mar 18th, 2011), Wolters Kluwer™ (updated Apr 4th, 2011) and others. To view content sources and attributions, refer to our editorial policy. Copyright © 2000-2011 Drugs.com. All rights reserved.	TRUSTe CERTIFIED PRIVACY Verify Here We comply with the HONcode standard for trustworthy health information. Verify here Ad Choices Terms of Use · Privacy Policy