

ADVERTISEMENT



Find us on:
facebook®

Get news & drug alerts [GO](#)



Drug Information Online

Search

Browse all medications [A](#) [B](#) [C](#) [D](#) [E](#) [F](#) [G](#) [H](#) [I](#) [J](#) [K](#) [L](#) [M](#) [N](#) [O](#) [P](#) [Q](#) [R](#) [S](#) [T](#) [U](#) [V](#) [W](#) [X](#) [Y](#) [Z](#) [Advanced Search](#) [Phonetic Search](#)

[Home](#)

[Drugs A to Z](#)

[Pill Identifier](#)

[Interactions Checker](#)

[News & Alerts](#)

[Health Professionals](#)

[Community](#)

Mednotes

[Drugs A to Z](#) [Drugs by Condition](#) [Drugs by Class](#) [Side Effects](#) [Dosage](#) [Pregnancy](#) [Natural Products](#) [Vet Drugs](#) [Symptom Checker](#) [Harvard Health Topics](#) [Twitter](#) [Facebook](#) [RSS](#)

[Home](#) > [Drugs by Condition](#) > [M](#) > [Muscle Spasm](#) > Flexeril Consumer Information [Print](#) [Save or Share](#)

Flexeril

Generic Name: cyclobenzaprine (syeh kloe BEN zay preen)
Brand Names: *Amrix*, *Fexmid*, *Flexeril*

Ads by Google

- [SUBOXONE® Sublingual Film](#)
[SUBOXONE.com](#) (buprenorphine and naloxone) CIII: Is here: learn how it's different
- [Breakthrough Back Pain..](#)
[www.TexasSpineClinic.com](#) Imagine Being Pain Free. Make Your Dream a Reality. FDA Cleared!
- [Hydrocodone Rapid Detox](#)
[www.Opiates.com/hydrocodone](#) Safe Detox from Hydrocodone with Renowned Waismann Method.

What is Flexeril?

Flexeril (cyclobenzaprine) is a muscle relaxant. It works by blocking nerve impulses (or pain sensations) that are sent to your brain.

Flexeril is used together with rest and physical therapy to treat skeletal muscle conditions such as pain or injury.

Flexeril may also be used for other purposes not listed in this medication guide.

Important information about Flexeril

Do not take Flexeril if you have used an MAO inhibitor such as isocarboxazid (Marplan), tranylcypromine (Parnate), phenelzine (Nardil), or selegiline (Eldepryl, Emsam) within the past 14 days. Serious, life-threatening side effects can occur if you take Flexeril before the MAO inhibitor has cleared from your body.

You should not take Flexeril if you have recently had a heart attack, or if you have a heart rhythm disorder, congestive heart failure, heart block, or an overactive thyroid.

Flexeril can cause side effects that may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be awake and alert. Avoid drinking alcohol, which can

ADVERTISEMENT

Save up to 65%
on your *Prescriptions.*



plus get a
FREE
Pharmacy Card

Click Here

SIMPLE SAVINGS CARD™

Related Video



[Rheumatoid Arthritis](#)
Learn the signs of RA and how to relieve the pain.

Click to Play

Sign In or Register

Forgot password?

username

password

Sign In

Related Flexeril Information

10+ years

Approval History
FDA approved 1977

B

Pregnancy Category
No proven risk in humans

Rx

Availability
Prescription only

N/A

CSA Schedule
Not a controlled drug

Reviews

Average User Rating

67 User Reviews

6.6

Rate it!

Related Pages

[Detailed Side Effects](#)
[Drug Interactions](#)
[Drug Images](#)

[Pregnancy & Breastfeeding](#)
[Dosage Information](#)
[Support Group Q & A](#)

Drug Class

Skeletal muscle relaxants

ADVERTISEMENT

Related Drugs

Muscle Spasm
[cyclobenzaprine](#)
[Soma](#)
[Valium](#)
[diazepam](#)
[baclofen](#)
[methocarbamol](#)
[tizanidine](#)
[carisoprodol](#)
[Skelaxin](#)
[Robaxin](#)
[Zanaflex](#)
[metaxalone](#)
[More...](#)

Sciatica
[cyclobenzaprine](#)
[naproxen](#)
[acetaminophen](#)
[Tylenol](#)
[ibuprofen](#)
[aspirin](#)
[diclofenac](#)
[Paracetamol](#)

[indomethacin](#)
[Voltaren](#)
[Naprosyn](#)
[Aleve](#)
[More...](#)

increase some of the side effects of Flexeril.

Before taking Flexeril

Do not take Flexeril if you have used an MAO inhibitor such as isocarboxazid (Marplan), tranylcypromine (Parnate), phenelzine (Nardil), or selegiline (Eldepryl, Emsam) within the past 14 days. Serious, life-threatening side effects can occur if you take Flexeril before the MAO inhibitor has cleared from your body. Do not use Flexeril if you have recently had a heart attack, or if you have:

- a heart rhythm disorder;
- congestive heart failure;
- heart block; or
- an overactive thyroid.

Before using Flexeril, tell your doctor if you are allergic to any drugs, or if you have:

- problems with urination;
- enlarged prostate;
- glaucoma; or
- liver disease.

If you have any of these conditions, you may need a dose adjustment or special tests to safely take Flexeril.

FDA pregnancy category B. Flexeril is not expected to be harmful to an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant during treatment. It is not known whether cyclobenzaprine passes into breast milk or if it could harm a nursing baby. Do not use this medication without telling your doctor if you are breast-feeding a baby.

See also: [Flexeril pregnancy and breastfeeding warnings](#) (in more detail)

Older adults may be more sensitive to the side effects of Flexeril.

Ads by Google

Back Pain Medication

Safe Outpatient Procedure. Relief From Back Pain In Less Than 1 Week!

www.LaserSpineInstitute.com/Back

Managing Side Effects

There is hope. Chat privately with our oncology info experts today.

CancerCenter.com/CareThatNeverQuits

Effective Drug Treatment

Drug Treatment and Rehabilitation Programs. Call (877) 680-8223

www.TheRecoveryPlace.net/Treatment

How should I take Flexeril?

Take this medication exactly as it was prescribed for you. Do not take Flexeril in larger amounts, or take it for longer than recommended by your doctor. Follow the directions on your prescription label.

Take this medicine with a full glass of water.

Flexeril is only part of a complete program of treatment that may also include rest, physical therapy, or other pain relief measures. Follow your doctor's instructions.

Drug Images



[View larger images](#)

Get Updates by Email

Email me about Flexeril:

News and Warnings related to this drug

Also send the Drugs.com monthly newsletter
FDA Medwatch Alerts for all medications

[Get Notified](#)

ADVERTISEMENT

Lower Your Blood Pressure - The All Natural Way

The only non-drug medical device clinically proven to lower blood pressure www.resperate.com

\$350,000 in Life Insurance

Protect Your Family Today. Coverage as low as \$13.04/month. Free Quote Here BestQuoteMarket.com

Weight Loss Medicine

Immediate weight loss - reduces fat absorption - less fat means less pounds www.pharmcom.com

Help and Support

Looking for answers? [Ask a question](#) or go join the [Flexeril support group](#) to connect with others who have similar interests.

Store Flexeril at room temperature away from moisture, heat, and light.

See also: [Flexeril dosage](#) (in more detail)

What happens if I miss a dose?

Take the missed dose as soon as you remember. If it is almost time for your next dose, skip the missed dose and take the medicine at the next regularly scheduled time. Do not take extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention if you think you have used too much of this medicine. An overdose of Flexeril can be fatal.

Overdose symptoms may include drowsiness, fast heartbeat, tremors or shaking, slurred speech, confusion, nausea, vomiting, hallucinations (seeing things), chest pain, or seizure (convulsions).

What should I avoid while taking Flexeril?

Flexeril can cause side effects that may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be awake and alert. Avoid drinking alcohol, which can increase some of the side effects of Flexeril. Cold or allergy medicine, narcotic pain medicine, sleeping pills, and medicine for seizures, depression or anxiety can add to sleepiness caused by Flexeril. Tell your doctor if you regularly use any of these medicines, or any other muscle relaxer.

[Ads by Google](#)

[Neuropathy Breakthrough](#)

Natural Solution That Works. Learn How To Relieve Your Pain.

[Neuropathy.RealFoodNutrients.com](#)

[Instant Back Pain Relief](#)

Free yourself from back pain today get your free trial pack here.

[www.elmoreoil.com.au](#)

[Sciatica Pain Relief](#)

Do you Suffer from Sciatica Pain? Request An Instant Sciatica Guide!

[MySciaticaExercises.com/Sciatica](#)

Flexeril side effects

Get emergency medical help if you have any of these signs of an allergic reaction to Flexeril: hives; difficulty breathing; swelling of your face, lips, tongue, or throat. Stop using Flexeril and call your doctor at once if you have any of these serious side effects:

- fast, pounding, or uneven heartbeats;
- chest pain or heavy feeling, pain spreading to the arm or shoulder, nausea, sweating, general ill feeling;
- sudden numbness or weakness, especially on one side of the body;
- sudden headache, confusion, problems with vision, speech, or balance;
- feeling light-headed, fainting;
- confusion, weakness, lack of coordination;
- nausea, stomach pain, low fever, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes);
- seizure (convulsions);
- unusual thoughts or behavior, hallucinations (seeing things); or

- easy bruising or bleeding, unusual weakness.

Less serious Flexeril side effects may include:

- dry mouth or throat;
- blurred vision;
- drowsiness, dizziness, tired feeling;
- loss of appetite, stomach pain, nausea;
- diarrhea, constipation, gas; or
- muscle weakness.

This is not a complete list of side effects and others may occur. Tell your doctor about any unusual or bothersome side effect. You may report side effects to FDA at 1-800-FDA-1088.

See also: [Flexeril side effects](#) (in more detail)

What other drugs will affect Flexeril?

Many drugs can interact with Flexeril. Below is just a partial list. Tell your doctor if you are using:

- atropine (Donnatal, and others), benztropine (Cogentin), dimenhydrinate (Dramamine), methscopolamine (Pamine), or scopolamine (Transderm-Scop);
- a bronchodilator such as ipratroprium (Atrovent) or tiotropium (Spiriva);
- glycopyrrolate (Robinul);
- guanethidine (Ismelin);
- mepenzolate (Cantil);
- tramadol (Ultram);
- bladder or urinary medications such as darifenacin (Enablex), flavoxate (Urispas), oxybutynin (Ditropan, Oxytrol), tolterodine (Detrol), or solifenacin (Vesicare); or
- irritable bowel medications such as dicyclomine (Bentyl), hyoscyamine (Anaspaz, Cystospaz, Levsin, and others), or propantheline (Pro-Banthine).

This list is not complete and there may be other drugs that can interact with Flexeril. Tell your doctor about all your prescription and over-the-counter medications, vitamins, minerals, herbal products, and drugs prescribed by other doctors. Do not start a new medication without telling your doctor.

More Flexeril resources

- [Flexeril Side Effects](#) (in More Detail)
- [Flexeril Dosage](#)
- [Flexeril Use in Pregnancy & Breastfeeding](#)
- [Drug Images](#)
- [Flexeril Drug Interactions](#)
- [Flexeril Support Group](#)
- [67 Reviews for Flexeril](#) - [Add your own review/rating](#)

- Professional [Flexeril Prescribing Information](#) (FDA)
- Advanced [Flexeril Advanced Consumer](#) (Micromedex) - Includes Dosage Information
- Professional [Flexeril Monograph](#) (AHFS DI)
- Med Facts [Flexeril MedFacts Consumer Leaflet](#) (Wolters Kluwer)
- Professional [Cyclobenzaprine Prescribing Information](#) (FDA)
- Professional [Amrix Prescribing Information](#) (FDA)
- Med Facts [Amrix Extended-Release Capsules MedFacts Consumer Leaflet](#) (Wolters Kluwer)

Compare Flexeril with other medications

- [Fibromyalgia](#)
- [Migraine](#)

- [Muscle Spasm](#)
- [Temporomandibular Joint Disorder](#)

- [Sciatica](#)

Where can I get more information?

- Your pharmacist can provide more information about Flexeril.

 [Printable Version](#)

 [Email](#)

[Recommend](#)

 [Tweet](#)

 [Save or Share](#)

Ads by Google

Out Patient Drug Treatment

Change Your Life In A Safe Retreat. Call Us 24/7. Insurance Welcome.
[www.TheTreatmentCenter.com](#)

Male Hormone Treatment

Since 1999 - Hormone treatment-men age 35+ National Network of Doctors
[www.renewman.com](#)

- Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Flexeril only for the indication prescribed.
- Every effort has been made to ensure that the information provided is accurate, up-to-date, and complete, but no guarantee is made to that effect. Drug information contained herein may be time sensitive. Multum's drug information is an informational resource designed to assist licensed healthcare practitioners in caring for their patients and/or to serve consumers viewing this service as a supplement to, and not a substitute for, the expertise, skill, knowledge and judgment of healthcare practitioners. The absence of a warning for a given drug or drug combination in no way should be construed to indicate that the drug or drug combination is safe, effective or appropriate for any given patient. The information contained herein is not intended to cover all possible uses, directions, precautions, warnings, drug interactions, allergic reactions, or adverse effects.

Copyright 1996-2010 Cerner Multum, Inc. Version: 3.11. Revision Date: 4/12/2009 4:38:46 PM.

 enter a search term

Search

Ads by Google [Drug Rehab](#) [Drug Treatment](#) [Drug Clinic](#) [Drug Center](#)

Support	About Drugs.com	Your Privacy
Advertise With Us Frequently Asked Questions Pharmaceutical Companies Widgets for your Website Content Submissions About Drugs.com Contact Us Sitemap Follow us on:  	Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines & natural products. This material is provided for educational purposes only and is not to be used for medical advice, diagnosis or treatment. Data sources include Micromedex™ (updated Feb 3rd, 2011), Cerner Multum™ (updated Jan 20th, 2011), Wolters Kluwer™ (updated Jan 12th, 2011) and others. To view content sources and attributions, refer to our editorial policy. Copyright © 2000-2011 Drugs.com. All rights reserved.	Terms of Use · Privacy Policy  TRUSTe CERTIFIED PRIVACY Verify Here  We comply with the HONcode standard for trustworthy health information. Verify here  Ad Choices