



Search enter a search term

Browse all medications A B C D E F G H I J K L M N O P Q R S T U V W X Y Z Advanced Search Phonetic Search

- Drugs A to Z
- Pill Identifier
- Interactions Checker
- News & Alerts
- Health Professionals
- Community

Mednotes

Drugs A to Z Drugs by Condition Drugs by Class Side Effects Dosage Pregnancy Natural Products Vet Drugs Symptom Checker Harvard Health Topics

Home > Drugs A to Z > I > To > Toradol Consumer Information

Print Save or Share

Toradol

Generic Name: ketorolac (KEE toe ROLE ak)
Brand Names: Toradol

Ads by Google

SUBOXONE® Sublingual Film
SUBOXONE.com (buprenorphine and naloxone) CIII: Now FDA approved: find out more at

Spinal Pain?
www.LaserSpineInstitute.com/Spine Safe Endoscopic Alternative For Your Spine Pain-Get Your Life Back!

Femara® HCP Information
Femara.com/HealthcareProviders Find Patient Support Info on Femara® (letrozole) 2.5 mg tablets.

What is Toradol?

Toradol is in a group of drugs called nonsteroidal anti-inflammatory drugs (NSAIDs). It works by reducing hormones that cause inflammation and pain in the body.

Toradol is used short-term (5 days or less) to treat moderate to severe pain, usually after surgery. It is used alone or in combination with other medicines.

Toradol may also be used for purposes other than those listed in this medication guide.

Important information about Toradol

Toradol can increase your risk of life-threatening heart or circulation problems, including heart attack or stroke. This risk will increase the longer you use Toradol. Do not use this medicine just before or after having heart bypass surgery (also called coronary artery bypass graft, or CABG).

Seek emergency medical help if you have symptoms of heart or circulation problems, such as chest pain, weakness, shortness of breath, slurred speech, or problems with vision or balance.

Toradol can also increase your risk of serious effects on the stomach or intestines, including bleeding or perforation (forming of a hole). These conditions can be fatal and gastrointestinal effects can occur without warning at any time while you are taking Toradol. Older adults may

Sign In or Register Forgot password?

username password Sign In

Related Toradol Information

Approval History Drug history at FDA

Pregnancy Category Risk cannot be ruled out

Availability Prescription only

CSA Schedule Not a controlled drug

Reviews Average User Rating

45 User Reviews 7.4 Rate it!

- Related Pages
- Detailed Side Effects
 - Drug Interactions
 - Support Group Q & A
 - Pregnancy & Breastfeeding
 - Drug Images

Drug Class

Nonsteroidal anti-inflammatory agents

Related Drugs

Pain

- tramadol
- oxycodone
- naproxen
- acetaminophen
- Tylenol
- Percocet
- Vicodin
- amitriptyline
- OxyContin
- clonidine
- aspirin
- More...

Postoperative Pain

- fentanyl
- ketorolac
- Toradol IV/IM
- Toradol IM
- Sublimaze
- More...

have an even greater risk of these serious gastrointestinal side effects.

Call your doctor at once if you have symptoms of bleeding in your stomach or intestines. This includes black, bloody, or tarry stools, or coughing up blood or vomit that looks like coffee grounds.

Do not drink alcohol while taking Toradol. Alcohol can increase the risk of stomach bleeding caused by ketorolac.

Before taking Toradol

Do not use Toradol if you are allergic to ketorolac, aspirin, or other NSAIDs, or if you have:

- severe kidney disease;
- a bleeding or blood clotting disorder;
- a closed head injury or bleeding in your brain;
- a stomach [ulcer](#) or a history of stomach or intestinal bleeding; or
- if you are breast-feeding a baby.

Do not take Toradol if you are also taking pentoxifylline (Trental) or probenecid (Benemid). Do not take this medicine with aspirin or other NSAIDs such as ibuprofen (Motrin, Advil), naproxen (Aleve, Naprosyn), diclofenac (Voltaren), diflunisal (Dolobid), etodolac (Lodine), flurbiprofen (Ansaïd), indomethacin (Indocin), ketoprofen (Orudis), mefenamic acid (Ponstel), meloxicam (Mobic), nabumetone (Relafen), or piroxicam (Feldene).

Taking an NSAID can increase your risk of life-threatening heart or circulation problems, including heart attack or stroke. This risk will increase the longer you use an NSAID. Do not use Toradol just before or after having heart bypass surgery (also called coronary artery bypass graft, or CABG).

NSAIDs can also increase your risk of serious effects on the stomach or intestines, including bleeding or perforation (forming of a hole). These conditions can be fatal and gastrointestinal effects can occur without warning at any time while you are taking an NSAID. Older adults may have an even greater risk of these serious gastrointestinal side effects.

Before taking Toradol, tell your doctor if you are allergic to any drugs, or if you have:

- a history of heart attack, stroke, or blood clot;
- heart disease, congestive heart failure, high blood pressure;
- liver or kidney disease,
- ulcerative colitis or Crohn's disease;
- asthma;
- polyps in your nose;
- if you have recently had surgery; or
- if you smoke.

If you have any of these conditions, you may need a dose adjustment or special tests to safely take Toradol.

FDA pregnancy category C. This medication may be harmful to an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant during treatment. Taking Toradol during labor can increase the risk of bleeding during childbirth. Do not take this medication during pregnancy unless your doctor has told you to.

This medication can affect fertility (your ability to have children). Do not take Toradol while you are trying to get pregnant.

Toradol can pass into breast milk and may harm a nursing baby. Do not take this medicine without telling your doctor if you are breast-feeding a baby. Do not give this medicine to anyone younger than 18 years old.

Ads by Google

[Lantus® \(insulin glargine](#)

[rDNA origin] injection) Get Info - Lantus Might Be Right For You

www.LANTUS.com

Drug Images



Toradol 10 MG

[View larger images](#)

Get Updates by Email

Email me about Toradol:

News and Warnings related to this drug

Also send the Drugs.com monthly newsletter
FDA Medwatch Alerts for all medications

Get Notified

ADVERTISEMENT

Lower Your Blood Pressure - The All Natural Way

The only non-drug medical device clinically proven to lower blood pressure www.resperate.com

\$350,000 in Life Insurance

Protect Your Family Today. Coverage as low as \$13.04/month. Free Quote Here BestQuoteMarket.com

Weight Loss Medicine

Immediate weight loss - reduces fat absorption - less fat means less pounds www.pharmcom.com

Related News and Articles

Health Tip: NSAIDs Induce Ulcers in Some

7 Nov 2010

-- Peptic ulcers are sores that form inside the stomach or intestines. Taking nonsteroidal anti-inflammatory drugs (NSAIDs) for chronic pain can contribute to ...

Health Tip: Taking NSAIDs

27 Oct 2010

-- Nonsteroidal anti-inflammatory drugs (NSAIDs) are over-the-counter or prescription drugs designed to alleviate pain and reduce inflammation. The American ...

Statins, Painkillers May Upset PSA Test Results

7 Aug 2010

FRIDAY, Aug. 6 -- Some of the most widely prescribed drugs in the United States may skew results of prostate cancer screening tests, possibly causing errors in ...

Common Pain Relievers Raise Heart Risk for Healthy Folks

[shoulder pain relief](#)
Excellence In Ortho Shoulder Care And
Replacement. Visit Ingalls.
[IngallsHealthSystem.org/Shoulder](#)

[Back Pain Treatment](#)
Read the Benefits of Our Minimally Invasive
AccuraScope™ Procedure!
[www.NorthAmericanSpine.com](#)

How should I take Toradol?

Take Toradol exactly as it was prescribed for you. Do not take the medication in larger amounts, or take it for longer than recommended by your doctor. Follow the directions on your prescription label. This medication is not for treating minor aches and pains.

Toradol is usually given first as an injection, and then as an oral (by [mouth](#)) medicine. Toradol injection is given through a needle into a muscle or a vein. Your doctor, nurse, or other healthcare provider will give you this injection.

The tablet should be taken with a full glass of water. Toradol is normally given for 5 days or less, including both the injection and oral forms combined. Long-term use of Toradol can damage your kidneys or cause bleeding. If you need to have any type of surgery, tell the surgeon ahead of time if you have recently used this medication. Store the tablets at room temperature away from moisture and heat.

What happens if I miss a dose?

Since Toradol is taken as needed for pain, you may not be on a dosing schedule. And if you receive Toradol injection in a hospital setting, it is not likely that you will miss a dose.

If you are taking the medication regularly, take the missed dose as soon as you remember. If it is almost time for your next dose, skip the missed dose and take the medicine at your next regularly scheduled time. Do not take extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention if you think you have used too much of this medicine. Overdose symptoms may include nausea, vomiting, stomach pain, drowsiness, black or bloody stools, coughing up blood, urinating less than usual, shallow breathing, and fainting.

What should I avoid while taking Toradol?

Do not use any other over-the-counter cold, allergy, or pain medication without first asking your doctor or pharmacist. Many medicines available over the counter contain aspirin or other medicines similar to ketorolac (such as ibuprofen, ketoprofen, or naproxen). If you take certain products together you may accidentally take too much of this type of medication. Read the label of any other medicine you are using to see if it contains aspirin, ibuprofen, ketoprofen, or naproxen. Do not drink alcohol while taking Toradol. Alcohol can increase the risk of stomach bleeding caused by Toradol.

[Ads by Google](#)

[Save Up to \\$30 a Month](#)

On Eligible Prescriptions for Knee & Hand OA Pain - T&C's Apply
[www.goaheadrubitin.com](#)

Toradol side effects

Get emergency medical help if you have any of these signs of an allergic reaction: hives; difficulty breathing; swelling of your face, lips, tongue, or throat. Stop taking Toradol and seek medical attention or call your doctor at once if you have any of these serious side effects:

8 Jun 2010
Healthy people who take nonsteroidal anti-inflammatory drugs to relieve minor aches and pains may raise their risk of dying from heart-related problems, a ...

[Pain Med May Affect Breast Cancer Recurrence](#)

2 Jun 2010
The painkiller a woman takes after a mastectomy may influence the odds of breast cancer recurrence, a new study suggests. Researchers at the Catholic ...

[View more...](#)

Help and Support

Looking for answers? [Ask a question](#) or go join the [Toradol support group](#) to connect with others who have similar interests.

chest pain, weakness, shortness of breath, slurred speech, problems with vision or balance;

- black, bloody, or tarry stools;
- coughing up blood or vomit that looks like coffee grounds;
- swelling or rapid weight gain;
- urinating less than usual or not at all;
- nausea, stomach pain, low fever, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the [skin](#) or eyes);
- fever, sore throat, and headache with a severe blistering, peeling, and red skin rash;
- the first sign of any [mouth sores](#) or skin rash, no matter how mild;
- pale skin, easy bruising, severe tingling, numbness, pain, muscle weakness; or
- fever, headache, neck stiffness, chills, increased sensitivity to light, purple spots on the skin, and/or seizure (convulsions).

Less serious side effects may include:

- upset stomach, mild nausea or vomiting, diarrhea, constipation;
- mild heartburn, stomach pain, bloating, gas;
- dizziness, headache, drowsiness;
- [sweating](#); or
- ringing in your ears.

This is not a complete list of side effects and others may occur. Tell your doctor about any unusual or bothersome side effect. You may report side effects to FDA at 1-800-FDA-1088.

See also: [Toradol side effects](#) (in more detail)

What other drugs will affect Toradol?

Tell your doctor if you are taking an antidepressant such as citalopram (Celexa), duloxetine (Cymbalta), escitalopram (Lexapro), fluoxetine (Prozac, Sarafem, Symbyax), fluvoxamine (Luvox), paroxetine (Paxil), sertraline (Zoloft), or venlafaxine (Effexor). Taking any of these drugs with Toradol may cause you to bruise or bleed easily.

Before taking Toradol, tell your doctor if you are taking any of the following drugs:

- a blood thinner such as warfarin (Coumadin);
- lithium (Eskalith, Lithobid);
- methotrexate (Rheumatrex, Trexall);
- thiothixene (Navane);
- alprazolam (Xanax);
- diuretics (water pills) such as furosemide (Lasix);
- muscle relaxers;
- steroids (prednisone and others);
- seizure medications such as carbamazepine (Carbatrol, Tegretol) or phenytoin (Dilantin);
- a heart or blood pressure medication such as candesartan (Atacand), eprosartan (Teveten), irbesartan (Avapro, Avalide), losartan (Cozaar, Hyzaar), valsartan (Diovan), telmisartan (Micardis), or olmesartan (Benicar);
- aspirin or other NSAIDs such as etodolac (Lodine), flurbiprofen (Ansaid), indomethacin (Indocin), ketoprofen (Orudis), ketorolac (Toradol), mefenamic acid (Ponstel), meloxicam (Mobic), nabumetone (Relafen), naproxen (Aleve, Naprosyn), piroxicam (Feldene), and others; or
- an ACE inhibitor such as benazepril (Lotensin), captopril (Capoten), fosinopril (Monopril), enalapril (Vasotec), lisinopril (Prinivil, Zestril), ramipril (Altace), and others.

This list is not complete and there may be other drugs that can interact with Toradol. Tell your doctor about all the prescription and over-the-counter medications you use. This includes vitamins, minerals, herbal products, and drugs prescribed by other doctors. Do not start using a new medication without telling your doctor.

More Toradol resources

- [Toradol Side Effects](#) (in more detail)
- [Toradol Use in Pregnancy & Breastfeeding](#)
- [Drug Images](#)
- [Toradol Drug Interactions](#)
- [Toradol Support Group](#)
- [45 Reviews for Toradol](#) - [Add your own review/rating](#)

Professional [Toradol Prescribing Information](#) (FDA)

Med Facts [Toradol MedFacts Consumer Leaflet](#) (Wolters Kluwer)

Advanced [Toradol Advanced Consumer](#) (Micromedex) - Includes Dosage Information

Advanced [ketorolac Nasal Advanced Consumer](#) (Micromedex) - Includes Dosage Information

Professional [Ketorolac Prescribing Information](#) (FDA)

Professional [Ketorolac Tromethamine Monograph](#) (AHFS DI)

Compare Toradol with other medications

- [Pain](#)
- [Postoperative Pain](#)

Where can I get more information?

- Your pharmacist can provide more information about Toradol.

[Printable Version](#)

[Email](#)

[Recommend](#)

[Tweet](#)

[Save or Share](#)

Ads by Google

[#1 Addiction Rehab Center](#)
Adult, Addiction Rehab & Detox All Private Rooms, Insurance OK
[TransformationsTreatment.com](#)

[Lower Spine Pain Relief](#)
866-740-6914 Our Surgeons and Staff Are Leaders in Laser Spine Surgery!
[www.LaserSpineProcedure.com](#)

- Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.
- Every effort has been made to ensure that the information provided by Cerner Multum, Inc. ('Multum') is accurate, up-to-date, and complete, but no guarantee is made to that effect. Drug information contained herein may be time sensitive. Multum information has been compiled for use by healthcare practitioners and consumers in the United States and therefore Multum does not warrant that uses outside of the United States are appropriate, unless specifically indicated otherwise. Multum's drug information does not endorse drugs, diagnose patients or recommend therapy. Multum's drug information is an informational resource designed to assist licensed healthcare practitioners in caring for their patients and/or to serve consumers viewing this service as a supplement to, and not a substitute for, the expertise, skill, knowledge and judgment of healthcare practitioners. The absence of a warning for a given drug or drug combination in no way should be construed to indicate that the drug or drug combination is safe, effective or appropriate for any given patient. Multum does not assume any responsibility for any aspect of healthcare administered with the aid of information Multum provides. The information contained herein is not intended to cover all possible uses, directions, precautions, warnings, drug interactions, allergic reactions, or adverse effects. If you have questions about the drugs you are taking, check with your doctor, nurse or pharmacist.

Copyright 1996-2009 Cerner Multum, Inc. Version: 8.02. Revision Date: 4/12/2009 4:36:47 PM.

enter a search term

Search

Ads by Google [Alcohol Rehab](#) [Alcohol Detox](#) [Side Effects](#) [Ibuprofen Pain](#)

Support	About Drugs.com	Your Privacy
Advertise With Us Frequently Asked Questions Pharmaceutical Companies Widgets for your Website Content Submissions About Drugs.com Contact Us Sitemap Follow us on:  	Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines & natural products. This material is provided for educational purposes only and is not to be used for medical advice, diagnosis or treatment. Data sources include Micromedex™ (updated Apr 2nd, 2011), Cerner Multum™ (updated Mar 18th, 2011), Wolters Kluwer™ (updated Apr 4th, 2011) and others. To view content sources and attributions, refer to our editorial policy. Copyright © 2000-2011 Drugs.com. All rights reserved.	 TRUSTe CERTIFIED PRIVACY Verify Here  HON @ CODE We comply with the HONcode standard for trustworthy health information. Verify here  Ad Choices Terms of Use · Privacy Policy